




A treatise of algebra : in two books.

<https://hdl.handle.net/1874/428367>




Bibl. Utenhov.


Octavo n.º 85.76

2

Jonathan Cuthbertson
Book

Bought by him at Newcastle
upon Tyne, Price Seven Shillings
April 9th Anno Domini 1769.





110

A
T R E A T I S E
O F *em*
A L G E B R A,
In TWO BOOKS.

B O O K I.
CONTAINING,
The Fundamental Principles of this ART.
Together with
All the Practical Rules of OPERATION.

B O O K II.
CONTAINING,
A great VARIETY of PROBLEMS,
In the most important
BRANCHES of the MATHEMATICS.

*Vix quicquam in universa Mathesi ita difficile aut arduum
occurrere posse, quò non inoffenso pede per hanc methodum
penetrare liceat.*

SCHOOT. Pref. to DES CARTES.

L O N D O N :
Printed for J. N O U R S E, in the Strand;
Bookseller in Ordinary to His MAJESTY.

MDCCLXIV.

BRIT. LIBRARY

TREATISE

OF

ALGEBRA

IN TWO BOOKS

BOOK I

CONTAINING

The Fundamental Principles of the Art

All the Rules of the Art

BOOK II

CONTAINING

A Full Variety of PROBLEMS

AND

BRANCHES of the MATHEMATICS

By JOHN WALLIS, M.A. Fellow of the Royal Society, and of the University of Cambridge.
Second Edition, corrected and enlarged.
London, Printed by J. Sturges, in the Strand, 1703.

LONDON

Printed for A. Woodman, in the Strand,
Opposite to the Old Swan.

MDCCIII.

T H E
P R E F A C E.

THE subject of the following book is Algebra, a science of universal use in the Mathematics. Its business and use is to solve difficult problems, to find out rules and theorems in any particular branch of science; to discover the properties of such quantities as are concerned in any subject we have a mind to consider. It properly follows these two fundamental branches, Arithmetic and Geometry, but is vastly superior in nature to both, as it can solve questions quite beyond the reach of either of them.

THIS is an art truly sublime, and of an unlimited extent; for if the conditions of a problem be never so complex, and though the quantities concerned are never so much entangled with one another, yet the Algebraist can find means to dissolve and separate them; or if they be ever so remote, his art can furnish him with methods to bring them together and compare them. It is true, he is often obliged to traverse by many roundabout ways, to get the relation of the quantities concerned; yet by certain rules he can pursue the computation of his problem through all these intricate turnings and windings; and by his skill and sagacity can hunt it through all these labyrinths, till he arrives safely

at the end of the chace, viz. the solution of the problem.

THE extent of this curious art is so great that it has gained the title of Universal Mathematics; and is called by way of eminence, The Great Art; and has been esteemed the very apex of human reason. It is also called Specious Arithmetic, Universal Arithmetic, The Analytic Art, The Art of Resolution and Equation; with a view to some or other of its properties or operations.

THE nature of this excellent art is such, that it may be applied to any subject, provided the principles of that subject, it is applied to, be understood. Its great beauty is, that it deals in generals. For whilst other branches go no farther than their own particular subject, and can only find solutions in particular cases; this art finds out general solutions, general rules, general theorems, and general methods.

THIS noble science has also this peculiar property, that it not only investigates rules in all the other parts of the Mathematics; but by the most subtle art and invention, it finds out its own rules, models them according to any form, and varies them at pleasure, so as to answer any end proposed. It would be in vain to attempt to enumerate all the uses of this admirable art.

By making use of letters instead of numbers, it has one great advantage above arithmetic, viz. that in the several operations of arithmetic, the numbers are lost or swallowed up, and changed into others: but here they are preserved distinct, visible, and unchanged. By which means general rules are drawn from particular solutions, to answer all cases of like nature.

The P R E F A C E.

By help of algebraic characters, geometrical demonstrations are often rendered more short, compendious, and clear. So that by this means we avoid the tediousness of a long verbal process, which otherwise we should necessarily be involved in; and which never fails to darken and obscure the subject.

It is highly probable the ancients made use of some sort of analysis, whereby they found out their noble theories. For it is hardly possible so many fine theorems in Geometry, should be groped out or stumbled on, without some such method. But as it was then only in its infancy, it must have been far short of the perfection we have it in at present.

As to the Reader's qualifications, it is absolutely necessary that he understand Arithmetic and Geometry, as the keys to all the rest. And it is also necessary that he understand the principles of every branch of science, to which he would apply algebraic calculations; otherwise it would be in vain to attempt the solution of any problems therein, by the help of Algebra.

THEN as to the method I have followed, it is this. I have gathered together the most valuable rules and precepts, which lie scattered up and down in all the best books of Algebra; and what was deficient, I have supplied as well as I could. Then I have thrown all these precepts and rules of working, into so many problems; which I have reduced into as short a compass, and expressed in as plain terms as possible, so as they may be clear and intelligible. And the method I have taken I suppose will appear to be very simple and easy, and will readily be apprehended by such people as have found confusion and difficulty in other methods. I believe I have omitted nothing that is fundamental; and if any thing of less moment is passed by, it is either
because

because it is of little use, or is supplied by some other method or rule. And all the rules and problems are in such order, that the easiest appear first, and lead on to the harder, which follow in due course afterwards: these make up the first book. And the second book contains the application of Algebra to all sorts of problems, of which there is great variety, and many of them perfectly new; others that are not so, have generally new solutions to them. So I hope I have delivered both the principles and the practice at large, and yet have not clogged the Reader with any superfluity.

W. Emerson.



T H E

THE
CONTENTS.

	Page
D efinitions	1
Characters	3
Notation	5
Axioms	6

BOOK I.

The fundamental Principles.

Sect. I. The Operations in Integers	7
Sect. II. Fractions	31
Sect. III. Surds	48
Sect. IV. Of managing Equations	90
Sect. V. Substitution, Extermination, &c.	110
Sect. VI. Infinite Series	132
Sect. VII. Several fundamental Problems	198
Prob. LXV. To find two quantities whose sum and difference is given	198
Prob. LXVI. To find the least common dividend	198
Prob. LXVII. To find the difference of the squares, having the sum and difference given	201
Prob. LXVIII. Two quantities given, to find the square of the sum	201
Prob. LXIX. Two quantities given to find the square of the difference	202
Prob. LXX. Given the sum and difference to find the rectangle	202
Prob. LXXI. Given the n^{th} power of a binomial, to find the difference between the square of the sum of the odd terms, and the square of the sum of the even terms	203
Prob. LXXII. To find any root of a binomial surd	204
Prob. LXXIII. To explain the properties of ∞ , and infinity	209
Prob. LXXIV. To find the value of a fraction, when both numerator and denominator are ∞	212
Prob. LXXV. To find whole numbers answering the equation $ax=by+c$	215
Prob. LXXVI. To find a number that being divided by given numbers, will leave given remainders	219
Prob	

Prob. LXXVII. To find the limits of an equation containing several unknown quantities	pag. 225
Prob. LXXVIII. To find the limits in two such equations	228
Prob. LXXIX. The investigation of the rule of alligation	231
Prob. LXXX. Investigation of the rule of false	234
Prob. LXXXI. Investigation of the rule of exchange	236
Prob. LXXXII. To find rational squares, cubes, &c.	237
Prob. LXXXIII. To find the maxima and minima of quantities	241
Prob. LXXXIV. To turn numbers into logarithmic series	244
Prob. LXXXV. To turn logarithms into numerical series	247
Prob. LXXXVI. To demonstrate a proposition synthetically, from the analytical solution	250
Sect. VIII. The resolution of equations, and extraction of their roots in numbers	251
Sect. IX. The geometrical construction of equations	297
Sect. X. To investigate a problem algebraically	316

B O O K II.

The Solution of particular Problems.

	Page
Sect. I. Numerical Problems	328
Sect. II. Problems concerning interest and annuities	348
Sect. III. Problems in arithmetical and geometrical progression	358
Sect. IV. Unlimited problems	368
Sect. V. Problems for finding rational squares, cubes, &c.	375
Sect. VI. Geometrical problems	385
Sect. VII. Problems in Plain Trigonometry	419
Sect. VIII. Problems in Spherical Trigonometry	434
Sect. IX. Problems of the Loci	450
Sect. X. Mechanical problems	464
Sect. XI. Physical Problems	475
Sect. XII. Problems concerning series	489
Sect. XIII. Problems concerning exponential quantities	497
Sect. XIV. Problems of Maxima and Minima	507

ALGEBRA.

DEFINITIONS.

1. *ALGEBRA* is a general method of computing Problems, by help of the letters of the alphabet, and other characters. It is of the same nature as Arithmetic, but more general, and therefore it is called *Universal Arithmetic*, as likewise the *Analytic Art*. The peculiar practice of this method is, to assume the quantity sought as if it was known, and proceeding to work by the rules of this art, till at last the quantity sought, or some powers thereof, is found equal to some given quantity, and consequently itself becomes known.
2. *Like quantities*, are those that consist of the same letters; as a , $4a$, $-3a$. Also bb , $3bb$, $-11bb$; also $2abc$, $15abc$, $-abc$; &c.
3. *Unlike quantities*, are those consisting of different letters, or of the same letters, differently repeated. As a , b , $2c$, $-3d$. Also a , $2aa$, $-5aaa$.
4. *Given quantities*, are those whose values are known.
5. *Unknown quantities*, are those whose values are not known.
6. *Simple quantities*, are those consisting of one term only; as $5b$, $3a^2c$, $13dec$, &c.
7. *Compound quantities*, are those consisting of several terms, as $a+b$, $2a-3c$, $a+7b-3d$, &c.

8. *Positive quantities*, are those to be added.

9. *Negative quantities*, are those to be subtracted.

10. *Like signs*, are either all +, or all —, (See the Characters.)

11. *Unlike signs* are + and —.

12. *The Coefficient*, is the number prefixed to any letter or letters in any term. As 3 is the coefficient of $3aa$. If no number be prefixed, then 1 must be understood, as a signifies $1aa$.

13. *A Binomial quantity*, is one consisting of two terms, as $2a + 3b$. *A Trinomial* of 3 terms, as $a + b - c$. *A Quadrinomial* of four, &c. *A Residual* is a binomial, where one of the quantities is negative.

14. *Power* of a quantity, is its square cube, biquadrate, &c.

15. *An Equation*, is the mutual comparing of one thing with another, by the sign of equality put between them.

16. *A dependent Equation*, is an equation which may be deduced from some others.

17. *An independent Equation*, is one that cannot, by any means, be produced from the others.

18. *Pure Equation*, is an equation containing but one power of the unknown quantity; as *a simple Equation*, a *pure Quadratic*, a *pure Cubic*, &c.

19. *An affected Equation*, is that which contains several powers of the unknown quantity; and is denominated according to the highest power in it; as an *affected Quadratic*; an *affected Cubic*; an *affected fourth Power*, &c. Thus a simple equation contains only the simple quantity itself. A quadratic, a quantity of 2 dimensions; a cubic, a quantity of 3 dimensions; a biquadratic, of 4 dimensions, &c.

20. *Index* or *Exponent*, is the number set over a letter shewing what power it is: as a^3 ; here 3 shews

shews it is the third power; or that a^3 is equivalent to aaa . And thus a^4 is the same as $aaaa$; a^5 the same as $aaaaa$, &c. the index always shewing how oft the letter is repeated.

21. *A Fraction*, consists of two quantities placed one above another, with a line between them, as $\frac{a}{b}$; the upper (a) is called the *numerator*, the lower (b) the *denominator*.

22. *A Surd*, is a quantity that has not a proper root, as square root of a (\sqrt{a}), cube root of bb ($\sqrt[3]{bb}$), &c. roots of compound quantities that contain other surds are called, *Universal Surds*.

23. *A rational quantity*, is a quantity that has no radical sign.

Characters used in Algebra.

- + *more, to be added*, being the sign of addition. This is called an affirmative sign. Thus $a + b$ signifies b added to a .
- *less, abating*, the sign of subtraction. This is also called a negative sign. Thus $a - b$, signifies b subtracted from a .

These signs always affect the quantity following; and are always to be interpreted in a contrary signification. If $+$ signifies *upward, forward, gain, increase, above, before, addition*, &c. then $-$ is to be interpreted *downward, backward, loss, decrease, below, behind, subtraction*, &c. And if $+$ be understood of these, then $-$ is to be interpreted of the contrary.

CHARACTERS.

- \cup *difference*; as $a \cup b$, signifies the difference between a and b .
 \times *multiplied by*; as $a \times b$, signifies a multiplied by b . Likewise ab , signifies a multiplied by b . All letters joined together signifies a multiplication. For brevity's sake points are often used instead of \times , as $n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}$, signifies $n \times \frac{n-1}{2} \times \frac{n-2}{3}$.
 \div *divided by*, as $a \div b$, signifies a divided by b , and $\frac{a}{b}$ signifies the same.
 $=$ *equal to*, as $a + b = 2d$, signifies a and b equal to $2d$.
 \sqsupset *greater than*, as $a \sqsupset b$, is a greater than b .
 \sqsubset *lesser than*, as $a \sqsubset b$, is a less than b .
 $\sqrt{\quad}$ *a root*, as \sqrt{a} , is square root of a . $\sqrt[3]{a}$, cube root of a . $\sqrt[4]{a}$, fourth root of a , &c. it is called a *Radical Sign*.
 \odot *involved to*, as $\odot 2$, involved to the square; $\odot 3$ involved to the cube, &c.
 ω *extracted*. $\omega 2$, square root. $\omega 3$, cube root, &c.

$\overline{a+b+c}$, *a line*, or *vinculum*, drawn over several quantities a , b , c , denotes them to be esteemed a compound quantity.

EXPLANATION.

$aa - bb + 3cd$, signifies bb subtracted from aa , and $3cd$ added.

$\overline{aa - bb - cd - dd}$, signifies, that $cc - dd$ is subtracted from $aa - bb$.

$$\overline{aa + 2ab}$$

EXPLANATION.

5

$\overline{aa + 2ab} \in \overline{rr - ss}$, signifies the difference between $aa + 2ab$ and $rr - ss$.

\overline{abcc} signifies the product of a and b and cc .

$\overline{a + b} \times aa$, signifies the sum of $a + b$ multiplied by aa .

$a + \overline{b} \times aa$, signifies the product of b into aa is to be added to a .

$\overline{aa - 2ab}^2$, signifies the square of the compound quantity $aa - 2ab$.

$\sqrt[3]{bb + cc}$ signifies the square root of $bb + cc$.

$\sqrt[3]{2ab - cc}$, signifies the cube root of $2ab - cc$.

$\frac{aa}{a - b}$, signifies aa divided by $a - b$.

$\sqrt{\frac{a^3}{xx - aa}}$ signifies the square root of a^3 divided by $xx - aa$.

a^3b^2 signifies $aaa \times bb$, or the cube of a multiplied by the square of b .

$3ax - xx \sqrt{5ax}$, signifies the square root of $5ax$ multiplied by $3ax - xx$; and so of others.

Quantities that have no sign prefixed, must be understood to have the sign $+$; leading quantities seldom have the signs put down, when they are affirmative.

If AB and CD be two lines; then $AB \times CD$, in a geometrical sense, signifies the rectangle made by the lines AB and CD .

Also $\frac{AB}{CD}$, signifies the ratio that AB has to CD .

NOTATION.

1. In the computation of problems, put the first letters of the alphabet, $b, c, d, f, g, h, \&c.$ for known quantities, and the last letters of the alphabet for unknown ones. Yet some put vowels for

unknown quantities, and the rest of the alphabet for known ones.

2. For general forms, put the capitals A, B, C, D, &c. for the general quantities.

3. Or in universal forms, let the quantities be denoted by the Greek capitals, $\Gamma, \Delta, Z, \Theta, \Lambda, \Pi, \Sigma, \Upsilon, \Phi, \Psi, \Omega$, and indices, coefficients, &c. by the small letters, $\delta, \epsilon, \zeta, \theta, \lambda, \mu, \nu, \pi, \tau, \phi$.

4. In case of necessity, make use of any other sort of letters, or of any characters, that have names, as $h, u, s, o, q, p, v, e, n, f, x$, &c.

A X I O M S.

1. If equal quantities be added to equal quantities, the sums will be equal.

2. If equal quantities be taken from equal quantities, the remainders will be equal.

3. If equal quantities be multiplied by equal quantities, the products will be equal.

4. If equal quantities be divided by equal quantities, the quotients will be equal.

5. The equal powers or roots of equal quantities, are equal.

6. If to or from equal quantities, unequal ones be added or subtracted; the sums or remainders will be unequal.

7. If equal quantities be multiplied or divided by unequal quantities; the products or quotients will be unequal.

8. Quantities severally equal to a third, are equal to one another.

9. The whole is equal to all the parts taken together.

10. If a quantity be added, and the same quantity subtracted, they destroy one another, and are both reduced to nothing.

B O O K I.

The fundamental Principles of Algebra.

S E C T. I.

The primary Operations of Algebra in Integers.

P R O B L E M I.

To add several Quantities together.

1 R U L E.

IF the quantities are like and have like signs ; add all the coefficients together, for the coefficient to that quantity, and prefix the same sign.

Ex. 1.

to + 5a	to -16ab	to + 4a - 3x
add + 7a	add - 5ab	add + 5a - x
Sum +12a	add - 2ab	add + a - 5x
	-23ab	Sum +10a - 9x

Ex. 2.

to + 135ab ²	- 202xxy ³
add + 17abb	- 105xxy ³
+ 3abb	- 17xxy ³
+ abb	- 324xxy ³
Sum + 155abb	

2 R U L E.

If like quantities with unlike signs ; add all the affirmative coefficients, into one sum ; and all the

negative ones into another; subtract the lesser sum from the greater, and to the difference prefix the sign of the greater, with the proper quantity.

Ex. 3.

$$\begin{array}{r}
 \text{to } + 6a \\
 \text{add } - 3a \\
 \hline
 \text{Sum } + 3a
 \end{array}
 \qquad
 \begin{array}{r}
 - 16d \\
 + 3d \\
 \hline
 - 13d
 \end{array}
 \qquad
 \begin{array}{r}
 3a - 7b \\
 - 3a + 8b \\
 \hline
 0 + b
 \end{array}$$

Ex. 4.

$$\begin{array}{r}
 - 125ab \\
 - 37ab \\
 + ab \\
 + 99ab \\
 \hline
 - 162ab \\
 + 100ab \\
 \hline
 \text{Sum } - 62ab
 \end{array}
 \qquad
 \begin{array}{r}
 + 34x^2y \\
 - 8x^2y \\
 - x^2y \\
 + 92x^2y \\
 - 67x^2y \\
 \hline
 + 126x^2y \\
 - 76x^2y \\
 \hline
 + 50x^2y
 \end{array}$$

Ex. 5.

$$\begin{array}{r}
 - 2aa - 9bcd + dd + 2e \\
 + 7aa - 20bcd - dd + 5e \\
 + 3aa + 4bcd \\
 \hline
 \hline
 \text{Sum } + 8aa - 25bcd + 7e
 \end{array}$$

3 R U L E .

Set down all the unlike quantities with their proper signs.

Ex. 6.

$$\begin{array}{r}
 + 2a \\
 + 3b \\
 - c \\
 + d \\
 \hline
 \text{Sum } 2a + 3b - c + d
 \end{array}$$

Ex.

Ex. 7.

$$\begin{array}{r}
 + 13aa - 2a \\
 - 4ab = 3a \\
 + bc + 5a \\
 - 2dd + 6d
 \end{array}$$

$$\text{Sum } + \underline{13aa - 4ab + bc - 2dd + 6d.}$$

Ex. 8.

$$\begin{array}{r}
 2ee + 3ef - ff + 17 \\
 - 3ee + 5ef + 2ff - 11 \\
 + 6ee - ef + ff - 3
 \end{array}$$

$$\text{Sum } + \underline{5ee + 7ef + 2ff + 3}$$

The reason of this rule is evident for like signs; and in unlike signs, it follows from the nature of affirmative and negative quantities, that the difference ought to be taken, to make up the total. As if a man owes 10*l.* then 10*l.* ought to be deducted from his stock to find his real worth.

Cor. 1. *When several quantities are to be added together, it is the same thing, in whatever order they are placed.*

Thus $a + b - c = a - c + b = -c + a + b = b + a - c$, &c. for all these are the same.

Cor. 2. *Hence the sum of any number of affirmative quantities, is affirmative; and the sum of any number of negative quantities, is negative.*

P R O B L E M II.

To subtract quantities from one another.

R U L E.

Change the signs of all the quantities to be subtracted; and then add them all together by Prob. I. and their sum will be the remainder sought.

Ex.

Ex. 1.

$$\begin{array}{r}
 + 8a \\
 + 3a \\
 \hline
 \text{Rem. } 8a - 3a \\
 \text{or } 5a
 \end{array}
 \qquad
 \begin{array}{r}
 16b \\
 - 5b \\
 \hline
 16b + 5b \\
 \text{or } 21b
 \end{array}
 \qquad
 \begin{array}{r}
 - 11c \\
 + 3c \\
 \hline
 - 11c - 3c \\
 \text{or } - 14c
 \end{array}$$

— 20

— 6

— 20 + 6

or — 14

Ex. 2.

$$\begin{array}{r}
 \text{from } 6a - 3x + 6y - 7 \\
 \text{take } + 8a + 4x + 6y + 5 \\
 \hline
 \text{Rem. } - 2a - 7x + 0 - 12
 \end{array}$$

Ex. 3.

$$\begin{array}{r}
 \text{from } a + b \qquad a + b \\
 \text{take } a - b \qquad - a + b \\
 \hline
 \text{Rem. } + 2b \qquad 2a
 \end{array}$$

Ex. 4.

$$\begin{array}{r}
 \text{from } aa + 2ab + bb \\
 \text{take } + 4ab \\
 \hline
 \text{Rem. } aa - 2ab + bb
 \end{array}$$

Ex. 5.

$$\begin{array}{r}
 \text{from } aa - bb \\
 \text{take } cc - dd \\
 \hline
 \text{Rem. } aa - bb - cc + dd
 \end{array}$$

Ex. 6.

$$\begin{array}{r}
 \text{from } 3aa - 2a + cd - dd - ff \\
 \text{take } - 2aa - 5a - ab - 2dd \\
 \hline
 \text{Rem. } 5aa + 3a + cd + ab + dd - ff
 \end{array}$$

Cor.

Sect. I. MULTIPLICATION. 11

Cor. 1. Hence, To subtract one quantity from another, is the same thing as to add them together, when all the signs of the subtrahend are changed.

$$a - - b = a + b.$$

For it is the same thing to subtract $-$, as to add $+$; and to add $-$, as to subtract $+$. For suppose a man to owe 10 l; because it is a debt it must be writ -10 l. therefore if any body would take away this -10 , it is the same thing as if he added $+10$ to his stock: but before it is discharged, this -10 is the same, as $+10$ deducted out of his stock.

PROBLEM III.

To multiply one quantity by another.

R U L E.

Multiply every particular term (or simple quantity) of the multiplier, into every term of the multiplicand, one after another; so that the coefficients be multiplied into the coefficients; and the letters into the letters, by placing them all together, like letters in a word. And prefix $+$ to products of like signs, and $-$ to unlike ones. The sum of all is the product sought.

Ex. 1.

$+ a$	$- a$	$+ 3 a$	$- 4 c$
$+ b$	$- b$	$- 2 b$	$+ 5 d$
$+ a b$	$+ a b$	$- 6 a b$	$- 20 c d$

Ex. 2.

$a + b$	$a + b$	
$a + b$	$a - b$	
$a a + a b$	$a a + a b$	
$+ a b + b b$	$- a b - b b$	
$a a + 2 a b + b b$	$a a - b b$	Ex.

Ex. 3.

$$\begin{array}{r}
 3a - 2b \\
 5a + 4b \\
 \hline
 15aa - 10ab \\
 \quad + 12ab - 8bb \\
 \hline
 15aa + 2ab - 8bb
 \end{array}$$

Ex. 4.

$$\begin{array}{r}
 aa + ab - bb \\
 a - b \\
 \hline
 a^3 + aab - abb \\
 \quad - aab - abb + b^3 \\
 \hline
 a^3 \quad \quad - 2abb + b^3
 \end{array}$$

Ex. 5.

$$\begin{array}{r}
 ab - 3cd + rs \\
 5r - 7d \\
 \hline
 5rab - 15rcd + 5rrs - 7abd + 2icdd - 7rsd.
 \end{array}$$

Ex. 6.

$$\begin{array}{r}
 3aa - 2ab + 5 \\
 aa + 2ab - 3 \\
 \hline
 3a^4 - 2ba^3 + 5aa \\
 \quad + 6ba^3 - 4aabb + 10ab \\
 \quad \quad - 9aa - 6ab - 15 \\
 \hline
 3a^4 + 4ba^3 - 4baa - 4aa + 16ab - 15
 \end{array}$$

Ex. 7.

$$\begin{array}{r}
 aa + bb \\
 cc - dd \\
 \hline
 ccaa + ccbb \\
 \quad - ddaa - ddbb \\
 \hline
 ccaa - ddaa + ccbb - ddbb
 \end{array}$$

Ex.

Ex. 8.

$$\begin{array}{r}
 a^3 \\
 a^2 \\
 \hline
 a^5
 \end{array}
 \qquad
 \begin{array}{r}
 4 b^3 \\
 - 3 b^5 \\
 \hline
 - 12 b^8
 \end{array}$$

That every term in the multiplicand must be multiplied by every term in the multiplier, is thus made evident. Let $a + b$ be multiplied by $c + d$; it is plain, $a + b$ must be taken so often as there are supposed units in c and d , that is, as often as there are units in c , and also as oft as there are units in d . Therefore the product will be $a + b \times c + a + b \times d$. But for the same reason $a + b \times c = ac + bc$, also $a + b \times d = ad + bd$. Whence the product will be $ac + bc + ad + bd$; that is, the sum of all the products of every term multiplied by every term.

That like signs give $+$, and unlike signs $-$, in the product, will appear thus.

Case 1. Let $+a$ be multiplied by $+b$. Then since this multiplication supposes, that $+a$ is to be so often added together as there are units in $+b$; and the sum of any number of affirmatives is affirmative, therefore the whole sum is affirmative, that is $+a \times +b = +ab$.

Case 2. Let $+a$ be multiplied by $-b$. Now since this implies that $+a$ is to be as often subtracted as there are units in b ; and the sum of any number of negatives, is negative, therefore that whole sum, is negative, that is, $+a \times -b = -ab$.

Case 3. Let $-a$ be multiplied by $+b$. It is plain here, that $-a$ is to be so often taken as there are units in b ; and the sum of any number of negatives being negative, therefore the whole sum is negative; that is, $-a \times +b = -ab$.

Otherwise,

Otherwise, Let $d - a$ be multiplied by $+ b$; then (*Case 1.*) the product will be bd together with $- a \times + b$: but bd is too big, as being the product of d by b ; instead of $d - a$ by b ($d - a$ being less than d); therefore bd , being too much, the product $- a \times + b$ must be subtracted; that is, the true product will be $db - ab$; and consequently $- a b = - a \times + b$.

Case 4. Let $- a$ be multiplied by $- b$. Here $- a$ is to be subtracted as often as there are units in b : but subtracting negatives is the same as adding affirmatives (*Cor. 2. Prob. 2.*); consequently the product is $+ a b$.

Or thus. Since $a - a = 0$, therefore $\overline{a - a} \times - b = 0$, because 0 multiplied by any thing produces 0; therefore since $+ a - a \times - b = 0$; and the first term of the product is $- ab$ (*Case 2.*); therefore the last term of the product must be $+ ab$, to make the sum 0, or $- ab + ab = 0$; that is, $- a \times - b = + a b$.

Otherwise, Let $d - a$ be multiplied by $- b$. Then (*Case 2.*) the product will be $- bd$ together with $- a \times - b$; but $- bd$ the quantity to be subtracted is too big, being the product of d by $- b$, instead of $d - a$ by $- b$, ($d - a$ being less than d); therefore the quantity $- bd$ to be subtracted being too much, something must be restored, that is $- a \times - b$ must be added; and the true product will be $- bd + ab$; and therefore $+ ab = - a \times - b$.

Cor. 1. If several quantities are to be multiplied together; it is the same thing in whatever order it be done. Thus $abc = acb = cab = bca$, &c. for all these are equal.

Cor. 2. The powers of the same quantity are multiplied together, by adding their indices. Thus $a^2 \times a^3 = a^{2+3} = a^5$.

Cor. 3. Any odd number of —, multiplied together produce —; and any even number of —, produce +.

S C H O L I U M.

In the multiplication of compound quantities, it is the best way to set them down in order, according to the dimensions of some of the quantities. And in multiplying them, begin at the left hand, and multiply from the left hand towards the right, the way we write, which is contrary to the way we multiply numbers. But this will be most expeditious, and the several products will by this means be so ranged under one another, that like quantities will fall in the same places, which is the easiest way for adding them up together.

In many cases, the multiplication of compound quantities is only to be performed by writing their sums, each under a vinculum, and putting the sign (\times) of multiplication between. As if the square of $aa - xx$ was to be multiplied by $ag - bb$, and that by $ac + bd$, it may be written thus, $\overline{aa - xx} \times \overline{ag - bb} \times \overline{ac + bd}$.

P R O B L E M IV.

To divide one quantity by another.

R U L E.

In simple quantities, which will divide without a remainder; divide the number by the number, and put the answer in the quotient. Then throw out all the letters in the dividend which are found in the divisor, and place the remaining letters in the quotient. And like signs produce +, and unlike signs —, in the quotient.

Ex.

Ex. 1.

$$\begin{array}{r} aa) aab (b \\ \underline{aab} \\ 0 \end{array}$$

$$\begin{array}{r} 3ab) 15abcd (5acd \\ \underline{15abcd} \\ 0 \end{array}$$

Ex. 2.

$$\begin{array}{r} -aa) aab (-b \\ \underline{aab} \\ 0 \end{array}$$

$$\begin{array}{r} -3ab) -15abcd (5acd \\ \underline{-15abcd} \\ 0 \end{array}$$

Ex. 3.

$$\begin{array}{r} -3cdd) -6cddd (+2c \\ \underline{-6cddd} \\ 0 \end{array}$$

Ex. 4.

$$\begin{array}{r} +6a^2b^2) -18b^3a^3d^3c (-3abcd^3 \\ \underline{-18b^3a^3d^3c} \\ 0 \end{array}$$

Ex. 5.

$$\begin{array}{r} -5a^2b) 10a^2bbd (-2bd \\ \underline{10a^2bbd} \\ 0 \end{array}$$

Ex. 6.

$$\begin{array}{r} 9x^2y) -9x^2y^2b (-yb \\ \underline{-9x^2y^2b} \\ 0 \end{array}$$

Ex.

$$\begin{array}{r}
 \text{Ex. 7.} \\
 - 8 x x) - 16 x^3 (+ 2 x \\
 \underline{- 16 x^3} \\
 \hline
 0
 \end{array}$$

2 R U L E .

In compound quantities, range the terms of the divisor and dividend, according to the dimensions of some letter. Then, by Rule 1, divide the first term of the dividend by the first term of the divisor, placing the result in the quotient. Multiply the whole divisor by the quotient, and subtract it from the dividend, to which bring down the next term of the dividend, call this the *Dividual*.

Divide the first term of the dividual by the first term of the divisor; then multiply and subtract as before, and repeat the same process till all the quantities be brought down. This is in effect the very same rule as is used in arithmetic.

Ex. 3.

$$\begin{array}{r}
 a) ab + ac - a(b + c - 1 \text{ the quotient} \\
 \underline{ab} \\
 + ac \\
 + ac \\
 \hline
 - a \\
 - a \\
 \hline
 0
 \end{array}$$

Ex. 9.

$$\begin{array}{r}
 2b - 3c) 2baa - 3caa (aa \\
 \underline{2baa - 3caa} \\
 \hline

 \end{array}$$

C

Ex.

form of a fraction; throwing out such letters, as are found in all the terms of both the dividend and divisor.

Ex. 14.

$$a-x) a \left(\frac{a}{a-x} \text{ the quotient.} \right.$$

Ex. 15.

$$ax-xx) ax+xx \left(\frac{ax+xx}{ax-xx} = \frac{a+x}{a-x} \text{ quote.} \right.$$

Ex. 16.

$$1-x) 1 \quad (1+x+xx+x^3+x^4 + \&c. \text{ sine sine.}$$

$$\begin{array}{r} 1-x \\ \hline +x \\ \hline x-xx \quad \left(\text{or } 1+x+xx+x^3 + \frac{x^4}{1-x} \right. \\ \hline +xx \\ \hline xx-x^3 \\ \hline +x^3 \\ \hline x^3-x^4 \\ \hline +x^4, \&c. \end{array}$$

Ex. 17.

$$aa-ee) aae \quad \left(e + \frac{e^3}{aa} + \frac{e^5}{a^4} + \frac{e^7}{a^6-eea^4} \right.$$

$$\begin{array}{r} aae-e^3 \\ \hline +e^3 \\ \hline e^3 - \frac{e^5}{aa} \\ \hline +\frac{e^5}{aa} \\ \hline +\frac{e^5}{aa} - \frac{e^7}{a^4} \\ \hline +\frac{e^7}{a^4} \text{ Rem.} \end{array}$$

This and such like examples will be better understood after the next section.

Ex. 18.

$$a^3) 2a^5 \left(\frac{2a^5}{a^3} = 2a^2. \right.$$

Ex. 19.

$$-2b^2) 18b^6 \left(-\frac{18b^6}{b^2} = -9b^4. \right.$$

Ex. 20.

$$\begin{array}{r} \overline{aa-xx^2) \overline{aa-xx^2} \left(\overline{aa-xx^2} \right. \\ \underline{aa-xx^2} \\ 0 \end{array}$$

That like signs give +, and unlike signs —, in the quotient, will appear thus. The divisor multiplied by the quotient must produce the dividend. Therefore, 1. When both are +, the quo-

1. +) + (+ must produce + in the dividend.
 2. —) — (— must produce — in the dividend.
 3. +) — (— must produce + in the dividend.
 4. —) + (+ must produce — in the dividend.)

Again, 3. When the divisor is + and the dividend —, the quotient is —, because — × + must produce — in the dividend. 4. Lastly, If the divisor is —, and the dividend +, the quotient will be —, because — × — produces + in the dividend.

Cor. 1. One power of a quantity, is divided by another power thereof; by subtracting the index of the divisor, from the index of the dividend. Thus,

$$\frac{a^5}{a^3} = a^{5-3} = a^2. \quad \text{And} \quad \frac{16b^3}{12b^3} = \frac{4b^{3-3}}{3} = \frac{4b^{-2}}{3} = \frac{4}{3bb}.$$

Cor.,

Cor. 2. Hence any power of a quantity may be taken out of the denominator and put into the numerator, and the contrary; by changing the sign of the index.

Thus $\frac{a}{2b^2} = \frac{ab^{-2}}{2}$. And $\frac{b}{a^{-3}} = ba^3$.

Cor. 3. Hence — divided by +, or + divided by —, give the same quotient, viz. —, That is,

$\frac{-a}{b} = \frac{a}{-b} = -\frac{a}{b}$.

PROBLEM V.

To involve a quantity to any power.

RULE.

Multiply the quantity so often into itself as the index denotes. And where the root is +, all the powers are +. And where the root is —, all the odd powers are —, and all the even powers +.

Ex. 1.

a root	—	a ² root
a a square	—	a ⁴ square
a ³ cube	—	a ⁶ cube
a ⁴ 4th power	—	a ⁸ 4th power
&c.		&c.

— 2a ³ root
+ 4a ⁶ square
— 8a ⁹ cube
+ 16a ¹² 4th power.
&c.

Ex. 2.

a b root	—	3 a b b root
aabb square	—	+ 9 a a b ⁴ square
a ³ b ³ cube	—	— 27 a ³ b ³ cube,
&c.		&c.

Ex. 3.

Involve $a+b$ to the cube or 3d power.

$$\begin{array}{r}
 a + b \\
 a + b \\
 \hline
 aa + ab \\
 \quad + ab + bb \\
 \hline
 \text{square } aa + 2ab + bb \\
 a + b \\
 \hline
 a^3 + 2aab + abb \\
 \quad + aab + 2abb + b^3 \\
 \hline
 \text{cube } a^3 + 3aab + 3abb + b^3
 \end{array}$$

2 R U L E.

Multiply the index of the quantity, by the index of the power, and make the signs as in Rule 1.

Ex. 4.

$$\begin{array}{ll}
 \text{root } a \text{ or } a^1 & - 2 b b a; \text{ or } 2 b^2 a \\
 \text{square } a^{1 \times 2} \text{ or } a^2 & + 4 b^{2 \times 2} a^{1 \times 2} \text{ or } + 4 b^4 a^2 \\
 \text{cube } a^{1 \times 3} \text{ or } a^3 & - 8 b^{2 \times 3} a^{1 \times 3} \text{ or } - 8 b^6 a^3 \\
 \text{th} & \\
 m \text{ power } a^m & - 2^m \times b^{2m} a^m
 \end{array}$$

Ex. 5.

$$\begin{array}{ll}
 \text{root } \overline{a-x}^2 & \\
 \text{square } \overline{a-x}^{2 \times 2} \text{ or } \overline{a-x}^4 & \\
 \text{cube } \overline{a-x}^{2 \times 3} \text{ or } \overline{a-x}^6 & \\
 \text{th} & \\
 m \text{ power } \overline{a-x}^{2 \times m} \text{ or } \overline{a-x}^{2m} &
 \end{array}$$

3 R U L E.

3 R U L E.

In a binomial. The power will consist of 1 term more than the index of the power. The highest power of both is the index of the given power, and the index of the leading quantity continually decreases by 1 in every term, and in the following quantity, the indices of the terms are 0, 1, 2, 3, 4, &c.

Then for finding the unciæ or coefficients. The first is always 1; the second, the index of the power. And in general, if the coefficient of any term be multiplied by the index of the leading quantity, and divided by the number of terms to that place; it gives the coefficient of the next following term.

Lastly, When both terms of the root are +, all the terms of the power will be +; but if the second term be —, then all the odd terms will be +, and all the even terms —.

Ex. 6.

Involve a+e to the 5th power.

The several terms without the coefficients will be

$$a^5, a^4e, a^3ee, a^2e^3, ae^4, e^5; \text{ and the}$$

$$\text{coefficients } 1, 5, \frac{5 \times 4}{2}, \frac{10 \times 3}{3}, \frac{10 \times 2}{4}, \frac{5 \times 1}{5};$$

$$\text{that is, } 1, 5, 10, 10, 5, 1.$$

And therefore the 5th power is

$$a^5 + 5 a^4e + 10 a^3ee + 10 a^2e^3 + 5 ae^4 + e^5.$$

Ex. 7.

Involve a-x to the 4th power.

$$\text{the root is } a^4 - 4a^3x + \frac{4 \times 3}{2} a^2x^2 - \frac{6 \times 2}{3} ax^3 + \frac{4 \times 1}{4} x^4;$$

$$\text{that is, } a^4 - 4a^3x + 6a^2x^2 - 4ax^3 + x^4.$$

C 4

4 RULE.

4 R U L E.

In trinomials, quadrinomials, &c. Let one letter remain, and put another letter for the rest of the quantities; then involve this binomial by Rule 3; then instead of the powers of the assumed letter, find (by Rule 3.) the powers of the compound quantity it represents, which put in its stead.

Ex. 8.

Involve $a+b-x$ to the third power.

Put e for $b-x$, then the cube of $a+e$ is $a^3 + 3a^2e + 3aee + e^3$ (Rule 3), that is, $a^3 + 3a^2 \times \overline{b-x} + 3a \times \overline{b-x}^2 + \overline{b-x}^3$. But (Rule 3.) $\overline{b-x}^2 = \overline{bb-2bx+xx}$, and $\overline{b-x}^3 = \overline{b^3-3bbx+3bx^2-x^3}$. Therefore $\overline{a+b-x}^3 = a^3 + 3a^2b - 3a^2x + 3abb - 6abx + 3axx + b^3 - 3bbx + 3bx^2 - x^3$.

Cor. 1. *The n^{th} power of $a+e$, that is,*

$$\overline{a+e}^n = a^n + na^{n-1}e + n \times \frac{n-1}{2} a^{n-2}ee + n \times \frac{n-1}{2} \times \frac{n-2}{3} a^{n-3}e^2 + n \times \frac{n-1}{2} \times \frac{n-2}{3} \times \frac{n-3}{4} a^{n-4}e^3 + \&c.$$

This rule is proved by involving $a+e$ as far as you will, for the several powers will always agree with the rule.

Cor. 2. *All powers of an affirmative quantity, are affirmative. And all odd powers of a negative quantity, are negative; and all even powers affirmative.*

Cor. 3. *The index of the power of any quantity, is the product of the index of the power, and index of the quantity.*

Cor. 4. *The n^{th} power of any product, is equal to the n^{th} power of each factor, multiplied together.*

$$\overline{ab}^n = a^n \times b^n.$$

PROBLEM VI.

To extract the root of any quantity.

Evolution is just the reverse to involution; and is performed as follows.

1 RULE.

For simple quantities; extract the root of the coefficient for the numerical part, and divide the index of the letter or letters, by the index of the power, gives the index of the root.

Ex. 1.

The cube root of a^3 is a^1 or a .

the square root of $25a^4$ is $5a^2$ or $5aa$.

the square root of $2a^2b^6$ is $a^1b^3\sqrt{2}$
or $ab^3\sqrt{2}$.

the cube root of $-125b^9$ is $-5b^3$ or $-5b^3$.

2 RULE.

For the square root, of a compound quantity; range the terms according to the dimensions of some letter. Then find the root of the first term (1 Rule), and set it in the quotient: subtract its square, and bring down the next term, which divide by double the quotient, and set the answer in the quotient. Multiply the divisor and quotient by this last quotient, which subtract from the dividend, proceed thus, just as in common arithmetic.

Ex.

Ex. 4.

Extract the square root of $aa+xx$.

$$\begin{array}{r}
 aa+xx \left(a + \frac{xx}{2a} - \frac{x^4}{8a^3} + \frac{x^6}{16a^5} - \&c. \right. \\
 \hline
 2a + \frac{xx}{aa} \left. \right) \circ + xx \\
 \quad + xx + \frac{x^4}{4aa} \\
 \hline
 2a + \frac{xx}{a} - \frac{x^4}{8a^3} \left. \right) \circ - \frac{x^4}{4aa} \\
 \quad - \frac{x^4}{4aa} - \frac{x^6}{8a^4} + \frac{x^8}{64a^6} \\
 \hline
 2a + \left. \right) \circ + \frac{x^6}{8a^4} - \frac{x^8}{64a^6}
 \end{array}$$

3 R U L E.

In higher powers. Find the root of the first member, which place in the quotient: subtract its power, the remainder is the residual. Involve this root to the next lower power, and multiply it by the index of the given power, for a divisor; by this divide the first term of the residual, the quotient is the next term of the root. Then involve the whole root as before, and subtract: and repeat the operation, till all the terms of the root be had.

Ex. 5.

Extract the cube root of $x^6+6x^5-40x^3+96x-64$.
 $x^6+6x^5-40x^3+96x-64$ ($xx+2x-4$ root.

$$\begin{array}{r}
 x^6 \\
 \hline
 3x^4 \left. \right) 6x^5 (+2x \\
 \hline
 x^6+6x^5+12x^4+8x^3 = \overline{xx+2x} \\
 \hline
 3x^4 \left. \right) \circ -12x^4 (-4 \\
 \hline
 x^6+6x^5-40x^3+96x-64 = \overline{xx+2x-4} \\
 \hline
 \circ
 \end{array}$$

Ex.

volved, it is $16a^4 + 96a^3x + 216a^2x^2 + 216ax^3 + 81x^4$, which differs in the signs, from the quantity given. Therefore make $2a - 3x$ the root, which being involved succeeds; the power being $16a^4 - 96a^3x + 216a^2x^2 - 216ax^3 + 81x^4$.

5 R U L E.

When the quantity given has not such a root as is required, set it down in form of a surd.

Ex. 9.

Square root of a^2 , is $\sqrt{a^2}$.

Cube root of $15aa$, is $\sqrt[3]{15aa}$.

4th root of $2a^2x^3$, is $\sqrt[4]{2a^2x^3}$.

Ex. 10.

The cube root of $a^3 - 6a^2b + 12abb + 8b^3$, is $\sqrt[3]{a^3 - 6a^2b + 12abb + 8b^3}$.

Ex. 11.

What is the 5th root of $a^5 - x^5$.

the root is $\sqrt[5]{a^5 - x^5}$.

Cor. 1. The square root, or any even root, of an affirmative quantity, may be either $+$ or $-$.

For the square root of aa may be $+a$ or $-a$, for $+a \times +a = aa$, and $-a \times -a = aa$: also the 4th root of a^4 is $+a$ or $-a$, for the 4th power of $-a$ is $+a^4$, as well as of $+a$.

Cor. 2. Any odd root of a quantity, will have the same sign, as the quantity itself.

For the root of $+a^3$ and $-a^3$, will be $+a$ and $-a$; for $+a$ cubed is $+a^3$, and $-a$ cubed is $-a^3$.

Cor.

Cor. 3. *The square root, or any even root, of a negative quantity, is impossible.*

For neither $+a \times +a$, nor $-a \times -a$, can produce $-aa$.

Cor. 4. *The n^{th} root of a product, is equal to the n^{th} root of each of the factors, multiplied together:*

$$\sqrt[n]{AB} = \sqrt[n]{A} \times \sqrt[n]{B}.$$



S E C T. II.

O f F R A C T I O N S.

THE operations of algebraic fractions are exactly the same as those of vulgar fractions in arithmetic; therefore he that has made himself master of vulgar fractions, will easily understand how to manage all sorts of algebraic fractions, as in the following problems.

P R O B L E M VII.

To reduce a given quantity to a fraction of a given denominator.

R U L E.

Multiply that quantity by the given denominator, and under the product write the same denominator.

Ex. 1.

Let $a+b$ have the denominator x .

$$\frac{a+b \times x}{x} = \frac{ax+bx}{x}, \text{ answer.}$$

Ex. 2.

Let $xx-yy$ have the denominator 1.

$$\frac{xx-yy \times 1}{1} = \frac{xx-yy}{1}, \text{ answer.}$$

Ex. 3.

Let $\frac{a}{b}$ have the denominator $b-c$.

$$\frac{\frac{a}{b} \times \overline{b-c}}{b-c} = \frac{a - \frac{ac}{b}}{b-c}, \text{ answer.}$$

Cor.

Cor. The value of a fraction is not altered, by multiplying both numerator and denominator by the same quantity. Thus $\frac{rab}{rc} = \frac{rabd}{rcd} = \frac{ab}{c}$.

PROBLEM VIII.

To reduce a mixed number to a fraction.

R U L E.

Multiply the integral part by the denominator of the fraction, and to the product add the numerator, under which write the common denominator.

Ex. 1.

Let $a - \frac{b}{c}$ be given. Then $\frac{ac-b}{c}$ is the fraction required.

Ex. 2.

Suppose $a - x + \frac{aa - ax}{x}$.

Here $\frac{ax - xx + aa - ax}{x}$, or $\frac{aa - xx}{x}$ is that required.

PROBLEM IX.

To reduce an improper fraction to a whole or mixed number.

R U L E.

Divide the numerator by the denominator, as far as you can, gives the integral part; and place the remainder over the denominator for the fractional part.

Ex.

Ex. 1.

Given $\frac{ab-aa}{b}$. $b)ab-aa$ $\left(a-\frac{aa}{b}\right)$ answer.

-aa

Ex. 2.

Suppose $\frac{aa+xx}{a-x}$.

$a-x)aa+xx$ $\left(a+x+\frac{2xx}{a-x}\right)$, answer.

+ax+xx

+ax-xx

+2xx

PROBLEM X.

To find the greatest common divisor, for the terms of a fraction, or for any two quantities.

R U L E.

The quantities being ranged according to the dimensions of some letter; divide the greater by the lesser, and the last divisor by the last remainder, and so on continually till nothing remain; then the last divisor is that required. But observe, first to throw out of each divisor, all the simple divisors, (or others) that will divide it; and then proceed. The simple divisors are had by inspection.

Ex. 1.

Let $\frac{cd+dd}{aac+aad}$ be the fraction proposed.

$cd+dd)aac+aad$ (

or $c+d)aac+aad$ (aa

$aac+aad$

o

D

There-

PROBLEM XI.

To reduce a fraction to its lowest terms.

R U L E.

Find the greatest common measure (Prob. X), by which divide both numerator and denominator of the fraction; the quotients will be the numerator and denominator of the fraction required.

Ex. 1.

Let $\frac{cc+dd}{aac+aad}$ be proposed.

the greatest common divisor is $c+d$. Therefore

$$c+d) \frac{cd+dd}{aac+aad} = \frac{d}{aa} \text{ the fraction required.}$$

Ex. 2.

Let $\frac{a^3-abb}{aa+2ab+bb}$ be proposed.

Here $a+b$ is the greatest common divisor: then

$$a+b) \frac{a^3-abb}{aa+2ab+bb} = \frac{aa-ab}{a+b} \text{ the fraction sought.}$$

Ex. 3.

Suppose $\frac{a^4-b^4}{a^5-bba^3}$ to be given.

the greatest common divisor is $aa-bb$; then

$$aa-bb) \frac{a^4-b^4}{a^5-bba^3} \left(= \frac{aa+bb}{a^3} \text{ the fraction required.} \right.$$

PROBLEM XII.

To reduce fractions of different denominators, to fractions of the same value, having a common denominator.

I RULE.

Multiply each numerator, into all the other denominators, for a new numerator; then multiply all the denominators together for a common denominator.

Ex. 1.

Let $\frac{a}{b}$, $\frac{a+b}{c}$ be given.

these become $\frac{ac}{bc}$, $\frac{ab+bb}{bc}$.

Ex. 2.

Let $\frac{a}{b}$, $\frac{c}{d}$, $\frac{f}{g}$ be proposed.

they become $\frac{adg}{bdg}$, $\frac{c bg}{bdg}$, $\frac{fbd}{bdg}$

2 RULE.

Divide the denominators by their greatest common divisor, then multiply both numerator and denominator of each fraction, by all the other quotients, which will produce as many new fractions.

Ex. 3.

Suppose $\frac{a}{2bb}$, $\frac{c}{2b}$, $\frac{d}{b}$.

$\frac{2a}{4bb}$, $\frac{2bc}{4bb}$, $\frac{4bd}{4bb}$ the fractions required.

or $\frac{a}{2bb}$, $\frac{bc}{2bb}$, $\frac{2bd}{2bb}$

Ex.

Ex. 4.

$$\text{Given, } \frac{2aa}{aa-ab}, \frac{3ab-2bb}{2ac}.$$

$$\frac{4aac}{2aac-2abc}, \frac{3ab-2bb \times a-b}{2aac-2abc} \quad \text{that is,}$$

$$\frac{4aac}{2aac-2abc}, \frac{3aab-5abb+2b^2}{2aac-2abc}.$$

PROBLEM XIII.

To add fractional quantities together.

R U L E.

If the fractions have not a common denominator, reduce them to one (Prob. XII); then add the numerators, and under the sum, write the common denominator.

Ex. 1.

$$\text{Add } \frac{a}{b} \text{ to } \frac{c}{d}$$

$$\text{reduced } \frac{ad}{bd} \text{ and } \frac{bc}{bd}; \text{ then } \frac{ad+bc}{bd} = \text{sum.}$$

Ex. 2.

$$\text{Add } \frac{a}{b}, \frac{c}{d}, \frac{f}{g} \text{ together.}$$

$$\text{reduced } \frac{adg}{bdg}, \frac{bcg}{bdg}, \frac{bdf}{bdg}; \text{ then } \frac{adg+bcg+bdf}{bdg} = \text{sum.}$$

Ex. 3.

$$\text{Add } \frac{a-b}{3c} \text{ to } \frac{2b-4a+d}{3c},$$

$$\text{the sum} = \frac{a-b+2b-4a+d}{3c} = \frac{b-3a+d}{3c}.$$

D 3

Ex.

Ex. 4.

$$\begin{array}{r}
 \text{To } a - \frac{aa}{b} \\
 \text{add } b + \frac{a-b}{c} \\
 \hline
 \text{sum } a+b + \frac{ab-bb-cao}{cb}
 \end{array}$$

PROBLEM XIV.

To subtract one fraction from another.

R U L E.

Reduce them to a common denominator; then subtract the numerators: and under the difference, write the common denominator.

Ex. 1.

$$\begin{array}{l}
 \text{From } \frac{a+b}{d} \text{ subtract } \frac{c}{d}. \\
 \frac{a-b-c}{d} = \text{difference.}
 \end{array}$$

Ex. 2.

$$\begin{array}{l}
 \text{From } \frac{a+b}{d} = \frac{ab+bb}{bd} \\
 \text{subtract } \frac{aa}{b} = \frac{aad}{bd}; \\
 \text{then } \frac{ab+bb-aad}{bd} = \text{remainder.}
 \end{array}$$

Ex. 3.

$$\begin{array}{l}
 \text{From } \frac{a-b}{3c} \text{ take } \frac{2b-4a}{5d}. \\
 \text{reduced } \frac{5ad-5bd}{15cd}, \frac{6bc-12ac}{15cd}. \\
 \text{remainder } \frac{5ad-5bd-6bc+12ac}{15cd}
 \end{array}$$

Ex.

Ex. 4.

From $a - \frac{aa}{b}$, or $a - \frac{aac}{bc}$
 take $b + \frac{a-b}{c}$, or $b + \frac{ab-bb}{bc}$

difference $a-b + \frac{-aac-ab+bb}{bc}$

PROBLEM XV.

To multiply fractions.

I R U L E.

In fractions, multiply the numerators together for a new numerator; and multiply the denominators together for a new denominator.

Ex. 1:

Multiply $\frac{a}{b}$ by $\frac{c}{d}$.

then $\frac{a \times c}{b \times d}$ or $\frac{ac}{bd} =$ product.

Ex. 2.

Multiply $\frac{b}{c}$ by $\frac{a+b}{b+c}$.

here $\frac{b}{c} \times \frac{a+b}{b+c} = \frac{ab+bb}{bc+cc}$, product.

Ex. 3.

Multiply $\frac{aa-bb}{bc}$ by $\frac{aa+bb}{b+c}$.

then $\frac{aa-bb}{bc} \times \frac{aa+bb}{b+c} = \frac{a^4-b^4}{bbc+bcc}$, product.

D 4

2 R U L E.

2 R U L E.

When the numerator of one, and denominator of the other, can be divided by some common divisor, take the quotients instead thereof.

Ex. 4.

$$\text{Let } \frac{c}{aa} \text{ multiply } \frac{aabb}{3cdd}.$$

$$\text{reduced } \frac{1}{1} \times \frac{bb}{3dd} = \frac{bb}{3dd}, \text{ product.}$$

Ex. 5.

$$\text{Multiply } \frac{aa+2ab+bb}{cd-dd} \text{ by } \frac{dd}{a+b}.$$

$$\text{reduced } \frac{a+b}{c-d} \times \frac{d}{1} = \frac{ad+bd}{c-d}, \text{ product.}$$

3 R U L E.

If a fraction is to be multiplied by an integer, which happens to be the same with the denominator; take the numerator for the product.

Ex. 6.

$$\text{Multiply } \frac{aa-2bb}{a-b} \text{ by } a-b.$$

$$\text{quotient } aa-2bb.$$

4 R U L E.

When a fraction is to be multiplied by an integer; multiply the numerator by the integer.

Ex. 7.

$$\text{Multiply } \frac{aa+3bb}{3cd} \text{ by } xx.$$

$$\text{then } \frac{aaxx+3bbxx}{3cd} \text{ or } \frac{aa+3bb}{3cd} xx = \text{the prod.}$$

Ex.

Ex. 8.

Multiply $\frac{2a-2x}{3b}$ by $a+x$ then $\frac{2aa-2xx}{3b} = \text{product.}$

Ex. 9.

Multiply $a + \frac{b-c}{d}$.by $b - \frac{b+c}{d}$ product $ab + \frac{bb-bc}{d} - \frac{ab+ac}{d} - \frac{bb-cc}{dd}$.

Schol. By this rule, a compound fraction may be reduced to a simple one.

PROBLEM XVI.

To divide one fraction by another.

R U L E.

In fractions, multiply the denominator of the divisor by the numerator of the dividend, for a new numerator; also multiply the numerator of the divisor into the denominator of the dividend, for a new denominator.

Ex. 1.

Divide $\frac{a}{b}$ by $\frac{c}{d}$. $\frac{c}{d}) \frac{a}{b} \left(\frac{ad}{bc} \right.$ the quotient.

Ex.

Ex. 2.

Let $\frac{a+c}{a-b}$ divide $\frac{a+b}{a}$.

$$\frac{a+c}{a-b} \frac{a+b}{a} \left(\frac{aa-bb}{aa+ac} \right) \text{ quotient.}$$

2 R U L E.

If the fractions have a common denominator; take the numerator of the dividend, for a numerator; and the numerator of the divisor, for the denominator.

Ex. 3.

Divide $\frac{aa-bb}{a+d}$ by $\frac{2ab-bb}{a+d}$.

$$\text{quotient } \frac{aa-bb}{2ab-bb}.$$

Ex. 4.

Let $\frac{aa+2ab+bb}{c-d}$ divide $\frac{a^2-abb}{c-d}$,

then $\frac{a^2-abb}{aa+2ab+bb} = \text{quotient.}$

or $\frac{aa-ab}{a+b} = \text{quotient reduced.}$

3 R U L E.

When fractions are to be divided by integers; multiply the denominators of the fractions, by such integers.

Ex. 5.

Divide $\frac{a-b}{c}$ by d .

quotient is $\frac{a-b}{cd}$.

Ex.

Ex. 6.

Let $a+b$ divide $\frac{aa-2bb}{a-b}$.then $\frac{aa-2bb}{a-b \times a+b} = \frac{aa-2bb}{aa-bb}$, quotient.

4 R U L E.

When the two numerators, or the two denominators, can be divided by some common divisor; throw out such divisor, and proceed by Rule 1.

Ex. 7.

Let $\frac{a-b}{cd}$ divide $\frac{ac-bb}{c+d}$,reduced $\frac{1}{cd} \left) \frac{a+b}{c+d} \left(\frac{acd+bcd}{c+d} \right)$, quotient.

Ex. 8.

Let $\frac{aa+ab}{a-b}$ divide $\frac{a^4-b^4}{aa-2ab+bb}$.reduced $\frac{a}{1} \left) \frac{a^3-a^2b+abb-b^3}{a-b} \left(\frac{a^3-aab+abb-b^3}{aa-ab} \right)$
the quotient.that is, the quotient $= a + \frac{bb}{a}$.

From hence may be deduced the following corollaries.

Cor. 1. The value of any fractional quantity is not at all changed, by changing all the signs of both numerator and denominator. Thus $\frac{ab-ac}{r-c} = \frac{ac-ab}{c-r}$.

Cor. 2. The value of any compound fractional quantity, is equal to the sum of all the particular simple

simple fractions, that compose it. Thus

$$\frac{rx + 2cx - 11rx}{3r - 2x} = \frac{rx}{3r - 2x} + \frac{2cx}{3r - 2x} - \frac{11rx}{3r - 2x}$$

Cor. 3. If a fraction be multiplied by any given quantity; it is the same thing whether the numerator be multiplied by that quantity, or the denominator divided by it. $\frac{dab}{dc} \times d = \frac{dabd}{dc} = \frac{dab}{c}$.

Cor. 4. The product of two fractions, is equal to the fraction, that has the product of the numerators for the numerator; and the product of the denominators for its denominator.

$$\frac{a}{b+x} \times \frac{r-c}{x} = \frac{a \times r - c}{b+x \times x} = \frac{ar - ac}{bx + xx}$$

Cor. 5. If a fraction is to be divided by some quantity; it is the same thing whether the numerator be divided by it, or the denominator multiplied.

$$\text{For } \frac{2az}{x} \div r = \frac{2az}{rx}. \text{ And } \frac{2ar}{x} \div r = \frac{2a}{x}.$$

Cor. 6. If any sort of quantity is to be divided by a fraction; it is the same thing, as to multiply the said quantity, by the fraction inverted. Thus

$$ab \div \frac{r}{s} = ab \times \frac{s}{r}. \text{ And } \frac{a}{c} \div \frac{b}{r} \text{ or}$$

$$\frac{\frac{a}{c}}{\frac{b}{r}} = \frac{a}{c} \times \frac{r}{b} = \frac{ar}{bc}.$$

PROBLEM XVII.

To involve fractional quantities.

R U L E.

Involve the numerator into itself, for a new numerator; and the denominator into itself for a new denominator; each as often as the index of the power.

Ex.

Ex. 1.

<i>Involve</i>	$\frac{a}{b}$	<i>and</i>	$\frac{1}{a a}$
root	$\frac{a}{b}$		$\frac{1}{aa}$
square	$\frac{aa}{bb}$		$\frac{1}{a^4}$
cube	$\frac{a^3}{b^3}$		$\frac{1}{a^6}$
4th power	$\frac{a^4}{b^4}$		$\frac{1}{a^8}$
&c.			

Ex. 2.

Let $\frac{3bc}{2ad}$, *and* $\frac{-ad}{4bb}$, *be involved.*

root	$\frac{3bc}{2ad}$		$-\frac{ad}{4bb}$
square	$\frac{9bbcc}{4aadd}$		$+\frac{aadd}{16b^4}$
cube	$\frac{27b^3c^3}{8a^3d^3}$		$-\frac{a^3d^3}{64b^6}$
&c.			

Ex. 3.

Involve $\frac{aa-bc}{a+c}$ *to the square, &c.*

$\frac{a^4-2aabc+bbcc}{aa+2ac+cc}$, *the square.*

$\frac{a^6-3a^2bc+3a^2bbcc-b^3c^3}{a^3+3a^2c+3ac^2+c^3}$, *cube, &c.*

Ex.

Ex. 4.

Involve $\frac{a-x}{2b}$ to the 4th power.

it is $\frac{a^4 - 4a^3x + 6a^2x^2 - 4ax^3 + x^4}{16b^4}$.

or thus $\left(\frac{a-x}{2b}\right)^4$ or $\frac{a-x^4}{16b^4}$.

P R O B L E M XVIII.

To extract the root of a fraction.

R U L E.

Extract the proper root of both numerator and denominator, if it can be done. If not set the radical sign ($\sqrt{\quad}$) before one or both of them, as they happen to be surd.

Ex. 1.

What is the square root of $\frac{9a^2b^4}{4dd}$.

root $\frac{3abb}{2d}$.

Ex. 2.

What the cube root of $\frac{a^3 - 3a^2b + 3ab^2 - b^3}{a^3 + 3a^2b + 3ab^2 + b^3}$.

the root is $\frac{a-b}{a+b}$.

Ex. 3.

The square root of $\frac{aabb}{d^2}$, is $\sqrt{\frac{aabb}{d^2}}$ or $\frac{ab}{\sqrt{d^2}}$.

Ex.

Ex. 4.

What is the cube root of $\frac{-27a^3b^3}{a^3-b^3}$.

the root is $\frac{-3ab}{\sqrt[3]{a^3-b^3}}$

Ex. 5.

What is the cube root of $\frac{a^3+4abd-d^3}{8a^6b^3}$.

the root is $\sqrt[3]{\frac{a^3+4abd-d^3}{8a^6b^3}}$, or $\frac{\sqrt[3]{a^3+4abd-d^3}}{2aab}$

Ex. 6.

What is the 4th root of $\frac{x^4-y^4}{8ax^3-8x^2yy+y^4}$.

the root is $\sqrt[4]{\frac{x^4-y^4}{8ax^3-8x^2yy+y^4}}$

or $\frac{\sqrt[4]{x^4-y^4}}{\sqrt[4]{8ax^3-8x^2yy+y^4}}$

Cor. The n^{th} power or root of a fraction, is equal to the n^{th} power or root of the numerator, divided by the n^{th} power or root of the denominator.

$$\left(\frac{a}{x}\right)^n = \frac{a^n}{x^n}. \text{ And } \sqrt[n]{\frac{a}{x}} = \frac{\sqrt[n]{a}}{\sqrt[n]{x}}.$$

S E C T. III.

O f S U R D S.

SURDS are such quantities as have not a proper root. *Simple Surds* are those which consist but of one term. *Compound Surds* are those which consist of several simple ones. And *Universal Surds* are those consisting of several terms under any radical sign.

Surds are said to be *commensurable*, when they are as one number to another; and *incommensurable*, when their proportion cannot be expressed in numbers.

P R O B L E M XIX.

To designate or express the roots of quantities by fractional indices.

R U L E.

Divide the index of the quantity by the number expressing the root; the quotient is the index of the root required.

Ex. 1.

Let the quantity a be proposed.

then $\sqrt{a} = a^{\frac{1}{2}}$, $\sqrt[3]{a} = a^{\frac{1}{3}}$, $\sqrt[4]{a} = a^{\frac{1}{4}}$, &c.

Ex. 2.

Let $3ab^2$ be proposed.

$\sqrt{3ab^2} = \sqrt[2]{3abb^2} = a^{\frac{1}{2}}b\sqrt{3}$. $\sqrt[3]{3ab^2} = a^{\frac{1}{3}}b^{\frac{2}{3}}\sqrt[3]{3}$.

$\sqrt[4]{3ab^2} = a^{\frac{1}{4}}b^{\frac{1}{2}}\sqrt[4]{3}$. $\sqrt[5]{3abb} = a^{\frac{1}{5}}b^{\frac{2}{5}}\sqrt[5]{3}$, &c.

Ex.

Ex. 3.

Let a^3 be given.

then $\sqrt{a^3} = a^{\frac{3}{2}}$, $\sqrt[3]{a^3} = a^{\frac{3}{3}}$ or a , $\sqrt[4]{a^3} = a^{\frac{3}{4}}$, &c.

Ex. 4.

Let $\overline{aa-xx}$ be proposed.

then $\sqrt{aa-xx} = \overline{aa-xx}^{\frac{1}{2}}$,

$\sqrt[3]{aa-xx} = \overline{aa-xx}^{\frac{1}{3}}$, &c.

Ex. 5.

Let $\frac{1}{x}$ be given.

then $\sqrt{\frac{1}{x}} = \frac{1}{x^{\frac{1}{2}}}$, $\sqrt[3]{\frac{1}{x}} = \frac{1}{x^{\frac{1}{3}}}$, $\sqrt[4]{\frac{1}{x}} = \frac{1}{x^{\frac{1}{4}}}$
&c.

Ex. 6.

Let $\frac{a-b}{abbc^3}$ be given.

then $\sqrt{\frac{a-b}{abbc^3}} = \frac{\overline{a-b}^{\frac{1}{2}}}{a^{\frac{1}{2}}bc^{\frac{3}{2}}}$, $\sqrt[3]{\frac{a-b}{abbc^3}} = \frac{\overline{a-b}^{\frac{1}{3}}}{a^{\frac{1}{3}}b^{\frac{1}{3}}c}$, &c.

Ex. 7.

Let $\frac{aa}{b^3}$ be proposed.

$\sqrt{\frac{aa}{b^3}} = \frac{a}{b^{\frac{3}{2}}}$, $\sqrt[3]{\frac{aa}{b^3}} = \frac{a^{\frac{2}{3}}}{b}$, $\sqrt[4]{\frac{aa}{b^3}} = \frac{a^{\frac{1}{2}}}{b^{\frac{3}{4}}}$, &c.

Ex. 8.

Let $\frac{a+2x}{aa-xx}$ be proposed.

$$\begin{aligned} \sqrt{\frac{a+2x}{aa-xx}} &= \frac{\overline{a+2x}^{\frac{1}{2}}}{\overline{aa-xx}^{\frac{1}{2}}}, \quad \sqrt[3]{\frac{a+2x}{aa-xx}} = \frac{\overline{a+2x}^{\frac{1}{3}}}{\overline{aa-xx}^{\frac{1}{3}}} \\ &= \frac{\overline{a+2x}^{\frac{1}{3}}}{\overline{aa-xx}^{\frac{1}{3}}}. \end{aligned}$$

2 R U L E.

When any quantity is in the denominator of a fraction; set it in the numerator, and change the sign of the index.

Ex. 9.

$$\sqrt{\frac{1}{a}} = \frac{1}{a^{\frac{1}{2}}} = \frac{a^{-\frac{1}{2}}}{1} = a^{-\frac{1}{2}}.$$

Ex. 10.

Let $\frac{1}{a}$, $\frac{1}{aa}$, $\frac{1}{a^3}$, $\frac{1}{a^4}$, $\frac{1}{a^5}$, &c. be given.
then they become a^{-1} , a^{-2} , a^{-3} , a^{-4} , a^{-5} , &c.
respectively.

Ex. 11.

Given $\frac{ab}{x^2y^3}$. This becomes $abx^{-2}y^{-3}$.

Ex. 12.

Given $\frac{1}{aa-xx}$, $\frac{1}{aa-xx^2}$, $\frac{1}{aa-xx^3}$, &c.

they become $\overline{aa-xx}^{-1}$, $\overline{aa-xx}^{-2}$, $\overline{aa-xx}^{-3}$, &c.

Ex.

Ex. 13.

Let $\frac{aab}{a+x}$, $\frac{aab}{a+x^2}$, $\frac{aab}{a+x^3}$ be given.

they are $aab \times \overline{a+x}^{-1}$, $aab \times \overline{a+x}^{-2}$,
 $aab \times \overline{a+x}^{-3}$.

In order to explain this; let there be a rank of powers, as 1, a , aa , aaa , $aaaa$, $aaaaa$, &c. the same will (by Def. 20.) be denoted 1, a , a^2 , a^3 , a^4 , a^5 , &c. Now these quantities, a , a^2 , a^3 , &c. are in geometrical progression, and their indices, in arithmetic progression, as is plain. Now since 1, a , a^2 , a^3 , &c. are geometrical proportionals, therefore these will also be geometrical proportionals,

1, \sqrt{a} , $\sqrt{a^2}$, $\sqrt{a^3}$, $\sqrt{a^4}$, $\sqrt{a^5}$, $\sqrt{a^6}$, $\sqrt{a^7}$,
 $\sqrt{a^8}$, &c.

that is 1, $\sqrt[3]{a}$, a , $\sqrt[3]{a^3}$, aa , $\sqrt[3]{a^5}$, a^3 , $\sqrt[3]{a^7}$, a^4 , &c.

and 1, $\sqrt[4]{a}$, \sqrt{aa} , a , $\sqrt[4]{a^4}$, $\sqrt[4]{a^5}$, a^2 , $\sqrt[4]{a^7}$,
 $\sqrt[4]{a^8}$, &c.

and 1, $\sqrt[5]{a}$, $\sqrt[4]{aa}$, $\sqrt[4]{a^3}$, a , $\sqrt[5]{a^5}$, $\sqrt[5]{a^6}$,
 $\sqrt[5]{a^7}$, aa , &c. and so on.

Therefore by the rule of analogy, the indices of all these, are also in arithmetic progression.

Take any one of these series as 1, $\sqrt[3]{a}$, \sqrt{aa} , a , $\sqrt[3]{a^3}$, &c. these will be equivalent to 1, $a^{\frac{1}{3}}$, $a^{\frac{2}{3}}$, a , $a^{\frac{4}{3}}$, &c.

Suppose now the series 1, a , a^2 , a^3 , &c. continued backwards, the powers of a will come into the denominator; and the indices, which continually decrease, will then become negative, and will stand thus:

Powers' $\frac{1}{a^3}, \frac{1}{a^2}, \frac{1}{a^1}, 1, a, a^2, a^3, a^4, \&c.$

Indices $-3, -2, -1, 0, 1, 2, 3, 4, \&c.$
therefore $a^{-3}, a^{-2}, a^{-1}, a^0, a^1, a^2, a^3, a^4, \&c.$
will represent these powers; that is,

$$\frac{1}{a^3} = a^{-3}, \quad \frac{1}{a^2} = a^{-2}, \quad \frac{1}{a} = a^{-1}, \quad 1 = a^0, \quad \&c.$$

In like manner, let the series $1, a^{\frac{1}{3}}, a^{\frac{2}{3}}, a^1, a^{\frac{4}{3}}, \&c.$ be continued backwards; these powers, and their indices will be as follows:

$$\frac{1}{a^{\frac{4}{3}}}, \frac{1}{a}, \frac{1}{a^{\frac{2}{3}}}, \frac{1}{a^{\frac{1}{3}}}, 1, a^{\frac{1}{3}}, a^{\frac{2}{3}}, a^1, a^{\frac{4}{3}}, \&c.$$

$$-\frac{4}{3}, -1, -\frac{2}{3}, -\frac{1}{3}, 0, \frac{1}{3}, \frac{2}{3}, 1, \frac{4}{3}, \&c.$$

then, $a^{-\frac{4}{3}}, a^{-1}, a^{-\frac{2}{3}}, a^{-\frac{1}{3}}, a^0, a^{\frac{1}{3}}, a^{\frac{2}{3}}, a, a^{\frac{4}{3}}, \&c.$ will denote the same powers; that is,

$$\frac{1}{a^{\frac{4}{3}}} = a^{-\frac{4}{3}}, \quad \frac{1}{a} = a^{-1}, \quad \frac{1}{a^{\frac{2}{3}}} = a^{-\frac{2}{3}}, \quad \frac{1}{a^{\frac{1}{3}}} = a^{-\frac{1}{3}},$$

$$a^0 = 1, \quad \&c. \quad \text{And therefore the series,}$$

$$\frac{1}{\sqrt[3]{aaaa}}, \frac{1}{\sqrt[3]{aaa}}, \frac{1}{\sqrt[3]{aa}}, \frac{1}{\sqrt[3]{a}}, 1, \sqrt[3]{a},$$

$$\sqrt[3]{aa}, \sqrt[3]{aaa}, \sqrt[3]{aaaa}, \&c. \text{ may be expressed thus,}$$

$$\frac{1}{\sqrt[3]{a^4}}, \frac{1}{\sqrt[3]{a^3}}, \frac{1}{\sqrt[3]{a^2}}, \frac{1}{\sqrt[3]{a}}, 1, \sqrt[3]{a}, \sqrt[3]{a^2}, \sqrt[3]{a^3},$$

$$\sqrt[3]{a^4}, \&c.$$

$$\text{or thus, } \frac{1}{a^{\frac{4}{3}}}, \frac{1}{a}, \frac{1}{a^{\frac{2}{3}}}, \frac{1}{a^{\frac{1}{3}}}, 1, a^{\frac{1}{3}}, a^{\frac{2}{3}}, a^1, a^{\frac{4}{3}}, \&c.$$

or thus, $a^{-\frac{1}{2}}$, a^{-1} , $a^{-\frac{3}{2}}$, a^{-2} , a^0 , $a^{\frac{1}{2}}$, a^1 , $a^{\frac{3}{2}}$, a^2 , $a^{\frac{5}{2}}$, &c. and the same is equally true of any of the other series.

Cor. 1. *The powers of any quantity are a set of geometrical proportionals from 1; and their indices, a set of arithmetic proportionals from 0.*

thus, powers 1, a , a^2 , a^3 , a^4 , \ddots increasing.

indices 0, 1, 2, 3, 4, \ddots increasing.

also, powers 1, $\frac{1}{a}$, $\frac{1}{aa}$, $\frac{1}{a^3}$, $\frac{1}{a^4}$, \ddots decreas.

indices 0, -1, -2, -3, -4, \ddots decreas.

Cor. 2. *Hence the double, triple, quadruple, &c. the index of any quantity, is the index of the square, cube, biquadrate, &c. of that quantity.*

Cor. 3. *Hence also, the index of the product of any two powers (whole or fracted) of any quantity, is equal to the sum of the indices of these powers. And therefore to multiply any two powers together, is to add their indices. Thus $a^2 \times a^3 = a^5$, $a^2 \times a^{-3} = a^{-1}$, &c.*

Cor. 3. *The index of the quotient of two powers, dividing one another, is equal to the index of the dividend — the index of the divisor; whatever the indices be. And therefore, to divide by powers, is to*

subtract their indices. Thus $\frac{a^3}{a} = a^2$, and

$$\frac{a^2}{a^5} = a^{-3}. \quad \text{Also } \frac{a^3}{a^{-2}} = a^5, \text{ \&c.}$$

Cor. 4. *Any power is taken out of the denominator, and put into the numerator, by changing the sign of the index: and the contrary. Thus*

$$\frac{1}{a^3} = a^{-3}, \quad \frac{b}{a^2} = ba^{-2}. \quad \text{Also}$$

$$\frac{a^2}{b^3} = a^2 b^{-3} = \frac{1}{a^{-2} b^3}, \text{ \&c.}$$

Cor. 5. In fractional indices, the numerator shews the power, and the denominator the root.

Schol. In all the following problems, it will be the best way to reduce the surds to fractional indices.

P R O B L E M XX.

To reduce a rational quantity to the form of a surd.

R U L E.

Multiply the index of the quantity, by the index of the surd root given; to which set the radical sign, or index of the surd.

Ex. 1.

Reduce 6 to the form of $\sqrt{5}$.

Here $6^{1 \times 2}$ or $6^2 = 36$, and $\sqrt{36}$ is that required.

Ex. 2.

Reduce a to the form of $\sqrt[3]{b}$.

Here $a^{1 \times 3} = a^3$, and $\sqrt[3]{a^3}$ is the answer.

Ex. 3.

Reduce $a+b$ to the form of \sqrt{bc} .

Anfw. $\sqrt{a+b}$, or $\sqrt{aa+2ab+bb}$.

Ex. 4.

Reduce $\frac{a}{b\sqrt{c}}$ to the form of \sqrt{d} .

Anf. $\sqrt{\frac{aa}{bbc}}$ is of the form \sqrt{d} .

PROBLEM XXI.

To reduce quantities of different indexes, to other equal ones, that shall have a common index given.

R U L E.

Divide the indexes of the quantities by the given index; the quotients will be the new indexes for these quantities. Over these quantities with their new indices, place the index given.

Ex. 1.

Reduce $12^{\frac{1}{4}}$ and $7^{\frac{1}{6}}$ to the common index $\frac{1}{2}$.

$$\left. \begin{array}{l} \frac{1}{2} \left) \frac{1}{4} \left(\frac{1}{2} \text{ first index.} \\ \frac{1}{3} \left) \frac{1}{6} \left(\frac{1}{3} \text{ second index} \end{array} \right\} \begin{array}{l} \text{then } \sqrt[1]{12^{\frac{1}{2}}} \text{ and } \sqrt[1]{7^{\frac{1}{2}}} \text{ are} \\ \text{the quantities required.} \end{array}$$

Ex. 2.

Reduce a^3 and $b^{\frac{3}{2}}$, to the common index $\frac{1}{2}$.

$$\left. \begin{array}{l} \frac{1}{3} \left) \frac{2}{1} \left(6 \text{ first index.} \\ \frac{1}{3} \left) \frac{3}{2} \left(\frac{9}{2} \text{ sec. index.} \end{array} \right\} \begin{array}{l} \text{then } \sqrt[1]{a^6} \text{, and } \sqrt[1]{b^{\frac{9}{2}}} \text{ are the} \\ \text{furds.} \end{array}$$

PROBLEM XXII.

To reduce quantities of different indices, to others equal to them, that shall have the least common index.

R U L E.

Reduce the indices of the given quantities, to a common denominator, in the least terms. Then involve each quantity to the power of its numerator; and take the root denoted by the common denominator.

E 4

Ex.

Ex. 1.

Reduce $b^{\frac{1}{4}}$ and $c^{\frac{1}{6}}$ to the least common index.

$\frac{1}{4}$ and $\frac{1}{6}$ are $= \frac{3}{12}$ and $\frac{2}{12}$. Therefore

$$b^{\frac{1}{4}} = b^{\frac{3}{12}}, \text{ and } c^{\frac{1}{6}} = c^{\frac{2}{12}}.$$

or $b^{\frac{1}{4}}$ and $c^{\frac{1}{6}}$ become $\overline{b^3}^{\frac{1}{12}}$ and $\overline{c^2}^{\frac{1}{12}}$, or $\overline{bbb}^{\frac{1}{12}}$ and $\overline{cc}^{\frac{1}{12}}$.

Ex. 2.

Let $b^{\frac{2}{3}}$ and $a^{\frac{1}{5}}$ be given.

$\frac{2}{3}$ and $\frac{1}{5}$ are reduced to $\frac{6}{9}$ and $\frac{2}{9}$.

Therefore $b^{\frac{2}{3}}$ and $a^{\frac{1}{5}}$ become $b^{\frac{6}{9}}$ and $a^{\frac{2}{9}}$,

or $\overline{b^6}^{\frac{1}{9}}$ and $\overline{a^2}^{\frac{1}{9}}$, or $\overline{b^6}^{\frac{1}{9}}$ and $\overline{aa}^{\frac{1}{9}}$.

Ex. 3.

Let $\sqrt{a+b}$, and $\sqrt[3]{aa-xx}$ be proposed.

These are $\overline{a+b}^{\frac{1}{2}}$ and $\overline{aa-xx}^{\frac{1}{3}}$. The indices are reduced to $\frac{3}{6}$ and $\frac{2}{6}$. Therefore the surds

become $\overline{a+b}^{\frac{3}{6}}$ and $\overline{aa-xx}^{\frac{2}{6}}$; or

$\overline{a^3+3aab+3abb+b^3}^{\frac{1}{2}}$, and $\overline{a^4-2a^2xx+x^4}^{\frac{1}{3}}$, or $\sqrt{a^3+3aab+3abb+b^3}$, and $\sqrt[3]{a^4-2a^2xx+x^4}$.

PROBLEM XXIII.

To reduce surds to their most simple terms.

R U L E.

Divide by the greatest power contained in it, and set the root before the surd containing the remaining quantities.

Ex.

Ex. 1.

Reduce $\sqrt{48}$ to a simpler form.

$$\sqrt{48} = \sqrt{3 \times 16} = 4\sqrt{3} \text{ the surd required.}$$

Ex. 2.

Let $\sqrt{64aabc}$ be proposed.

$$\sqrt{64aa} = 8a. \text{ Then } \sqrt{64aabc} = 8a\sqrt{bc}.$$

Ex. 3.

Reduce $\sqrt{a^2x - a^2x^2}$.

Here $\sqrt{aa} = a$, and the surd becomes $a\sqrt{ax - xx}$
or $a\sqrt{ax - xx}$.

Ex. 4.

Let $\sqrt{\frac{a^2b - 4aabb + 4ab^2}{cc}}$ be given.

The surd is $\sqrt{\frac{aa - 4ba + 4bb}{cc}} \times ab$. And

$$\sqrt{\frac{aa - 4ba + 4bb}{cc}} = \frac{a - 2b}{c}. \text{ Therefore the surd}$$

becomes $\frac{a - 2b}{c} \times \sqrt{ab}$, or $\frac{a - 2b}{c} \sqrt{ab}$.

Ex. 5.

$$\text{Given } \sqrt{\frac{27a^2b^3}{8b - 8a}} = \sqrt[3]{\frac{27a^2b^3 \times a}{8 \times b - a}}$$

$$\text{reduced, becomes } \frac{3ab}{2} \sqrt[3]{\frac{a}{b-a}}.$$

PROBLEM XXIV.

To find whether two surds are commensurable or not.

R U L E.

Reduce them to the least common index, and the quantities to a common denominator, if fractions, except when like terms are commensurable.

Then divide them by the greatest common divisor, (or by such a one as will give one quotient rational;) then if both quotients be rational, the surds are commensurable; otherwise not.

Ex. 1.

Let $\sqrt{18}$ and $\sqrt{8}$ be proposed.

These are $\sqrt{2 \times 9}$ and $\sqrt{2 \times 4}$. Divide by 2, and the quotients are $\sqrt{9}$, and $\sqrt{4}$; that is, 3 and 2; therefore they are commensurable.

Ex. 2.

Let the surds be $\sqrt{\frac{50}{16}}$ and $\sqrt{\frac{72}{25}}$.

These are $\frac{\sqrt{50}}{4}$ and $\frac{\sqrt{72}}{5}$. Divide by 2, and the quotients are $\sqrt{25}$ and $\sqrt{36}$, that is 5 and 6; and the surds become $\frac{5}{4}\sqrt{2}$ and $\frac{6}{5}\sqrt{2}$, and are therefore commensurable, being as $\frac{5}{4}$ to $\frac{6}{5}$.

Ex. 3.

Let $\sqrt{48}$ and $\sqrt{8}$ be proposed.

Divide by 8, the quotients are $\sqrt{6}$ and $\sqrt{1}$ or 1; therefore they are incommensurable.

Ex.

Ex. 4.

Let $\sqrt{\frac{b}{c}}$ and $\sqrt{\frac{c}{b}}$ be given.

Here $\sqrt{\frac{b}{c}}$ and $\sqrt{\frac{c}{b}}$ are reduced to $\sqrt{\frac{b}{c}}$ and $\sqrt{\frac{c}{b^3}}$, and these to $\sqrt{\frac{b^4}{cb^3}}$ and $\sqrt{\frac{c^4}{cb^3}}$. Divide by $\frac{1}{cb^3}$, and the quotients are \sqrt{b} and $\sqrt{c^4}$, that is, bb and cc ; therefore the surds are commensurable.

Ex. 5.

Suppose $\sqrt{a^2 + aabb}$ and $\sqrt{aabb + b^2}$.

These are $\sqrt{aa \times aa + bb}$, and $\sqrt{bb \times aa + bb}$. Therefore dividing by $aa + bb$, the quotients are \sqrt{aa} , and \sqrt{bb} , or a and b , and therefore they are commensurable.

Ex. 6.

Let $\sqrt{\frac{16aa}{14b}}$ and $\sqrt{\frac{9aa}{8b}}$ be given.

That is, $\frac{4a}{\sqrt{14b}}$ and $\frac{3a}{\sqrt{8b}}$. Divide the denominators by 2, then they are reduced to $\frac{4a}{\sqrt{7b}}$ and $\frac{3a}{\sqrt{4b}}$ or $\frac{3a}{2\sqrt{b}}$, and are therefore incommensurable.

P R O B L E M XXV.

To add surd quantities together.

R U L E.

Reduce quantities with unlike indexes, to those of like indexes.

Also reduce fractions to a common denominator, or else to others that have rational denominators (or numerators).

Then reduce the quantities to the simplest terms (Prob. 23.) This being done; if the surd part be the same in all, annex it to the sum of the rational parts, with the sign (\times) of multiplication.

If the surd part is not the same in all, the quantities can only be added by the signs $+$ and $-$.

Ex. 1.

Add $\sqrt{6}$ to $2\sqrt{6}$.

The sum is $1+2 \times \sqrt{6}$ or $3\sqrt{6}$.

Ex. 2.

Add $\sqrt{8}$ to $\sqrt{50}$.

$\sqrt{8} = 2\sqrt{2}$, and $\sqrt{50} = 5\sqrt{2}$, and the sum
 $= 2+5 \times \sqrt{2} = 7\sqrt{2} = \sqrt{98}$.

Ex. 3.

Add $\sqrt[3]{500}$ to $\sqrt[3]{108}$.

$\sqrt[3]{500} = \sqrt[3]{4 \times 125} = 5\sqrt[3]{4}$. And
 $\sqrt[3]{108} = \sqrt[3]{4 \times 27} = 3\sqrt[3]{4}$. Therefore the sum
 $= 5+3 \times \sqrt[3]{4} = 8\sqrt[3]{4}$.

Ex. 4.

Add $\sqrt{48a^2b}$ to $\sqrt{3aab^3}$.

They are reduced to $4aa\sqrt{3b}$ and $ab\sqrt{3b}$.
 And the sum $= 4aa+ab \times \sqrt{3b}$.

Ex.

Ex. 5.

Given $\sqrt{4a}$ and $\sqrt[4]{a^6}$.

$$\sqrt{4a} = \overline{4a}^{\frac{1}{2}} = \overline{4a^2} = \overline{16aa}^{\frac{1}{4}} = \sqrt[4]{16aa} =$$

$$2\sqrt[4]{aa}. \quad \text{And } \sqrt[4]{a^6} = a\sqrt[4]{aa}. \quad \text{And their sum}$$

$$= \overline{a+2} \times \sqrt[4]{aa} = \overline{a+2} \times \sqrt{a}.$$

Ex. 6.

Add $\sqrt{\frac{24}{25}}$ to $\sqrt{\frac{2}{3}}$.

These reduced to a common denominator, become $\sqrt{\frac{72}{75}}$ and $\sqrt{\frac{50}{75}}$, or $\sqrt{\frac{2 \times 36}{75}}$ and $\sqrt{\frac{2 \times 25}{75}}$,

that is, $6\sqrt{\frac{2}{75}}$ and $5\sqrt{\frac{2}{75}}$, whose sum is

$$11\sqrt{\frac{2}{75}};$$

Or thus,

$$\text{Here } \sqrt{\frac{24}{25}} = \frac{\sqrt{24}}{5} = \frac{\sqrt{4 \times 6}}{5} = \frac{2}{5}\sqrt{6};$$

$$\text{Also } \sqrt{\frac{2}{3}} = \sqrt{\frac{6}{9}} = \frac{\sqrt{6}}{3}. \quad \text{And their sum}$$

$$= \frac{2}{5} + \frac{1}{3} \times \sqrt{6} = \frac{11}{15}\sqrt{6} = \frac{11}{5}\sqrt{\frac{2}{3}}.$$

Ex. 7.

Add $\sqrt[3]{\frac{1}{4}}$ to $\sqrt[3]{\frac{16}{27}}$.

These become $\sqrt[3]{\frac{1}{4}}$ and $\sqrt[3]{\frac{64}{108}}$, or $\sqrt[3]{\frac{1}{4}}$

and

and $4\sqrt[3]{\frac{1}{4 \times 27}}$ that is $\sqrt[3]{\frac{1}{4}}$ and $\frac{4}{3}\sqrt[3]{\frac{1}{4}}$;

whose sum is $1 + \frac{4}{3} \times \sqrt[3]{\frac{1}{4}} = \frac{7}{3}\sqrt[3]{\frac{1}{4}}$.

Or thus,

$$\sqrt[3]{\frac{1}{4}} = \sqrt[3]{\frac{27}{108}} = 3\sqrt[3]{\frac{1}{4 \times 27}} = \frac{3}{3}\sqrt[3]{\frac{1}{4}},$$

$$\text{and } \sqrt[3]{\frac{16}{27}} = \sqrt[3]{\frac{64}{108}} = 4\sqrt[3]{\frac{1}{4 \times 27}} = \frac{4}{3}\sqrt[3]{\frac{1}{4}}.$$

$$\text{And the sum} = \frac{7}{3}\sqrt[3]{\frac{1}{4}}.$$

Ex. 8.

Add $\sqrt[3]{\frac{b}{c}}$ to $\sqrt[3]{\frac{c}{b}}$.

These are reduced to $\left(\frac{b^4}{cb^3}\right)^{\frac{1}{2}}$ and $\left(\frac{c^4}{cb^3}\right)^{\frac{1}{2}}$ or

$\frac{bb}{b}\sqrt{\frac{1}{bc}}$ and $\frac{cc}{b}\sqrt{\frac{1}{bc}}$. And their sum is

$$\frac{bb+cc}{b}\sqrt{\frac{1}{bc}} = \frac{bb+cc}{b\sqrt{bc}}.$$

Ex. 9.

Add $\sqrt{ccddaa - ccddxx}\sqrt{2}$ to $\sqrt{d^4aa - d^4xx}\sqrt{2}$.

They are reduced to $cd\sqrt{aa - xx}\sqrt{2}$, and $dd\sqrt{aa - xx}\sqrt{2}$, and the sum is $cd + dd \times \sqrt{aa - xx}\sqrt{2}$.

Ex. 10.

To $2\sqrt[3]{aa} - \sqrt{a^3} + \sqrt{13}$.

Add $\sqrt[3]{aa} + 2\sqrt{a} - \sqrt{7}$

Sum $3\sqrt[3]{aa} - \sqrt{a^3} + \sqrt{13} + 2\sqrt{a} - \sqrt{7}$.

PRO-

PROBLEM XXVI.

To subtract surd quantities.

R U L E.

Reduce, as in the last rule; then subtract the rational quantities, and annex the difference to the common surd, with the sign (\times) of multiplication.

E X A M P L E S.

1. Subtract $\sqrt{6}$ from $2\sqrt{6}$, the remainder is
 $2 - 1 \times \sqrt{6} = \sqrt{6}$.

$$2. \sqrt{50} - \sqrt{8} = 5\sqrt{2} - 2\sqrt{2} = 3\sqrt{2}.$$

$$3. \sqrt[3]{500} - \sqrt[3]{108} = 5\sqrt[3]{4} - 3\sqrt[3]{4} = 2\sqrt[3]{4}.$$

$$4. \sqrt{48a^2b} - \sqrt{3aab^3} = 4aa\sqrt{3b} - ab\sqrt{3b} \\ = 4aa - ab \times \sqrt{3b}.$$

$$5. \sqrt{a^6} - \sqrt{4a} = \sqrt{a^3} - \sqrt{4a} = a\sqrt{a} - \\ 2\sqrt{a} = a - 2 \times \sqrt{a}.$$

$$6. \sqrt{\frac{24}{25}} - \sqrt{\frac{2}{3}} = \sqrt{\frac{72}{75}} - \sqrt{\frac{50}{75}} = 6\sqrt{\frac{2}{75}} \\ - 5\sqrt{\frac{2}{75}} = \sqrt{\frac{2}{75}}.$$

$$\text{or } \sqrt{\frac{24}{25}} - \sqrt{\frac{2}{3}} = \sqrt{\frac{24}{25}} - \sqrt{\frac{6}{9}} = \\ \frac{2}{5}\sqrt{6} - \frac{1}{3}\sqrt{6} = \frac{1}{15}\sqrt{6}.$$

$$7. \sqrt[3]{\frac{16}{27}} - \sqrt[3]{\frac{1}{4}} = \frac{4}{3}\sqrt[3]{\frac{1}{4}} - \sqrt[3]{\frac{1}{4}} =$$

$$\frac{1}{3}\sqrt[3]{\frac{1}{4}}. \text{ Or thus, } \sqrt[3]{\frac{16}{27}} - \sqrt[3]{\frac{1}{4}} = \sqrt[3]{\frac{64}{4 \times 27}}$$

$$= \sqrt[3]{\frac{1}{4}} = \frac{4}{3}\sqrt[3]{\frac{1}{4}} - \sqrt[3]{\frac{1}{4}} = \frac{1}{3}\sqrt[3]{\frac{1}{4}}.$$

$$8. \quad \frac{\sqrt{b}}{c} - \frac{\sqrt{c}}{b} = \frac{b}{1} \times \frac{\sqrt{1}}{bc} - \frac{cc}{b} \times \frac{\sqrt{1}}{bc} = \frac{bb-cc}{b} \times \frac{\sqrt{1}}{bc}.$$

$$9. \quad \sqrt{ccddaa-ccddxx} - \sqrt{a^2aa-d^2xx} = cd\sqrt{aa-xx} - dd\sqrt{aa-xx} = cd-ad\sqrt{aa-xx}.$$

$$10. \quad \text{From } 2\sqrt{oa} - \sqrt{a^3} + \sqrt{13} \\ \text{take } \sqrt{aa} + 2\sqrt{a} - \sqrt{7}.$$

$$\text{rem. } \sqrt{aa} - \sqrt{a^3} + \sqrt{13} - 2\sqrt{a} + \sqrt{7}.$$

PROBLEM XXVII.

To multiply surds.

RULE.

Surds by surds; if they have not the same index already, reduce them to the same; then multiply the quantities under the common index.

Ex. 1.

$$\text{Multiply } \sqrt{5} \text{ by } \sqrt{3}. \quad \begin{array}{r} \sqrt{5} \\ \sqrt{3} \\ \hline \end{array}$$

the product $\sqrt{15}$

Ex. 2.

$$\text{Multiply } \sqrt{\frac{2ab}{3c}} \text{ by } \sqrt{\frac{9ad}{2b}}.$$

$$\text{product} = \sqrt{\frac{18abd}{6bc}} = \sqrt{\frac{3aad}{c}}.$$

Ex.

Ex. 3.

Multiply \sqrt{d} by $\sqrt[3]{ab}$.

Reduced to $\sqrt[6]{a^2d^3}$ and $\sqrt[6]{aabb^2}$; the product
 $= \sqrt[6]{a^2b^2d^3} = \sqrt[6]{aabb^2d^3}$.

Ex. 4.

Multiply $a^{\frac{2}{3}}$ by $a^{\frac{1}{4}}$.Product $a^{\frac{2}{3}} \times a^{\frac{1}{4}} = a^{\frac{2}{3} + \frac{1}{4}} = a^{\frac{11}{12}}$.

2 R U L E.

A surd by a rational quantity; connect them with the sign (\times) of multiplication; or else reduce the rational quantity to the form of that surd, and multiply by Rule 1.

Ex. 5.

Multiply $\sqrt{4a-3x}$ by $2a$.The product is $2a \times \sqrt{4a-3x}$.

Or $2a = \sqrt{4aa}$, then the product =
 $\sqrt{16a^2-12aax}$.

3 R U L E.

When rational quantities are annexed to surds; multiply the rational by the rational, and the surd by the surd.

Ex. 6.

Multiply $\frac{a}{2b} \sqrt{a-x}$ by $\frac{c-d}{c-d} \sqrt{ax}$.

The product = $\frac{a \times c-d}{2b} \sqrt{a-x \times ax} =$

$$\frac{ac-ad}{2b} \sqrt{aax-axx}.$$

F

Ex.

Ex. 7.

Multiply $\frac{a}{b}\sqrt{ax}$ by $\sqrt{b-x}\sqrt{\frac{ax^3}{b}}$.

Here $\frac{a}{b}\sqrt{ax} = \frac{a}{b} \times ax^{\frac{1}{2}} = \frac{a}{b} \times \overline{ax^3}^{\frac{1}{6}}$.

And $\sqrt{b-x}\sqrt{\frac{ax^3}{b}} = \sqrt{b-x} \times \left(\frac{ax^3}{b}\right)^{\frac{1}{2}} =$

$\sqrt{b-x} \times \frac{\overline{aax^6}}{bb}^{\frac{1}{2}}$. Therefore

$\frac{a}{b} \times \overline{ax^3}^{\frac{1}{2}} \times$ into $\sqrt{b-x} \times \frac{\overline{aax^6}}{bb}^{\frac{1}{2}} =$

$\frac{ab-ax}{b} \times \frac{\overline{a^3x^9}}{bb}^{\frac{1}{2}} = \frac{ab-ax}{b} \sqrt{\frac{a^3x^9}{bb}} =$

$\frac{ab-ax}{b} \times \sqrt{\frac{a^3x^9}{bb}}$ the product.

Ex. 8:

Multiply $a + \sqrt{b-d}$
by $a - \sqrt{b}$

$$\begin{array}{r} aa + a\sqrt{b-d} \\ - a\sqrt{b-d} + d\sqrt{b} \\ \hline \end{array}$$

product $aa - ad - b + d\sqrt{b}$

Ex. 9.

Multiply $2a - 3a\sqrt{d}$
by $3c - 2c\sqrt{d}$

$$\begin{array}{r} 6ac - 9ac\sqrt{d} \\ - 4ac\sqrt{d} + 6acd \\ \hline \end{array}$$

product $6ac - 13ac\sqrt{d} + 6acd$

Ex.

Ex. 10.

Multiply $\sqrt{a - \sqrt{b - \sqrt{3}}}$
 by $\sqrt{a + \sqrt{b - \sqrt{3}}}$

product $\sqrt{aa - a\sqrt{b - \sqrt{3}} + a\sqrt{b - \sqrt{3}} - b + \sqrt{3}}$

or $\sqrt{aa - b + \sqrt{3}}$.

Schol. If impossible or imaginary roots be multiplied together, they always produce —, otherwise a real product would be raised from impossible factors, which is absurd. Thus,

$\sqrt{-a} \times \sqrt{-b} = \sqrt{-ab}$, and $\sqrt{-a} \times -\sqrt{-b} = -\sqrt{-ab}$, &c. Also $\sqrt{-a} \times \sqrt{-a} = -a$, and $\sqrt{-a} \times -\sqrt{-a} = +a$, &c.

PROBLEM XXVIII.

To divide surds.

I R U L E.

In surds of the same simple quantity; subtract their indices from each other.

Ex. 1.

Divide $a^{\frac{2}{3}}$ by $a^{\frac{1}{3}}$.

quotient $a^{\frac{2}{3} - \frac{1}{3}} = a^{\frac{1}{3}}$.

Ex. 2.

Divide $a^{\frac{1}{n}}$ by $\frac{1}{m}$.

quotient $a^{\frac{1}{n} - \frac{1}{m}} = a^{\frac{m-n}{mn}}$.

F 2

2 RULE.

2 R U L E.

If they be different quantities; reduce them to the same index, if they are not so already. Then divide the quantities under the common index.

Ex. 3.

Divide $\sqrt{15}$ by $\sqrt{5}$.

$5 \sqrt{3}$ the quotient.

Ex. 4.

Divide $\sqrt{\frac{3aad}{c}}$, by $\sqrt{\frac{2ab}{3c}}$.

$\frac{2ab}{3c} \sqrt{\frac{3aad}{c}}$ $\left(\sqrt{\frac{9ad}{2b}} \right)$, quotient.

Ex. 5.

Divide $\sqrt[6]{aabb^3}$ by \sqrt{d} .

$\sqrt{d} = \sqrt[6]{d^3}$. $d^3 aabb^3$ $\left(\sqrt[6]{aabb} = \sqrt[3]{ab} \right)$, quot.

3 R U L E.

If rational quantities are annexed; divide rational quantities by rational quantities, and surds by surds.

Ex. 6.

Divide $\sqrt{16a^3 - 12aax}$ by $2a$.

quotient $\frac{1}{2a} \sqrt{16a^3 - 12aax} = \sqrt{\frac{16a^3 - 12aax}{4aa}} = \sqrt{4a - 3x}$.

Ex.

Ex. 7.

Divide $\frac{ac-ad}{2b} \sqrt{aax-axx}$ by $\frac{a}{2b} \sqrt{a-x}$.

$$\frac{a}{2b} \left) \frac{ac-ad}{2b} \left(\frac{c-d}{1} \cdot a-x \right) aax-axx (\sqrt{ax}.$$

Then $\overline{c-d} \times \sqrt{ax} =$ quotient.

Ex. 8.

Divide $\frac{ab-ax}{b} x \sqrt[6]{\frac{a^5x^3}{bb}}$ by $\overline{b-x} \sqrt[3]{\frac{ax^3}{b}}$.

$$\sqrt[3]{\frac{ax^3}{b}} = \sqrt[6]{\frac{aax^6}{b^2}}.$$

$$b-x \left) \frac{ab-ax}{b} x \left(\frac{ax}{b} \cdot \frac{aax^6}{b^2} \right) \frac{a^5x^3}{bb} \left(\frac{a^3}{x^3} \right).$$

Then the quotient

$$= \frac{ax}{b} \sqrt[6]{\frac{a^3}{x^3}} = \frac{ax}{b} \sqrt{\frac{a}{x}} = \frac{a}{b} \sqrt{ax}.$$

Ex. 9.

$$\begin{array}{r} a-\sqrt{b} \quad aa-ad-b+d\sqrt{b} \quad (a+\sqrt{b}-d \\ \underline{aa-a\sqrt{b}} \qquad \qquad \qquad \text{quotient.} \end{array}$$

$$+a\sqrt{b}$$

$$+a\sqrt{b}-b$$

○

○

$$-ad$$

$$\underline{-ad+d\sqrt{b}}$$

○

Ex. 10.

$$\begin{array}{r}
 aa + a\sqrt{bc} \quad a^2b - abbc \quad (ab - b\sqrt{bc}) \\
 a^2b + a^2b\sqrt{bc} \\
 \hline
 -a^2b\sqrt{bc} - abbc \\
 -a^2b\sqrt{bc} - abbc \\
 \hline
 0
 \end{array}$$

Ex. 11.

Divide $\sqrt{aa - b + \sqrt{3}}$ by $\sqrt{a - \sqrt{b} - \sqrt{3}}$.

$$\begin{array}{r}
 a - \sqrt{b} - \sqrt{3} \quad) \quad aa - b + \sqrt{3} \quad (\sqrt{a + \sqrt{b} - \sqrt{3}} \\
 aa - a\sqrt{b} - \sqrt{3} \\
 \hline
 +a\sqrt{b} - \sqrt{3} - b + \sqrt{3} \\
 +a\sqrt{b} - \sqrt{3} - b + \sqrt{3} \\
 \hline
 0
 \end{array}$$

Ex. 12.

Divide $\sqrt{bbca + \sqrt{aab} - bc} - \sqrt{abc}$
by $\sqrt{bc + \sqrt{a}}$:

$$\begin{array}{r}
 \sqrt{bc + \sqrt{a}} \quad) \quad \sqrt{bbca + \sqrt{aab} - bc} - \sqrt{abc} \quad (\sqrt{ba} - \sqrt{bc} \\
 \sqrt{bbca + \sqrt{aab}} \\
 \hline
 0 \quad -bc - \sqrt{abc} \\
 -bc - \sqrt{abc} \\
 \hline
 0
 \end{array}$$

4 R U L E.

When the quantities will not divide, set them down in form of a fraction.

Ex.

Ex. 13.

Divide $\sqrt{bcd} + \sqrt{abb}$: by $\sqrt{ab} - \sqrt{4bc}$

The quotient is $\frac{\sqrt{bcd} + \sqrt{abb}}{\sqrt{ab} - \sqrt{4bc}}$

PROBLEM XXIX.

To involve surd quantities to any power.

R U L E.

Multiply the index of the quantity, by the index of the power to be raised.

Ex. 1.

Let $\sqrt{2}$ be cubed.

$\sqrt{2} = 2^{\frac{1}{2}}$. Then $2^{\frac{1}{2} \times 3}$ or $2^{\frac{3}{2}}$ is the cube, that is $\sqrt{2^3}$ or $\sqrt{8} =$ the cube of $\sqrt{2}$.

Ex. 2.

What is the square of $3\sqrt[3]{bcc}$.

$3\sqrt[3]{bcc} = 3 \times \overline{bcc^{\frac{1}{3}}}$. Its square =
 $9 \times \overline{bcc^{\frac{2}{3}}} = 9\sqrt[3]{bbc^2} = 9c\sqrt[3]{bbc}$.

Ex. 3.

What is the cube of $a\sqrt{a-x}$.

$a\sqrt{a-x} = a^1 \times \overline{a-x^{\frac{1}{2}}}$; cubed it is $a^3 \times \overline{a-x^{\frac{3}{2}}}$
 that is, the cube = $a^3\sqrt{a^3-3a^2x+3ax^2-x^3}$.

Ex. 4.

What is the 4th power of $\frac{a}{2b}\sqrt{\frac{2a}{c-b}}$.

Here $\frac{a}{2b}\sqrt{\frac{2a}{c-b}} = \frac{a}{2b} \times \sqrt{\frac{2a}{c-b}}^{\frac{1}{2}}$. And its 4th power is $\left(\frac{a}{2b}\right)^4 \times \left(\frac{2a}{c-b}\right)^{\frac{4}{2}} = \frac{a^4}{16b^4} \times \left(\frac{2a}{c-b}\right)^2 = \frac{4a^6}{16b^4 \times c-b^2} = \frac{a^6}{4b^4 \times cc - 2bc + bb^2}$.

2 R U L E.

If quantities are to be involved to a power denoted by the index of the surd root; take away the radical sign.

Ex. 5.

Let $\sqrt{\frac{4ab}{cc}}$ be squared.

Its square is $\frac{4ab}{cc}$.

Ex. 6.

What is the cube of $\sqrt[3]{a^3-b^3+3b\sqrt{abb}}$.

Answer, $a^3-b^3+3b\sqrt{abb}$.

3 R U L E.

Compound surds are involved as integers, observing the rule of multiplication of surds.

Ex. 7.

Let $3+\sqrt{5}$ be squared.

$$\begin{array}{r} 3+\sqrt{5} \\ 3+\sqrt{5} \\ \hline 9+3\sqrt{5} \\ +3\sqrt{5}+5 \\ \hline 14+6\sqrt{5} \end{array} \text{ Ex.}$$

the square

Ex. 8.

Let $a - \sqrt{b}$ be cubed.

$$\begin{array}{r}
 a - \sqrt{b} \\
 a - \sqrt{b} \\
 \hline
 aa - a\sqrt{b} \\
 \quad - a\sqrt{b} + b \\
 \hline
 aa - 2a\sqrt{b} + b \\
 a - \sqrt{b} \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 a^3 - 2aa\sqrt{b} + ab \\
 \quad - aa\sqrt{b} + 2ab - b\sqrt{b} \\
 \hline
 \end{array}$$

the cube

$$\begin{array}{r}
 a^3 - 3aa\sqrt{b} + 3ab - b\sqrt{b}. \\
 \hline
 \end{array}$$

P R O B L E M XXX.

To extract any root of a surd.

R U L E.

Divide the index of the quantity or quantities,
by the index of the root to be extracted.

Ex. 1.

Extract the square root of a^3 .

$$\text{The root} = a^{\frac{3}{2}} = \sqrt{a^3}.$$

Ex. 2.

Extract the cube root of ab^2 .

$$\text{The root is } a^{\frac{1}{3}}b^{\frac{2}{3}} = \sqrt[3]{abb}.$$

Ex. 3.

What is the 4th root of $3aa$.

$$\text{The root is } a^{\frac{2}{4}}\sqrt[4]{3} = a^{\frac{1}{2}}\sqrt[4]{3} = \sqrt{a} \times \sqrt[4]{3}.$$

Ex.

Ex. 4.

What is the cube root of $\sqrt{aa-xx}$.

The root is $\overline{aa-xx}^{\frac{1}{2} \times \frac{2}{3}} = \overline{aa-xx}^{\frac{1}{3}} = \sqrt[3]{aa-xx}$.

2 R U L E.

When the index of the root to be extracted, is the same as the index of the power of that quantity; take away that index, and the quantity itself is the root.

Ex. 5.

What is the square root of $3^2 a^2$.

Answ. $3a$, the root.

Ex. 6.

What is the cube root of $\overline{5ax-3xx}^3$

Answ. $5ax-3xx$, the root.

3 R U L E.

Compound surds are extracted as integers, due regard being had to the operations of simple surds. When no such root can be found, prefix the radical sign.

Ex. 7.

For the square root of $aa-4a\sqrt{b}+4b$.

$$\begin{array}{r}
 aa-4a\sqrt{b}+4b \quad (a-2\sqrt{b}) \\
 aa \\
 \hline
 2a-2\sqrt{b}) \circ -4a\sqrt{b}+4b \\
 \quad \quad \quad -4a\sqrt{b}+4b \\
 \quad \quad \quad \hline
 \quad \quad \quad 0
 \end{array}$$

Ex.

Ex. 8.

What is the cube root of $aa - \sqrt{ax - xx}$.

Answ. $\sqrt[3]{ax - \sqrt{ax - xx}}$, the root.

PROBLEM XXXI.

To change a binomial surd quantity into another.

RULE.

This reduction is performed by an equal involution, and evolution. Involve the binomial to the power denoted by the surd or surds, then set the radical sign of the same root before it.

Ex. 1.

To transform $2 + \sqrt{3}$ to another.

Its square, $4 + 3 + 4\sqrt{3} = 7 + 4\sqrt{3}$

the square root, $\sqrt{7 + 4\sqrt{3}}$.

Ex. 2.

Reduce $\sqrt{2 + \sqrt{3}}$ to a universal surd.

Its square $2 + 3 + 2\sqrt{6} = 5 + 2\sqrt{6}$

the root $\sqrt{5 + 2\sqrt{6}}$.

Ex. 3.

Let $\sqrt{a - 2\sqrt{x}}$ be given to reduce.

The square $a + 4x - 4\sqrt{ax}$

the root $\sqrt{a + 4x - 4\sqrt{ax}}$.

Ex. 4.

Let $\sqrt[3]{a} + \sqrt[3]{b}$ be given.

The cube $a + 3\sqrt[3]{aa^2} + 3\sqrt[3]{abb} + b$

the root $\sqrt[3]{a + 3\sqrt[3]{aa^2} + 3\sqrt[3]{abb} + b}$.

Cor.

Cor. $\sqrt{a} + \sqrt{b} = \sqrt{\frac{1}{2}(\sqrt{a} + \sqrt{b})^2}$; and in general $a^{\frac{1}{n}} + b^{\frac{1}{n}} = \sqrt[n]{\frac{1}{2}(a^{\frac{1}{n}} + b^{\frac{1}{n}})^n}$.

P R O B L E M XXXII.

To extract the square root of a binomial (or residual) surd, $A+B$, or $A-B$; or a trinomial, &c.

R U L E, for binomials.

Take $\sqrt{AA - BB} = D$. Then $\sqrt{A+B} = \sqrt{\frac{A+D}{2}} + \sqrt{\frac{A-D}{2}}$.

and $\sqrt{A-B} = \sqrt{\frac{A+D}{2}} - \sqrt{\frac{A-D}{2}}$

For if $\sqrt{\frac{A+D}{2}} + \sqrt{\frac{A-D}{2}}$ be involved by Prob. 29. it will produce $A + \sqrt{AA - DD}$, that is $A+B$, as it ought. And $\sqrt{\frac{A+D}{2}} - \sqrt{\frac{A-D}{2}}$ will also produce $A-B$.

Ex. I.

To extract the root of $7 + \sqrt{20}$.

Here $A=7$, $B=\sqrt{20}$, and $\sqrt{AA - BB} = \sqrt{29} = D$.

Then the square root of $7 + \sqrt{20} = \sqrt{\frac{7+\sqrt{29}}{2}} + \sqrt{\frac{7-\sqrt{29}}{2}}$.

Ex.

Ex. 2.

What is the square root of $3-2\sqrt{2}$.

Here $\sqrt{AA-BB} = \sqrt{1} = 1 = D$, and
 $\frac{A+D}{2} = 2$, $\frac{A-D}{2} = 1$. And

$\sqrt{3-2\sqrt{2}} = \sqrt{2-\sqrt{1}} = \sqrt{2-1}$, the root.

Ex. 3.

To extract the root of $27+\sqrt{704}$.

$\sqrt{AA-BB} = \sqrt{25} = D = 5$. And the
 root = $\sqrt{\frac{32}{2}} + \sqrt{\frac{22}{2}}$ that is,

$\sqrt{27+\sqrt{704}} = \sqrt{16+\sqrt{11}} = 4+\sqrt{11}$.

Ex. 4.

What is the square root of $6-2\sqrt{5}$.

Here $\sqrt{AA-BB} = \sqrt{36-20} = D = 4$.
 And $\sqrt{\frac{A+D}{2}} = \sqrt{5}$, and $\sqrt{\frac{A-D}{2}} = 1$.

And the root = $\sqrt{5-1}$.

Ex. 5.

Extract the root of $\sqrt{21+\sqrt{5}}$.

$\sqrt{AA-BB} = \sqrt{16} = D = 4$. And
 $\frac{A+D}{2} = \frac{\sqrt{21+4}}{2}$, $\frac{A-D}{2} = \frac{\sqrt{21-4}}{2}$.

And the root $\sqrt{\frac{\sqrt{21+4}}{2}} + \sqrt{\frac{\sqrt{21-4}}{2}}$.

Ex.

Ex. 6.

Extract the root of $aa+2x\sqrt{aa-xx}$.

Here $A = aa$, $B = 2x\sqrt{aa-xx}$. Then
 $\sqrt{AA-BB} = \sqrt{aa-4a^2x^2+4x^4} = aa-2xx = D$.
 Then $\frac{A+D}{2} = aa-xx$, and $\frac{A-D}{2} = xx$, and
 the root $= x + \sqrt{ax-xx}$.

Ex. 7.

What is the root of $6+\sqrt{8}-\sqrt{12}-\sqrt{24}$.

Let $A=6+\sqrt{8}$, $B=\sqrt{12}+\sqrt{24}$. Then
 $\sqrt{AA-BB}=D=\sqrt{44+12\sqrt{8}-36-2\sqrt{12}\times 24}$
 $=\sqrt{8}$. $\frac{A+D}{2} = 3 + \sqrt{8}$, $\frac{A-D}{2} = 3$.

And the root $= \sqrt{3+\sqrt{8}} - \sqrt{3}$. But
 $\sqrt{3+\sqrt{8}} = 1 + \sqrt{2}$, (see Ex. 2.); there-
 fore the root $= 1 + \sqrt{2} - \sqrt{3}$.

2 R U L E, for trinomials, &c.

For trinomial, quadrinomial surds, &c. divide half the product of any two radicals by a third, gives the square of one radical part of the root. This repeated with different quantities, will give the squares of all the parts of the root, to be connected by + and -. But if any quantity occur oftener than once; it must be taken but once.

For if $x+y+z$ be any trinomial surd, its square will be $x^2+y^2+z^2+2xy+2xz+2yz$; then if half the product of any two rectangles as $2xy \times 2xz$ (or $2x^2yz$) be divided by some third $2yz$, the quotient $\frac{2x^2yz}{2yz} = xx$, must needs be the square of one of the parts; and the like for the rest.

Ex.

Ex. 8.

To extract the square root of

$$6 + \sqrt{8} - \sqrt{12} - \sqrt{24}.$$

Here $\frac{\sqrt{8} \times \sqrt{12}}{2\sqrt{24}} = 1$, and $\frac{\sqrt{8} \times \sqrt{24}}{2\sqrt{12}} = \sqrt{4} = 2$,and $\frac{\sqrt{12} \times \sqrt{24}}{2\sqrt{8}} = \sqrt{9} = 3$. And the root is $1 + \sqrt{2} - \sqrt{3}$.

Ex. 9.

To find the square root of

$$12 + \sqrt{32} - \sqrt{48} + \sqrt{80} - \sqrt{24} + \sqrt{40} - \sqrt{60}.$$

Here $\frac{\sqrt{32} \times 48}{2\sqrt{80}} = \sqrt{\frac{24}{5}}$, this produces no-thing. Again, $\frac{\sqrt{32} \times 48}{2\sqrt{24}} = \sqrt{16} = 4$. And

$$\frac{\sqrt{40} \times 60}{2\sqrt{4}} = \sqrt{25} = 5; \text{ and } \frac{\sqrt{24} \times 40}{2\sqrt{60}} = \sqrt{4} = 2;$$

$$\text{and } \frac{\sqrt{48} \times 24}{2\sqrt{32}} = \sqrt{9} = 3; \text{ and } \frac{\sqrt{32} \times 80}{2\sqrt{40}} = \sqrt{16} = 4,$$

&c. therefore the parts of the root are $\sqrt{4}$, $\sqrt{5}$, $\sqrt{3}$, $\sqrt{2}$, $\sqrt{4}$, &c. and the root $2 + \sqrt{2} - \sqrt{3} + \sqrt{5}$; for being squared it produces the surd quantity given.

Cor. 1. In binomials, if D be a rational quantity, the root will consist of two surds, and the parts of each under the radical sign will consist of a rational quantity (D), and a surd (A).

Cor. 2. If both A and D be rational, the root will consist either of the two surds, or else of a rational part and a surd; which is the only case that is useful in this extraction.

P R O B L E M XXXIII.

To extract any root (c) of a binomial surd $A+B$,
or $A-B$.

R U L E.

Let $AA-BB=D$, take Q such, that $QD=n^c$,

the least integer power. Let $\sqrt[A+B]{\times\sqrt{Q}}=r$,
the nearest integer number.

Reduce $A\sqrt{Q}$ to the simplest form $p\sqrt{s}$.

Let $\frac{r+\frac{n}{r}}{2\sqrt{s}} = t$, the nearest integer.

Then the root = $\frac{t\sqrt{s} + \sqrt{tts-n}}{\sqrt{Q}}$, if it can be
extracted.

Note, + is for the binomial $A+B$, and -
for the residual $A-B$.

Ex. I.

What is the cube root of $\sqrt{968+25}$.

Here $D = 343 = 7 \times 7 \times 7$. $Q \times 7^3 = n^3$, and

$Q=1$, $n=7$. Then $\sqrt[A+B]{\times\sqrt{Q}} = \sqrt[3]{56} +$
 $= r = 4$. $A\sqrt{Q} = \sqrt{968} = 22\sqrt{2} = p\sqrt{s}$, and

$\sqrt{s} = \sqrt{2}$. $\frac{r+\frac{n}{r}}{2\sqrt{s}} = \frac{4+\frac{7}{4}}{2\sqrt{2}} = t = 2$. And

$t\sqrt{s} = 2\sqrt{2}$, $\sqrt{tts-n} = \sqrt{8-7} = 1$. $\sqrt[6]{Q} = 1$.

And the root $\frac{2\sqrt{2}+1}{1} = 2\sqrt{2}+1$, which
succeeds.

Ex.

Ex. 2.

Extract the cube root of $68 - \sqrt{4374}$.

Here $D = 250 = 5 \times 5 \times 5 \times 2$. And $5^3 \times 2^3 = 4D$
 $= QD = n^3$, and $Q = 4$, $n = 2 \times 5 = 10$. And

$$\sqrt[3]{A+B} \times \sqrt{Q} = \sqrt[3]{134 \times 2} = 6 = r.$$

$$A\sqrt{Q} = 136\sqrt{1} = p\sqrt{s}, \text{ and } \sqrt{s} = 1.$$

$$\frac{r + \frac{n}{r}}{2\sqrt{s}} = \frac{7\frac{2}{3}}{2} = \frac{23}{6} = 4 = t. \quad t\sqrt{s} = 4.$$

And the root

$$\frac{t\sqrt{s} - \sqrt{tts - n}}{\sqrt{2}} = \frac{-\sqrt{10 - 10}}{\sqrt{4}} = \frac{4 - \sqrt{6}}{\sqrt{2}}$$

for its cube is $68 - 27\sqrt{6}$.

Ex. 3.

Extract the 5th root of $23\sqrt{6} + 41\sqrt{3}$.

Here $D = 3$, $n = 3$, $Q = 81$, $r = 5$, $\sqrt{s} = \sqrt{6}$,

$$t = 1, \quad t\sqrt{s} = \sqrt{6}, \quad \sqrt{tts - n} = \sqrt{3},$$

$\sqrt{Q} = \sqrt{81} = \sqrt[5]{9}$. And the root to be tried

$$\frac{\sqrt{6} + \sqrt{3}}{\sqrt[5]{9}}.$$

SCHOLIUM.

If the quantity be a fraction or has a common divisor, extract the root of the denominator or of that common divisor, separately. They that would see the demonstration of this rule, may consult *Gravesand's* or *Mac Laurin's Algebra*. For as it seldom happens that such quantities have a proper root; it is not worth while spending any more time about them.

PROBLEM XXXIV.

A compound surd being given, consisting of two, three, or more terms, which are surd square roots: to find such a multiplier or multipliers, by which multiplying the given surd; the product will be rational.

R U L E.

Change the sign of one of the terms in a binomial, or trinomial, or the signs of two terms in a quadrinomial; and by this multiply the given surd.

Ex. 1.

Let $a + \sqrt{3}$ be given.

$$\begin{array}{r} \text{Multiply by } a - \sqrt{3} \\ \hline \text{product } aa - 3. \end{array}$$

Ex. 2.

Given $\sqrt{5} - \sqrt{x}$.

$$\begin{array}{r} \text{Multiply by } \sqrt{5} + \sqrt{x} \\ \hline \text{product } 5 - x \text{ rational.} \end{array}$$

Ex. 3.

Let $\sqrt{5} + \sqrt{3} - \sqrt{2}$ be given.

$$\begin{array}{r} \text{Multiply by } \sqrt{5} + \sqrt{3} + \sqrt{2} \\ \hline 5 + \sqrt{12} - \sqrt{10} \\ + \sqrt{15} + 3 - \sqrt{6} \\ + \sqrt{10} + \sqrt{6} - 2 \\ \hline \text{product } 6 + 2\sqrt{15} \\ \text{multiply by } -6 + 2\sqrt{15} \\ \hline \text{product } 60 - 36 = 24. \end{array}$$

Ex.

Ex. 4.

There is given $\sqrt{a} + \sqrt{b} - \sqrt{c} + \sqrt{d}$
 Multiply by $\sqrt{a} + \sqrt{b} + \sqrt{c} - \sqrt{d}$
 product $\frac{a+b-c-d+2\sqrt{ab}+2\sqrt{dc}}{}$

or
 multiply by $\frac{f+2\sqrt{ab}+2\sqrt{dc}}{f+2\sqrt{ab}-2\sqrt{dc}}$
 product $\frac{ff+4f\sqrt{ab}+4ab-4dc}{}$

or
 multiply by $\frac{g+4f\sqrt{ab}}{g-4f\sqrt{ab}}$
 product $\frac{gg-16ffab}{}$

In this process f is put for the rational part $a+b-c-d$; and g for $ff+4ab-4dc$.

Cor. A binomial becomes rational after one operation, a trinomial after two, and a quadrinomial after three, &c.

P R O B L E M XXXV.

A binomial being given, consisting of one or two surds, whose index or root is any power of 2; to find a multiplier or multipliers that shall make it rational.

R U L E.

Multiply it by its corresponding residual (that is when one sign is changed); and repeat the same operation, as long as there are surds.

Ex. 1.

Let $\sqrt{a} - \sqrt{b}$ be given.

Multiply by $\sqrt{a} + \sqrt{b}$
 product $\frac{a-b}{}$ rational.

G 2

Ex:

Ex. 2.

Let $\sqrt[4]{5} + \sqrt[4]{3}$ be proposed.Multiply by $\sqrt[4]{5} - \sqrt[4]{3}$

1 product	$\sqrt[4]{5} - \sqrt[4]{3}$
multiply by	$\sqrt[4]{5} + \sqrt[4]{3}$

2 product	$5 - 3 = 2$, rational.
-----------	-------------	-------------

Ex. 3.

Let there be given $\sqrt[8]{a} + \sqrt[8]{b}$.

Multiply by	$\sqrt[8]{a} - \sqrt[8]{b}$
-------------	-----------------------------

1 product	$\sqrt[4]{a} - \sqrt[4]{b}$
multiply by	$\sqrt[4]{a} + \sqrt[4]{b}$

2 product	$\sqrt{a} - \sqrt{b}$
mult. by	$\sqrt{a} + \sqrt{b}$

3 product	$a - b$	rational.
-----------	---------	-----------

Ex. 4.

Let $a + \sqrt[4]{b}$ be given.

Multiplier	$a - \sqrt[4]{b}$
------------	-------------------

1 prod.	$aa - \sqrt{b}$
mult.	$aa + \sqrt{b}$

2 prod.	$a^2 - b$
---------	-----------

Cor. The number of operations, is equal to the power of 2 in the index.

P R O B.

PROBLEM XXXVI.

Any binomial surd being given, to find a multiplier which shall produce a rational product.

R U L E.

If the surds have not the same index, reduce them to the same, (Prob. 21.)

Take the two quantities (throwing away the radical sign or index); change the sign of one of them. That done, involve these to the next inferior power denoted by the index of the root (Prob. 5. Rule 3), but leave out the unciæ or coefficients: then place the common radical sign before each quantity, but after its sign. And this will be your multiplier.

Shorter thus,

$$\text{Binomial } \sqrt[n]{A} \pm \sqrt[n]{B}.$$

$$\text{Multiplier } \sqrt[n]{A^{n-1}} \mp \sqrt[n]{A^{n-2}B} + \sqrt[n]{A^{n-3}B^2} \\ + \sqrt[n]{A^{n-4}B^3} + \&c.$$

The upper signs must be taken with the upper, and the lower with the lower; and the series continued to n terms.

Ex. I.

Let $\sqrt[3]{7} + \sqrt[3]{3}$ be given.

$$\text{Multiplier } \sqrt[3]{7 \times 7} - \sqrt[3]{7 \times 3} + \sqrt[3]{3 \times 3}$$

$$7 + \sqrt[3]{7 \times 7 \times 3}$$

$$- \sqrt[3]{7 \times 7 \times 3} - \sqrt[3]{7 \times 3 \times 3}$$

$$+ \sqrt[3]{7 \times 3 \times 3} + 3$$

$$\text{product } 7 + 3 = 10, \text{ rational.}$$

G 3

Ex.

Ex. 2.

Let $a - \sqrt[3]{2}$ be proposed. $= \sqrt[3]{a^3 - \sqrt[3]{2}}$.

$$\begin{array}{l} \text{Multiplier } \underline{aa + a\sqrt[3]{2} + \sqrt[3]{2} \times 2} \\ \text{product } \underline{\underline{a^3 - 2.}} \end{array}$$

Ex. 3.

Let $\sqrt[3]{a} + \sqrt[3]{b}$ be proposed.

$$\begin{array}{l} \text{Mult. } \underline{\sqrt[3]{aa} - \sqrt[3]{ab} + \sqrt[3]{bb}} \\ \text{product } \underline{\underline{a \qquad \qquad \qquad + b}} \end{array}$$

Ex. 4.

Let $\sqrt[4]{5} + \sqrt[4]{3}$ be given.reduced $\sqrt[4]{5} + \sqrt[4]{9}$, given.

$$\begin{array}{l} \text{Multiplier } \underline{\sqrt[4]{5^3} - \sqrt[4]{5^2 \times 9} + \sqrt[4]{5 \times 9^2} - \sqrt[4]{9^3}} \\ \text{product } \underline{\underline{5 \qquad \qquad \qquad - 9 = -4.}} \end{array}$$

Or thus,

$$\begin{array}{l} \text{Surd } \sqrt[4]{9} + \sqrt[4]{5}. \\ \text{mult. } \underline{\sqrt[4]{9^3} - \sqrt[4]{9^2 \times 5} + \sqrt[4]{9 \times 5^2} - \sqrt[4]{5^3}} \\ \text{product } \underline{\underline{9 \qquad - 5 \qquad = 4.}} \end{array}$$

Ex. 5:

Let $\sqrt[4]{a^3} - \sqrt[4]{b^3}$ be given.

$$\text{Multiplier } \sqrt[4]{a^9} + \sqrt[4]{a^6 b^3} + \sqrt[4]{a^3 b^6} + \sqrt[4]{b^9}.$$

Or

Or thus,

$$\begin{array}{l} \text{Surd} \quad \sqrt[4]{a^3} - \sqrt[4]{b^3}. \\ \text{mult.} \quad aa\sqrt[4]{a} + a\sqrt[4]{aab^2} + b\sqrt[4]{a^2bb} + bb\sqrt[4]{b} \\ \hline \text{product} \quad a^3 - b^3. \end{array}$$

Ex. 6.

Let $\sqrt[3]{a} - \sqrt[3]{b}$ be proposed.

reduced to $\sqrt[6]{a^3} - \sqrt[6]{bb}$
 put $x = a^3, y = bb$.

$$\begin{array}{l} \text{Surd} \quad \sqrt[6]{x} - \sqrt[6]{y} \\ \text{mult.} \quad \sqrt[6]{x^5} + \sqrt[6]{x^4y} + \sqrt[6]{x^3y^2} + \sqrt[6]{x^2y^3} + \sqrt[6]{xy^4} + \sqrt[6]{y^5}. \\ \hline \text{product} \quad x - y = a^3 - bb. \end{array}$$

PROBLEM XXXVII.

A fraction being given whose denominator is a compound surd; to reduce it to another whose denominator is rational.

R U L E.

Find such a multiplier (by Prob. 34, 35, or 36), as will make the denominator rational. By this multiply both numerator and denominator.

Ex. 1.

Let $\frac{3}{\sqrt{5} - \sqrt{2}}$ be proposed.

Here $\frac{3 \times \sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2} \times \sqrt{5} + \sqrt{2}} =$
 $\frac{3\sqrt{5} + 3\sqrt{2}}{5 - 2 = 3} = \sqrt{5} + \sqrt{2}.$

G 4

Ex.

Ex. 2.

Let there be given $\frac{\sqrt{6}}{\sqrt{7+\sqrt{3}}}$.

Multiply both terms by $\sqrt{7-\sqrt{3}}$, the fraction becomes $\frac{\sqrt{42}-\sqrt{18}}{7-3=4}$.

Ex. 3.

Suppose $\frac{\sqrt{2}}{3-\sqrt{2}}$.

Multiply by $3+\sqrt{2}$, then $\frac{3\sqrt{2+2}}{9-2=7}$ is the fraction required.

Ex. 4.

Let $\frac{ab-b\sqrt{bc}}{a+\sqrt{bc}}$ be proposed.

Multiply by $a-\sqrt{bc}$, then $\frac{acb-2ab\sqrt{bc}+b^2c}{aa-bc}$ is the fraction sought.

Ex. 5.

Let $\frac{3\sqrt{a+2\sqrt{b}}}{5-\sqrt{3}}$ be given.

Multiply by $5+\sqrt{3}$; then $\frac{15\sqrt{a+2\sqrt{b}}+3\sqrt{3a+2\sqrt{3b}}}{25-3=22}$.

Ex. 6.

Suppose $\frac{10}{\sqrt{7-\sqrt{5}}}$.

Multiply by $\sqrt[3]{7^2}+\sqrt[3]{7}\times\sqrt[3]{5}+\sqrt[3]{5^2}$, and the fraction becomes $\frac{10\sqrt[3]{49}+10\sqrt[3]{35}+10\sqrt[3]{25}}{7-5} =$
 $5\sqrt[3]{49}+5\sqrt[3]{35}+5\sqrt[3]{25}$. Ex.

Ex. 7.

$$\text{Let } \frac{\sqrt{ab}}{\sqrt{5} + \sqrt{3}}$$

Multiply by $\sqrt{5} - \sqrt{5^2 \cdot 3} + \sqrt{5 \cdot 3^2} - \sqrt{3^3}$,

And the fraction is

$$\frac{\sqrt{125} - \sqrt{75} + \sqrt{45} - \sqrt{27}}{5 - 3 = 2} \sqrt{ab}.$$

Or thus,

Multiply the terms of the fraction $\frac{\sqrt{ab}}{\sqrt{5} + \sqrt{3}}$

by $\sqrt{5} - \sqrt{3}$, and it becomes $\frac{\sqrt{5} - \sqrt{3}}{\sqrt{5} - \sqrt{3}} \sqrt{ab}$;

again multiply the terms of the last fraction by $\sqrt{5} + \sqrt{3}$, and it becomes

$$\frac{5^{\frac{3}{2}} - 5^{\frac{1}{2}} 3^{\frac{1}{2}} + 3^{\frac{1}{2}} 5^{\frac{1}{2}} - 3^{\frac{3}{2}}}{5 - 3 = 2} \sqrt{ab}.$$

Ex. 8.

Let $\frac{8}{\sqrt{3} + \sqrt{2} + 1}$ be the fraction.

Multiply by $\sqrt{3} + \sqrt{2} - 1$, and the fraction will

$$\text{be } \frac{8\sqrt{3} + 8\sqrt{2} - 8}{5 + 2\sqrt{6} - 1} = \frac{4\sqrt{3} + 4\sqrt{2} - 4}{2 + \sqrt{6}}.$$

Again, multiply by $-2 + \sqrt{6}$, and it becomes

$$\frac{-8\sqrt{3} - 8\sqrt{2} + 8 + 4\sqrt{18} + 4\sqrt{12} - 4\sqrt{6}}{6 - 4 = 2}$$

$$= 4 + 2\sqrt{18} + 2\sqrt{12} - 2\sqrt{6} - 4\sqrt{3} - 4\sqrt{2}$$

$$= 4 + 6\sqrt{2} + 4\sqrt{3} - 2\sqrt{6} - 4\sqrt{3} - 4\sqrt{2}$$

$$= 4 + 2\sqrt{2} - 2\sqrt{6}.$$

S E C T. IV.

Several Methods of managing Equations.

AN Equation is the mutual comparing of two equal quantities, by the help of this character ($=$); the part on the left hand is called the *first side* of the equation; that on the right, the *second side*. And the single quantities are called *terms* of the equation.

An equation is either two ranks of quantities equal to one another, and separated by this mark ($=$); or one rank equal to nothing. And they are to be considered either, as the last conclusion to which we come in the solution of a problem; or as the means whereby we come to it. In the first case, the equation is composed of only one unknown quantity mixed with known ones, and may be called the *final equation*. But those of the last sort involve several unknown quantities; and therefore they are to be so managed and reduced, that out of all the rest there may emerge a new equation, with only one unknown quantity, which is that we seek. And this is to be made as simple as it can, in order to find the value of the unknown quantity.

An equation is named according to the dimension of the highest power of the unknown quantity in it. A *simple equation* is that which contains only the quantity itself; as $a=b-c$. A *quadratic equation*, is when the highest power is a square, as $aa-ba=d$. A *cubic equation*, when the highest power is a cube, as $a^3+ba^2-ca=d$. A *fourth power* when the highest power is such, as $a^4-3a^3+a=d$, &c.

P R O B L E M XXXVIII.

To turn proportional quantities into equations; and equations into proportions.

In the solution of problems, it often happens, that we have several quantities in geometrical proportion, which are to be reduced into an equation; which will be done thus:

R U L E.

Multiply the extremes together for one side of the equation, and the two means for the other side; or the square of the mean, when there are but 3 terms.

On the contrary in a given equation, divide each side into two factors; and make the two factors of one side the two means; and the two factors of the other side, the extremes.

Ex. 1.

If $a:b::c+f:d$. Then $ad=bc+bf$.

Ex. 2.

Let $a+b : a-b :: \frac{c}{d}\sqrt{aa-xx} : \frac{r}{s}$.

Then $\frac{ar+br}{s} = \frac{ca-cb}{d}\sqrt{aa-xx}$.

Ex. 3.

If $ad = bc+bf$. Then $a:b::c+f:d$.

Ex.

Ex. 4.

$$\text{If } \frac{ar+br}{s} = \frac{ca-cb}{d} \sqrt{aa-xx}.$$

$$\text{Then } \frac{a+b}{s} : \frac{ca-cb}{d} :: \sqrt{aa-xx} : r$$

$$\text{or } a-b : a+b :: \frac{r}{s} : \frac{c}{d} \sqrt{aa-xx}.$$

Ex. 5.

$$\text{Let } bc+bd = da-cg.$$

$$\text{Then } 1 : b :: c+d : da-cg.$$

$$\text{or } b : \sqrt{da-cg} :: \sqrt{da-cg} : c+d.$$

P R O B L E M XXXIX.

To reduce an equation.

When a question is brought to an equation, the unknown quantities are generally mixed and entangled with the known ones; and therefore the equation must be so ordered that the unknown quantity may stand clear, on the first side of the equation; and the known ones on the second side. Which is done thus:

1 R U L E.

When any quantity is on both side the equation, throw it out of both.

Ex. 1.

$$\text{If } 3x+6b = 4c-d + 9b.$$

$$\text{Throw out } 6b. \text{ Then } 3x = 4c-d + 3b.$$

2 R U L E.

When the known and unknown quantities are both on one side; transpose any of them to the contrary side, and change its sign.

Ex.

Ex. 2.

If $5x + 3b = rx + bd.$

Then $5x = rx + bd - 3b$

And $5x - rx = bd - 3b.$

For to transpose a quantity with a contrary sign, is the same thing as to add it, or else to subtract it from both sides; therefore the quantities on each side, remain still equal, by Axiom 1. and 2.

3 R U L E.

If there be fractions in the equation, multiply both sides by the denominators.

Ex. 3.

Suppose $\frac{aa}{b} + c - f = \frac{dx}{a}.$

Multiply by b , $aa + cb - fb = \frac{bdx}{a}$

multiply by a , $a^3 + bca - bfa = bdx.$

This process is plain from Axiom 3.

4 R U L E.

When any quantity is multiplied into both sides of the equation, or into the highest term of the unknown quantity; divide the whole equation thereby.

Ex. 4.

If $7ba^3 + bcaa = bcda.$

Divide by ba , $7aa + ca = cd.$

divide by 7, $aa + \frac{ca}{7} = \frac{cd}{7}.$

The truth of this appears by Axiom 4.

5 R U L E.

5 R U L E.

If the unknown quantity is affected with a surd; transpose the rest of the terms; then involve each side according to the index of the surd.

Ex. 5.

$$\text{If } \sqrt{aa-ba} + c = d.$$

$$\text{Then } \sqrt{aa-ba} = d-c.$$

$$\text{squared } aa - ba = dd - 2dc + cc.$$

This process is plain from Axiom 5.

6 R U L E.

When the side containing the unknown quantity is a pure power; or if being affected, it has a rational root: then extract such root on both sides of the equation.

Ex. 6.

$$\text{If } a^3 = b^3 - bbc.$$

$$\text{Cube root. } a = \sqrt[3]{b^3 - bbc}.$$

Ex. 7.

$$\text{If } xx + 6x + 9 = 20b.$$

$$\text{Square root } x + 3 = \pm \sqrt{20b}.$$

$$\text{and } x = \pm \sqrt{20b} - 3.$$

Schol. All these rules are to be used promiscuously, as one has occasion for them; till the equation be duly cleared.

P R O B L E M XL.

To explain the nature and origin of affected equations.

1. Any affected equation may be considered as made up of as many simple equations, as the dimension

Sect. IV. *Nature of EQUATIONS.* 95

mention of the highest power is. Suppose $x=a$,
 $x=b$, and $x=c$, &c. then $x-a=0$, $x-b=0$,
 $x-c=0$. And if all these be multiplied together,
then $\overline{x-a} \times \overline{x-b} \times \overline{x-c} = 0$; that is,

$$\begin{array}{r} x^3 - ax^2 + abx - abc = 0, \text{ a cubic equation,} \\ -b \quad +ac \\ -c \quad +bc \end{array}$$

whose roots are a, b, c .

In like manner, $\overline{x-a} \times \overline{x-b} \times \overline{x-c} \times \overline{x-d} = 0$,
produces a biquadratic equation,

$$\begin{array}{r} x^4 - ax^3 + abx^2 - abcx + abcd = 0, \\ -b \quad +ac \quad -abd \\ -c \quad +bc \quad -acd \\ -d \quad +da \quad -bcd \\ \quad +db \\ \quad +dc \end{array}$$

whose roots are a, b, c, d .

These two equations may be written or de-
noted thus, $x^3 - px^2 + qx - r = 0$, and
 $x^4 - p^2x^3 + qx^2 - rx + s = 0$. And any such e-
quation being found in the solution of a problem;
the business is then to resolve it into its original
compounding simple equations, and so to find the
roots a, b, c , &c. For each of these simple equa-
tions gives one value of x , or one root. And if
any one of these values of x be substituted in the
equation instead of x , all the terms of the equa-
tion will vanish and be $=0$. For since

$\overline{x-a} \times \overline{x-b} \times \overline{x-c}$, &c. $=0$. It is plain, when
one of the factors $\overline{x-a}$ is $=0$, the whole pro-
duct will be $=0$. And of consequence there are
three roots in the cubic equation, and four in the
biquadratic; and in general there are as many
roots, as is the dimension of the highest power in
it, and no more.

2. If it happen that the roots $a, b, c, \&c.$ be equal to one another, then $\overline{x-a^3}$ will be $=0$, or $\overline{x-a^4}$ $=0$, $\&c.$ and $x-a$ is had by evolution, since the given equation is generated by involution.

3. That there are no more roots than these is plain; for if you put any quantity, as f for x , which is equal to none of the roots $a, b, c, \&c.$ Then since neither $f-a, f-b,$ nor $f-c, \&c.$ is 0, their product cannot vanish or be $=0$, but must be some real product; and therefore f is no root of the equation.

4. Since the square root of a negative quantity is impossible; therefore if we have such an equation as this, $xx + aa = 0$, or $xx = -aa$, then $x = \pm\sqrt{-aa}$, which are two impossible roots of that equation. So that a quadratic equation has either two impossible roots or none. And therefore in any equation, there is always an even number of impossible roots; since each quadratic that goes to the compounding it, must have either two or none. Therefore no equation can have an odd number of impossible roots. Hence therefore the number of real roots in a cubic equation, will either be one or three; in a biquadratic, four, two, or none. In a fifth power, 5, 3 or 1; $\&c.$

5. From the foregoing equations it is plain, that the coefficient of the first term (or that of the highest power) is 1. The coefficient of the second term (or next highest power), is the sum of all the roots, $a, b, c, \&c.$ with their signs changed. The coefficient of the third term, the sum of the products of every two of the roots. The coefficient of the fourth term, the sum of the products of every three of them, with contrary signs, $\&c.$ The odd terms having always the same sign, and the even terms a contrary one. And the absolute

absolute number is always the product of all the roots together.

6. Hence it follows, that when the sum of all the negative roots is equal to the sum of all the affirmative, the second term vanishes, and the contrary. And if all the negative rectangles be equal to all the affirmative ones, the third term vanishes. And if all the negative solids be equal to all the affirmative ones, the fourth term vanishes, out of the equation; and so forward.

7. But the roots of equations may be either $+$ or $-$, yet still the same rules hold good. For let the sign of any of them as c be changed into $-c$, that is, let $x+c=0$; then in the cubic equation the second term will be $-a-b+c$; that is, the sum of the roots with a contrary sign; the third term will be $+ab-ac-bc$, that is, the sum of the products of all the roots; and so of the rest.

8. Hence also in every equation cleared of fractions and surds, each of the roots, each of the rectangles of any two of the roots, each of the solids under any three of them, each of the products of any four of the said roots, &c. are all of them just divisors of the last term or absolute number. Therefore when no such divisor can be found, it is evident there is no root, no rectangle of roots, no solid of roots, &c. but what is surd. For in the cubic equation, a , b , c , and ab , ac , bc , are all of them divisors of the last term abc : and so of higher powers.

9. In any equation, change the signs of all the terms but the first; then let the coefficients of the first, second, third, &c. terms be 1 , p , q , r , s , t , v , &c. respectively.

Then observing the signs, we shall have

$p = \text{sum of the roots, } a+b+c, \&c.$

$pA+2q = \text{sum of the squares of the roots}$
 $= a^2+b^2 \&c.$

$pB+qA+3r = \text{the sum of their cubes}$
 $= a^3+b^3 \&c.$

$pC+qB+rA+4s = \text{the sum of the biquadrates,}$
 $\&c.$

Where A, B, C, $\&c.$ are the first, second, third,
 $\&c.$ terms.

For $+p=a+b+c \&c. = A.$

Also pA or $\overline{a+b+c}^2 = a^2+b^2+c^2+2ab +$
 $2ac+2cd=B-2q.$ Therefore $B=pA+2q, \&c.$

To go through the calculations of the rest would
 be tedious, and of little use.

10. In equations of the third and fourth power, we find, when the roots are all affirmative, the signs are + and - alternately; so that there are as many changes of the signs as is the index of the power, or as the number of roots. But if the roots are all negative, the signs are all + throughout, there being no changes of the signs. Whence in these cases, there are as many affirmative roots, as changes of the signs in all the terms, from + to -, and from - to +. And the same rule holds in general, that is, there are as many affirmative roots in any equation as there are changes of the signs. But the equation is supposed to be compleat, that is to want no terms, and to have numeral coefficients. And likewise the number of negative roots is known thus; as often as two of the signs +, or two of the signs - stand next one another, so often there is a negative root. It would be needless to trouble the reader with the proof of these things; since it can only be done in particular cases, and not in a general way.
 And

Se^t. IV. *Nature of EQUATIONS.* 99

And besides when impossible roots happen to lie hid in the equation, they cause the rule to fail.

11. When the roots are all affirmative, the terms of the equation are alternately + and — through the equation; but when the roots are all negative, the signs are all +; and therefore, as by changing the signs of the roots, the signs of the alternate terms are changed; so on the contrary, changing the signs of the alternate terms, changes the signs of all the roots. And this holds in general, as will be evident by producing two equations from the same roots, with contrary signs.

12. Since any adfected equation, as $x^3 - px^2 + qx - r = 0$, is made up of simple equations, such as $x - a = 0$, $x - b = 0$, &c. Therefore if one root as a be known, the whole equation may be exactly divided by $x - a$; and so reduced to a lower dimension. Also when all the roots a, b, c are found out, then will the continual product of $x - a, x - b, x - c$, exactly produce the same equation. It is no wonder that an equation has several roots; because in such cases, there are more solutions to a problem than one. So that in one case of it, $x = a$, in another case $x = b$, in a third $x = c$, &c. and they are all comprehended in the general equation. And hence though there be several roots in an equation, yet only one of them will answer one case, or the particular question proposed.

12. That any root substituted for x in the given equation, will make the whole equation to vanish, by destroying all the terms, is proved thus. Let the equation be,

$$\begin{array}{r} x^3 - ax^2 + abx - abc = 0. \\ - b \quad + ac \\ - c \quad + bc \end{array}$$

And let the roots be a, b, c , as before. Then substitute any one, as a , instead of x , and the equation will become

$$\begin{array}{r} a^3 - a^3 + baa - abc = 0, \\ -baa + caa \\ -caa + abc \end{array}$$

Where the terms manifestly destroy one another. And the same will happen, by substituting b or c , for x .

13. If the last term of an equation vanishes (as $a b c$, Art. 12), then one root will be 0; for then the whole equation may be divided by the unknown quantity x or $x - 0$. If the two last terms vanish ($abx + acx + bcx$, and $-abc$), then two roots are $= 0$; if the three last terms vanish, then three roots will be 0; &c.

And on the contrary, if one, two, or three roots, &c. be $= 0$, the last term, the two last, or the three last terms, &c. will vanish out of the equation, and the remaining part of the equation will contain the rest of the roots. Thus in the equation, Art. 12. if the roots b, c be $= 0$; there remains only $x^3 - a + b + c \times x^2 = 0$, or $x - a = 0$, an equation containing the remaining root a .

14. And in any power of a binomial, if each term be multiplied by the index of the unknown quantity therein; it will thereby be reduced to the next inferior power. To prove this, we must observe, that the coefficients of a binomial, are the very same, whether you reckon forward from the beginning, or backward from the end; that is, the first and last are the same; the second and last but one; the third and last but two, &c. For the coefficients of any power of $x + b$, are the same as of $b + x$. In the quadratic $xx + 2bx + bb$,
the

Sect. IV. *Nature of EQUATIONS.* 101

the coefficients are 1, 2, 1. In the cubic $x^3 + 3x^2b + 3xb^2 + b^3$, they are 1, 3, 3, 1. In the fourth power they are 1, 4, 6, 4, 1. In the fifth power, 1, 5, 10, 10, 5, 1; and so on.

Therefore, let any power of $x+b$ be denoted

$$\text{thus, } x^n + nx^{n-1}b + \frac{n \cdot n-1}{2} x^{n-2}bb + \frac{n \cdot n-1 \cdot n-2}{2 \cdot 3} x^{n-3}b^3, \text{ \&c. } + \frac{n \cdot n-1 \cdot n-2}{2 \cdot 3} x^1 b^{n-3} + \frac{n \cdot n-1}{2} x^2 b^{n-2} + nx b^{n-1} + b^n;$$

n being the index of the power, and let m be that of the next inferior power, or $m=n-1$. Now let each term be multiplied by the index of x in each term; that is, by $n, n-1, n-2, \text{ \&c.}$ and we shall have

$$nx^n + n \cdot n-1 \cdot x^{n-1}b + \frac{n \cdot n-1 \cdot n-2}{2} x^{n-2}bb + \frac{n \cdot n-1 \cdot n-2 \cdot n-3}{2 \cdot 3} x^{n-3}b^3, \text{ \&c. } + \frac{n \cdot n-1 \cdot n-2}{2} x^1 b^{n-3} + n \cdot n-1 \cdot x^2 b^{n-2} + nx b^{n-1} + 0.$$

And dividing all by nx , it becomes

$$x^{n-1} + n-1 \cdot x^{n-2}b + \frac{n-1 \cdot n-2}{2} x^{n-3}bb + \frac{n-1 \cdot n-2 \cdot n-3}{2 \cdot 3} x^{n-4}b^3, \text{ \&c. } + \frac{n-1 \cdot n-2}{2} x^1 b^{n-3} + n-1 \cdot x b^{n-2} + b^{n-1};$$

that

$$\text{is, restoring } m, x^m + m \cdot m-1 b + \frac{m \cdot m-1}{2} x^{m-2}bb + \frac{m \cdot m-1 \cdot m-2}{2 \cdot 3} x^{m-3}b^3, \text{ \&c. } + \frac{m \cdot m-1}{2} x^2 b^{m-2} + m x b^{m-1} + b^m,$$

which is manifestly the m^{th} power of $x+b$.

15. Also if the equation resulting from the last operation be taken, and its several terms again multiplied by the index of x in each term; it will be reduced to the next power below that, and so on for more operations. And therefore after each operation one root will be destroyed; or so many roots will be destroyed as there are operations, and the rest will remain.

16. And further: If there be several equal roots of one sort, and also several equal ones of another sort, in any equation. And if the terms of that equation be multiplied by the several indexes of the unknown quantity in each term; an equation will arise wherein one of the equal roots of each sort will be destroyed. And in general, whatever roots there be in any equation, if the terms be respectively multiplied by the indexes of the unknown quantity therein, an equation will come out wherein one root of every sort will be destroyed, whether there be equal roots, or all different. But these things being of little consequence, I shall not detain the reader any longer about them.

17. As impossible roots are such as are produced from the square roots of negative quantities: so impossible equations are those produced from impossible roots; as this equation $a^4 - 4a^3 + 6a^2 + 10a + 22 = 0$, which is produced from these two, $aa + 2a + 2 = 0$, and $aa - 6a + 11 = 0$; the former produced from $a + 1 + \sqrt{-1}$, and $a + 1 - \sqrt{-1}$; and the latter from $a - 3 + \sqrt{-2}$, and $a - 3 - \sqrt{-2}$. These sort of equations have roots that are barely impossible.

Likewise, there are equations that are doubly impossible, or impossible equations of the second degree. And these are produced from equations involving two degrees of impossibility, as this $a^4 + 4a^3 + 8aa + 8a + 5 = 0$, which is produced from the

Sect. IV. *Nature of EQUATIONS.* 103

the equations, $aa + 2a + 2 + \sqrt{-1} = 0$, and $aa + 2a + 2 - \sqrt{-1} = 0$. Such as these cannot be reduced into rational quadratics, as the other may.

P R O B L E M X L I.

To increase or diminish the roots of an equation, by any given quantity.

R U L E.

For the unknown letter substitute a new letter, — the given increment, or + the given decrement. And substitute the powers thereof, in the equation, instead of the powers of the unknown letter.

Ex. 1.

Let $x^3 - px^2 + qx - r = 0$, be given; and let the roots be lessened by the quantity e .

Suppose $y = x - e$, or $x = y + e$. Then

$$\left. \begin{array}{r} x^3 = y^3 + 3ey^2 + 3e^2y + e^3 \\ -px^2 = -py^2 - 2pey - pe^2 \\ +qx = +qy + qe \\ -r = -r \end{array} \right\} = 0, \text{ which}$$

is the equation required.

Ex. 2.

Increase the roots by 4, of this equation

$$a^3 + a^2 - 10a + 8 = 0.$$

Suppose $a + 4 = e$, or $a = e - 4$.

$$\begin{array}{r} \text{Then } a^3 = e^3 - 12e^2 + 48e - 64 \\ + a^2 = \quad ce - 8e + 16 \\ - 10a = \quad -10e + 40 \\ + 8 = \quad \quad \quad + 8 \end{array}$$

$e^3 - 11ee + 30e = 0$, the equation required; reduced, $e^2 - 11e + 30 = 0$, a quadratic.

Cor. 1. *The last term of the transformed equation, is the very same as the equation given, having e in the place of r (in Ex. 1.)*

Cor. 2. *When the last term vanishes, the number assumed (-4 , Ex. 2.) is one of the roots in the equation proposed.*

Schol. *By this rule, all the roots of an equation may be made affirmative; by increasing them by a proper quantity.*

P R O B L E M XLII.

To multiply or divide the roots of an equation, by a given number or quantity.

R U L E.

Assume a new letter; and divide or multiply it by the given number; and substitute its powers in the equation, instead of the unknown quantity.

Ex. 1.

Multiply by 3, this equation $y^3 - \frac{4}{3}y - \frac{146}{27} = 0$.

Suppose $y = \frac{1}{3}z$, then substituting $\frac{1}{3}z$ for y ,

we have $\frac{z^3}{27} - \frac{4}{9}y - \frac{146}{27} = 0$, or reduced
 $z^3 - 12y - 146 = 0$.

Ex. 2.

Divide by $\sqrt{3}$, the equation $x^3 - 2x + \sqrt{3} = 0$.

Let $x = y\sqrt{3}$, which put for x , we have

$3y^3\sqrt{3} - 2y\sqrt{3} + \sqrt{3} = 0$, or $3y^3 - 2y + 1 = 0$.

Cor. *By this rule, fractions or surds may be taken out of an equation; by dividing the new letter by the common denominator; or by multiplying the new letter by the surd quantity.*

P R O -

PROBLEM XLIII.

To change the roots of an equation into their reciprocals.

R U L E.

In the given equation, instead of the root, substitute a unit divided by some other letter.

Example.

Let $3y^3 - 2y + 1 = 0$, be given.

Put $y = \frac{1}{z}$, then $\frac{3}{z^3} - \frac{2}{z} + 1 = 0$.

reduced $3 - 2z^2 + z^3 = 0$.

or $z^3 - 2z^2 + 3 = 0$.

Schol. By this rule the greatest root is changed into the least, and the least into the greatest, &c.

PROBLEM XLIV.

To compleat a deficient equation.

An equation is *compleat*, when it has all its terms, or those containing all the powers of the unknown quantity; and *deficient*, when any power is wanting.

R U L E.

Increase or diminish the roots of the equation, by some given quantity (by Prob. 41).

Example.

Suppose $a^3 + 2a - 5 = 0$, deficient.

Let $e + 1 = a$, then

$$a^3 = e^3 + 3ee + 3e + 1$$

$$+ 2a = \quad \quad + 2e + 2$$

$$\underline{-5 = \quad \quad \quad -5}$$

$$e^3 + 3ee + 5e - 2 = 0, \text{ compleat.}$$

Schol.

Schol. An equation may be rendered compleat, by multiplying by the same letter with some quantity added, as $a+1$; but then it raises the equation a degree higher.

P R O B L E M XLV.

To depress an equation to a lower dimension; one of its roots being given.

R U L E.

Put the equation $=0$, and divide it by the unknown quantity — the root given.

Example.

Given $a^3+a^2-10a+8=0$, one root $a=-4$.

$a+4=0$) $a^3+a^2-10a+8=0$ ($aa-3a+2=0$ the equation req.

$$a^3+4a^2$$

$$\underline{-3a^2-10a}$$

$$\underline{-3a^2-12a}$$

$$+2a+8$$

$$\underline{+2a+8}$$

$$0$$

2 R U L E.

Put a new letter added to that root, equal to the unknown quantity; and substitute that and its powers in the equation.

Example.

Let $a^3+a^2-10a+8=0$, be given, and $a=-4$.

Put $a=e-4$. Then

$$+ a^3 = e^3 - 12e^2 + 48e - 64$$

$$+ a^2 = + ee - 8e + 16$$

$$- 10a = - 10e + 40$$

$$+ 8 = 8$$

$$\underline{0 = e^3 - 11e^2 + 30e + 0}$$

reduced

$$e^3 - 11e^2 + 30e = 0.$$

PRO-

PROBLEM XLVI.

To find how many roots are affirmative, and how many negative, in a given equation.

R U L E.

Range the terms of the equation according to the dimensions of the unknown quantity. And if the equation is not compleat, make it so by Prob. 44.

Then observe how often + follows —, or — follows +, that is, how many changes of the signs there are; and there are so many affirmative roots in the equation.

Also, as often as two like signs stand together, so often there is a negative root.

Ex. 1.

$$\text{Given } x^4 - x^3 - 19xx + 49x - 30 = 0:$$

Here the signs are + — — + —, and there are three changes; from the first to the second, from the third to the fourth, and from the fourth to the fifth term: therefore there are three affirmative roots. Also, in the second and third terms, two negatives stand together, and in none else, consequently there is one negative root.

Ex. 2.

$$\text{Suppose } x^4 + 5x^3 - 7x^2 - 29x + 30 = 0.$$

The signs are + + — — +
 roots neg. af. neg. af.

So there are two affirmative, and two negative roots.

Ex.

Ex. 3.

Let the equation be $a^3 - 7a + 6 = 0$.

This equation being defective is to be completed.

$$\begin{array}{r}
 a^3 * - 7a + 6 = 0. \\
 \text{mult. by } a + 1 = 0. \\
 \hline
 a^4 * - 7a^2 + 6a \\
 + a^3 * - 7a + 6. \\
 \hline
 a^4 + a^3 - 7a^2 - a + 6 = 0.
 \end{array}$$

So there are two affirmative, and two negative roots in this last equation, and one of the negative roots being -1 , (by the multiplication of $a + 1 = 0$,) therefore, the given equation contains two affirmative roots, and one negative.

The reason of this rule appears from Art. 10. Prob. 40.

S C H O L I U M.

This rule does not hold good, if there be impossible roots in the equation; except so far as these impossible roots may be taken for ambiguous ones, that is, for either affirmative or negative roots. As in the equation $x^3 - 6x^2 + 13x - 10 = 0$, which by this rule gives three affirmative roots, but in reality it has but one root, which is 2, the rest are imaginary.

There are also some rules whereby to judge how many impossible roots are in an equation, but they are so very tedious, and of so little use, that I shall not trouble the reader with them. See *Newton's Universal Arithmetic*, p. 197.

PROBLEM XLVII.

To change the affirmative roots into negatives, and the negatives into affirmatives.

R U L E.

Place cyphers for the deficient terms, if there be any; then change the signs of all the even terms, that is, of the second, fourth, sixth, &c. terms of the equation.

Ex. 1.

Given $x^3 + 8x + 24 = 0$.

That is, $x^3 + 0 + 8x + 24 = 0$.

transformed $x^3 - 0 + 8x - 24 = 0$.

In the given equation $x = -2$, in the transformed equation $x = +2$.

Ex. 2.

Suppose $x^4 - 4x^3 - 19x^2 + 106x - 120 = 0$.

transformed $x^4 + 4x^3 - 19x^2 - 106x - 120 = 0$.

In the former equation the roots are 2, 3, 4 and -5; and in the latter 5, -2, -3, and -4.

The reason of this process is plain from Art. 11. Prob. 40. and may be demonstrated thus. In the given equation, we have $+x$ for the root. Now suppose $-x$ to be a root. Let this be substituted in the given equation, and it produces $-x^3 - 8x + 24 = 0$, that is, $x^3 + 8x - 24 = 0$, as in Exam. 1. And $x^4 + 4x^3 - 19x^2 - 106x - 120 = 0$, as in Exam. 2. For it is plain, all the odd powers of x will now be negative, which before were affirmative, the rest remaining as before. Whence the signs of all the odd powers will be changed, according to the rule.

S E C T. V.

Ranging the terms; working by general forms; substitution and restitution; taking away any term of an equation; extermination of unknown quantities; the designation of quantities by letters; registering the steps.

P R O B L E M XLVIII.

To range the terms of an equation, or dispose of them in the best manner for any operation.

R U L E.

THIS is done by placing these terms foremost that contain the highest power of the unknown quantity; and in the following places, those of less dimensions; so that the powers in the several terms may continually decrease from the highest, according to the series of the natural numbers. But in many cases, the contrary method is to be followed, and the lowest power taken first.

Ex. 1:

$$\text{Let } az^3 + z^4 - bz^3 - b^4 + ab^3 = 0.$$

$$\text{Place it thus, } z^4 + az^3 \quad * \quad * \quad + ab^3 = 0.$$

$$\quad \quad \quad -b \quad \quad \quad -b^4$$

Ex. 2:

$$\text{Suppose } x^4 + ax^3 + bx^2 - bx^3 + cx = dx - ab^3 + b^4.$$

$$\text{ranged } x^4 + ax^3 + bx^2 + cx + ab^3 = 0.$$

$$\quad \quad \quad -b \quad \quad \quad -d + b^4$$

PROBLEM XLIX.

To work by a general form.

R U L E.

Write down each letter or quantity in the general form, and after it (with the sign =), each letter it represents in that particular case; which will give several equations.

Then cast your eye over the general form, and observe the general quantities therein, and look for them on the first side of the equations; and what you find them equal to, on the right hand, write down, instead of them, each one by one, till you have gone through the general form; and you will have the solution.

When the quantities are many, it will be the best way to write down the general form first, and the particular one under it, each quantity under its correspondent; then it will appear by inspection what letters to substitute.

*Ex. 1.**To involve $aa-xx$ to the 5th power.*

This is to be done by the general form in Cor. 1. Prob. 5. therefore we have

$$a = aa$$

$$e = -xx$$

$$n = 5,$$

Whence $a+e = aa-xx = a^5 + 5 \times aa^4 \times -xx + 5 \times \frac{5-1}{2} \times aa^3 \times x^4 + 5 \times \frac{5-1}{2} \times \frac{5-2}{3} aa^2 \times -x^6 + 5 \times \frac{5-1}{2} \times \frac{5-2}{3} \times \frac{5-3}{4} aa \times x^8 + 5 \times \frac{5-1}{2} \times \frac{5-2}{3} \times \frac{5-3}{4} \times \frac{5-4}{5} \times -x^{10} = a^{10} - 5a^8x^2 + 10a^6x^4 - 10a^4x^6 + 5a^2x^8 - x^{10}$, the power required. *Ex.*

Ex. 2.

Extract the square root of $28 - \sqrt{300}$.

This is to be done by the form in 1 Rule, Prob. 32.

Here $A=28$, $B=\sqrt{300}$, $D=\sqrt{784-300}=22$,
 $\sqrt{\frac{A+D}{2}} = \sqrt{\frac{28+22}{2}} = 5$, $\sqrt{\frac{A-D}{2}} =$
 $\sqrt{\frac{28-22}{2}} = \sqrt{3}$. Therefore $\sqrt{A-B} = 5 - \sqrt{3}$,
 the root required.

Ex. 3.

To find a quantity, by which if $\sqrt[5]{2} - \sqrt[5]{6}$ be multiplied, the product will be rational.

This is to be done by Prob. 36.

Here $n=5$, $A=2$, $B=6$.

And the multiplier $\sqrt[5]{16} + \sqrt[5]{8 \times 6} + \sqrt[5]{4 \times 36} +$
 $\sqrt[5]{2 \times 216} + \sqrt[5]{1296}$.
 mult. $\sqrt[5]{16} + \sqrt[5]{8 \times 6} + \sqrt[5]{4 \times 36} + \sqrt[5]{2 \times 216} + \sqrt[5]{1296}$
 by $\sqrt[5]{2} - \sqrt[5]{6}$

 $\sqrt[5]{32} + \sqrt[5]{96} + \sqrt[5]{8 \times 36} + \sqrt[5]{4 \times 216} + \sqrt[5]{2592}$
 $-\sqrt[5]{96} - \sqrt[5]{8 \times 36} - \sqrt[5]{4 \times 216} - \sqrt[5]{2592} - \sqrt[5]{7776}$

 $2 - 6 = -4$. product.

PROBLEM L.

To shorten the work by substitution and restitution.

In any operation, when the quantities grow very numerous, or very much compounded, it will make the work very tedious; and therefore it ought to be made shorter as follows,

RULE.

R U L E.

Assume a new letter to represent or stand for any number of given quantities; and likewise some different letter to stand for the coefficient of any power of the unknown quantity; do so for as many of the coefficients as are compounded. Likewise, put letters for the numbers concerned; then work with these instead of the original quantities, which will make the work easier. And this is called *Substitution*.

When the operation is over, each number or compound quantity must be restored again instead of its letter; and this is called *Restitution*.

Ex. 1:

$$\text{Let } aa + ba - ca + da = dc.$$

Put $s = b - c + d$. Then the equation becomes

$$aa + sa = dc.$$

Ex. 2:

$$\text{Let } \frac{bx}{a - 2x} \times \sqrt{aa - xx} = \frac{bx}{cx^2 - dx^2 + cx}$$

Put $c - d = p$. Then

$$\frac{bx}{a - 2x} \times \sqrt{aa - xx} = \frac{bx}{pxx + cx}$$

multiply by $pxx + cx$. Then

$$apxx - 2px^3 + acx - 2cxxx \times \sqrt{aa - xx} = bxx.$$

Put $ap - 2c = q$. Then

$$\frac{qxx - 2px^3 + acx}{acx + qxx - 2px^3} \times \sqrt{aa - xx} = bx.$$

or
$$\frac{qxx - 2px^3 + acx}{acx + qxx - 2px^3} \times \sqrt{aa - xx} = bx.$$

Squared $\frac{qxx - 2px^3 + acx}{acx + qxx - 2px^3} \times \sqrt{aa - xx} = bxx$, &c.
where the values of p, q , may be restored.

PROBLEM LI.

To take away the second term of an equation.

R U L E.

Divide the coefficient of the second term by the index of the highest power; annex the quotient, with its sign changed, to some new letter, which substitute for the root, in the given equation.

Ex. 1.

Suppose $a^3 + aa - 10a + 8 = 0$.

Put $e - \frac{1}{3} = a$. Then

$$\begin{array}{r} a^3 = e^3 - e^2 + \frac{1}{3} + \frac{1}{27} \\ + aa = + ee - \frac{2}{3} - \frac{1}{9} \\ - 10a = - 10e + \frac{10}{3} \\ + 8 = + 8 \\ \hline 0 = e^3 * - 10\frac{1}{3}e + 11\frac{11}{27}, \text{ the equation required.} \end{array}$$

Ex. 2.

Let $y^4 - 8ay^3 + a^4 = 0$, be given.

Let $y = x + \frac{8a}{4} = x + 2a$.

$$\begin{array}{r} \text{then } y^4 = x^4 + 8ax^3 + 24a^2x^2 + 32a^3x + 16a^4 \\ - 8ay^3 = - 8ax^3 - 48a^2x^2 - 96a^3x - 64a^4 \\ + a^4 = + a^4 \\ \hline 0 = x^4 * - 24a^2x^2 - 64a^3x - 47a^4 = 0. \end{array}$$

Schol. Hence by this and the 43d problem, an equation may be found, which wants the last term but

but one. For if the second term be taken away by this problem, and the equation transformed by Prob. 43, you will have the equation required.

PROBLEM LII.

To take away any term out of an equation.

R U L E.

Take a new letter for the root, to which add an unknown quantity; and substitute this sum and the powers thereof, into the given equation. Then any term put equal to nothing, will determine the value of that assumed unknown quantity.

Ex. 1.

Suppose $x^4 - 3x^3 + 3x^2 - 5x - 2 = 0$.

Put $y + e = x$.

$$\begin{array}{r} \text{Then } x^4 = y^4 + 4y^3e + 6y^2e^2 + 4ye^3 + e^4 \\ -3x^3 = -3y^3 - 9y^2e - 9ye^2 - 3e^3 \\ +3x^2 = +3y^2 + 6ye + 3e^2 \\ -5x = -5y - 5e \\ +2 = +2 \end{array} = 0$$

Then, if the second term is to be taken away, make $4y^3e - 3y^3 = 0$, or $4e = 3$; therefore $e = \frac{3}{4}$.

Ex. 2.

The same supposed; to take away the third term.

Here we shall have $6y^2e - 9y^2e + 3yy = 0$; reduced, $2ee - 3e + 1 = 0$, the resolving of which quadratic equation gives the value of e . Then $y + e$ gives the value of x , so that the third term may vanish.

Ex. 3.

The same thing still supposed; to take away the fourth or fifth term.

For the fourth term, $4e^3 - 9e^2 + 6e - 5 = 0$, a cubic equation whose root is e ; and $y + e = x$, makes the fourth term vanish.

For the fifth term, $e^4 - 3e^3 + 3e^2 - 5e + 2 = 0$, a fourth power whose root is e . Then $y + e = x$, which substituted in the equation, makes the last term vanish.

Cor. 1. Hence the third, fourth, fifth, &c. term, may be taken out of the equation; by resolving a quadratic, cubic, fourth power, &c. equation.

Cor. 2. Hence if the last term of an equation (as $e^4 - 3e^3 + 3e^2 - 5e + 2$) be $= 0$, then one root (x) is $= 0$; for then $x = 0$, or x will divide the equation. If two of the last terms be $= 0$, two values of the root will be $= 0$, and so on. But if the last term does not vanish, there is no root $= 0$.

Schol. After the same rule any term may be made equal to any given quantity; by putting the said term equal to that quantity.

P R O B L E M LIII.

To exterminate a single letter, or a quantity of one dimension, out of several equations.

I R U L E.

Seek the value of the quantity to be expelled, in two equations; and put these values equal to one another.

Ex. 1.

Let $a+x=b+y$
and $2x+y=3b$ } to exterminate y .

By transposing b , $a+x-b=y$, and by transposing $2x$, $y=3b-2x$. Therefore $a+x-b=3b-2x$.

And by reduction $3x=4b-a$, and $x=\frac{4b-a}{3}$.

Ex. 2.

Let $ax-2by=ab$
and $xy=bb$, } to exterminate y .

Here $ax-ab=2by$, and $y=\frac{ax-ab}{2b}$.

Also $y=\frac{bb}{x}$; therefore $\frac{ax-ab}{2b}=\frac{bb}{x}$, and re-

ducing $xx-bx=\frac{2b^2}{a}$.

2 R U L E.

Find, by reduction, the value of one unknown quantity, in one equation; and substitute that value for it, in all the other equations. Proceed thus, with another unknown quantity, &c.

Ex. 3.

Let $a+x=2b-y$
and $3ax-yx=d$ } to expell y .

By the first equation $y=2b-a-x$, put this value in the second equation; then

$3ax - x \times 2b - a - x = d$, that is, $3ax - 2bx + ax + xx = d$, or $4ax - 2bx + xx = d$.

Ex. 4.

Suppose $x+y+z=a$
 $3y=x+2z$
 $ax=xy$ } to expunge z and y .

By the first equation, $z=a-x-y$,

By the second equation, $3y=x+2a-2x-2y$,

By the third, $a \times a - x - y = xy$, or

$$aa - ax - ay = xy.$$

The former reduced $5y = 2a - x$.

and since $aa - ax - ay = xy$.

From these to expunge y .

By the former $y = \frac{2a-x}{5}$, and by the latter

$aa - ax = ay + xy$, and $y = \frac{aa-ax}{a+x}$. Therefore

$\frac{2a-x}{5} = \frac{aa-ax}{a+x}$, in which equation there is only one unknown quantity x .

Cor. 1. By each given equation, one unknown quantity may be taken away. And consequently when there are as many equations as unknown quantities, they may be all taken away but one.

Cor. 2. If there be more unknown quantities than equations, there will remain in the last equation more unknown quantities by 1, than that excess amounts to.

PROBLEM LIV.

To exterminate an unknown quantity of several dimensions.

R U L E.

Find the value of its greatest power in two equations; then if they are not the same, multiply the lesser

lesser power, so that it may become equal to the greater. Then put these values equal to each other, and there will come out a new equation, with a less power of the unknown quantity. And by repeating this operation, the quantity will at last be taken away.

Ex. I:

Let $ace + be + c = 0$
and $fee + ge + b = 0$ } to expunge e .

By transposing and dividing $-ee = \frac{be+c}{a}$, and

$$-ee = \frac{ge+b}{f}. \text{ Therefore } \frac{be+c}{a} = \frac{ge+b}{f}.$$

And multiplying, $bef + cf = age + ab$, and by transposing $bfe - age = ab - cf$, and dividing,

$$e = \frac{ab - cf}{bf - ag}. \text{ And multiplying by } -e,$$

$$-ee = \frac{-abe + cfe}{bf - ag}. \text{ Whence } \frac{be+c}{a} = \frac{-abe + cfe}{bf - ag}.$$

And multiplying alternately $bbfe + bcf - abge - age = acfe - aabe$. And transposing and dividing,

$$e = \frac{cgc - bcf}{bbf - abg - acf + aab}. \text{ Therefore}$$

$$\frac{ab - cf}{bf - ag} = \frac{ag - bcf}{bbf - abg - acf + aab}. \text{ Then multiply-}$$

ing and reducing, $hbba + cga + bbf b = 0$.

$$\begin{aligned} & - 2cfb - bgfc \\ & - bgb + cff \end{aligned}$$

2 R U L E.

For two quadratic equations.

$$ax^2 + bx + c = 0.$$

$$\text{and } fx^2 + gx + b = 0.$$

to exterminate x . Here a, b, c, f, g, b , are either given

given quantities, or composed of given quantities, and some other unknown quantity y . Thus

make $bf-ag=A$, $bb-cg=B$, and $cf-ab=D$.
then $AB+DD=0$.

To prove this rule, we have $-x^2 = \frac{bx+c}{a} = \frac{gx+b}{f}$, which reduced is $\overline{bf-ag} \times x + \overline{cf-ab} = 0$; that is, $Ax+D=0$. Whence $Ax^2+Dx=0$; therefore $-xx = \frac{Dx}{A} = \frac{bx+c}{a}$, which reduced is $x = \frac{cA}{aD-bA}$. In like manner $-xx = \frac{Dx}{A} = \frac{gx+b}{f}$, which reduced is $x = \frac{bA}{fD-gA}$. Whence $\frac{cA}{aD-bA} = \frac{bA}{fD-gA}$. And this reduced is $\overline{cf-ab} \times D + \overline{bb-cg} \times A = 0$, that is $AB+DD=0$.

The Newtonian Rule is,

$$ab \times \overline{ab-bg-2cf} + bf \times \overline{bb-cg} + c \times \overline{agg+cff} = 0.$$

3 R U L E.

For a cubic and a quadratic equation,

$$ax^3+bx^2+cx+d=0,$$

$$\text{and } fx^2+gx+b=0.$$

Make $\overline{fc-ab} = D$, $\overline{fb-ag} = A$. Then

$$\overline{fD-gA} \times \overline{bD-fAg} + \overline{dff-bA}^2 = 0.$$

For multiplying the first equation by f , and the second by ax , and subtracting one from the other, we have

$$\overline{bf-ag} \times x^2 + \overline{fc-ab} \times x + \overline{fd} = 0; \text{ and since } \overline{fx^2+gx+b} = 0, \text{ these two equations come under the}$$

Sect. V. EXTERMINATION. 121

the last rule, making $a = bf - ag$, $b = fc - ab$, $c = fd$.

And $A = f \times fc - ab - g \times bf - ag$. $B = b \times fc - ab - fdg$.

$D = ffd - b \times bf - ag$. Whence by that rule,

$$\frac{f \times fc - ab - g \times bf - ag}{ffd - b \times bf - ag} \times \frac{b \times fc - ab - fdg}{bD - fdg + ffd - bA} + \frac{ffd - b \times bf - ag}{bD - fdg + ffd - bA} = 0, \text{ that is, according to the present designation of the letters A, B, C; } fD - gA \times bD - fdg + ffd - bA^2 = 0.$$

The Newtonian Rule is,

$$\left. \begin{aligned} &abb \times ab - bg - 2cf + bfb \times bb - cg - 2df \\ &+ cb - dg \times agg + cff + df \times 3agb + bgg + dff \end{aligned} \right\} = 0.$$

4 RULE.

For a quadratic and a fourth power.

$$ax^4 + bx^3 + cx^2 + dx + e = 0.$$

$$\text{and } fx^2 + gx + b = 0.$$

Make $A = bf - ag$. $D = cf - ab$. Then

$$\frac{df^3 - gfd + gg - fbx \times A \times dbff - cgff - bbA}{+ef^3 + gbA - f bD^2} = 0.$$

For multiply the first equation by f , and the latter by axx ; their difference will be $bf - ag \times x^3 + cf - ab \times x^2 + dfx + ef = 0$. Or $Ax^3 + Dx^2 + dfx + ef = 0$. And since $fx^2 + gx + b = 0$. Therefore these two equations come under the last rule; in which writing A for a , D for b , df for c , ef for d ; and lastly $fD - gA$ instead of A , and $ffd - bA$ for D , you will get the rule, as above.

The Newtonian Rule is,

$$\left. \begin{aligned} &ab^3 \times ab - bg - 2cf + bfb \times bb - cg - 2df \\ &+ agg + cff \times cbh - dgk + egg - 2efb \\ &+ dfb \times 3agb + bgg + dff \\ &+ eff \times 2abb + 3bgb - dfg + cff \\ &- efgb \times bg + 2ab \end{aligned} \right\} = 0.$$

5 RULE.

5 R U L E.

For two cubic equations.

$$ax^3 + bx^2 + cx + d = 0.$$

and $fx^3 + gx^2 + bx + k = 0.$

Make $A = bf - ag$, $C = df - ak$, $D = cf - ab$.

and $P = cA^2 - aAC - bAD + aDD$.

$$Q = cAC - aCC - dAD.$$

$$R = dAA - bAC + aCD.$$

Then $PQ + RR = 0.$

For multiply the first equation by f , and the latter by a , and their difference will be found $\overline{bf - ag} \times x^2 + \overline{fc - ab} \times x + \overline{fd - ak} = 0$; that is, $Ax^2 + Dx + C = 0$. And since $ax^3 + bx^2 + cx + d = 0$; these two equations come under the third rule; in which writing A , D , C for f , g , b , respectively; and likewise $cA - aC$ for A , and $bA - aD$ for D ; the rule will be evident.

The Newtonian Rule is,

$$\begin{aligned} &+ \overline{ab - bg - 2cf} \times \overline{aabb - acbb} \\ &+ \overline{bdfb} \times \overline{ak + bk - cg - 2df} \\ &+ \overline{aakk} \times \overline{bk - ak + 2gc + 3df} \\ &+ \overline{bbfk} \times \overline{bk - 2dg} \\ &+ \overline{cdb - dag - cck + 2bdk} \times \overline{agg + cff} \\ &+ \overline{3agb + bgg + dff - 3afk} \times \overline{ddf} \\ &+ \overline{bcfk} \times \overline{cg + df - 3ak - bb} \\ &- \overline{agk} \times \overline{bbk + 3adb + cdf} = 0. \end{aligned}$$

6 R U L E.

For a cubic and a fourth power.

$$ax^4 + bx^3 + cx^2 + dx + e = 0.$$

and $fx^3 + gx^2 + bx + k = 0.$

Make $A = fb - ag$, $C = fd - ak$, $D = cf - ab$.

Then

Then put

$$P = C \times \overline{fD-gA}^2 - A \times \overline{fD-gA} \times \overline{ffe-kA} - D \times \overline{ffe-kA} + A \times \overline{fC-bA}^2$$

$$Q = C \times \overline{fD-gA} \times \overline{ffe-kA} - A \times \overline{ffe-kA}^2 - ef \times \overline{fD-gA} \times \overline{fC-bA}$$

$$R = ef \times \overline{fD-gA}^2 - D \times \overline{fD-gA} \times \overline{ffe-kA} + A \times \overline{ffe-kA} \times \overline{fC-bA}$$

Then $PQ + RR = 0$.

Or thus,

Put $E = \overline{fD-gA}$, $F = \overline{ffe-kA}$, $G = \overline{fC-bA}$.

$$P = \overline{CE-AG} \times E + \overline{AF-DE} \times G.$$

$$Q = \overline{CE-AG} \times F - fe \times EG.$$

$$R = \overline{feE-DF} \times E + AFG.$$

Then $PQ + RR = 0$, as before.

For multiplying the first equation by f , and the last by a , the difference is $Ax^3 + Dx^2 + Cx + ef = 0$. And since $fx^3 + gx^2 + bx + k = 0$; it will come under Rule 5, in which write A, D, C, ef , for a, b, c, d respectively; and likewise $fD-gA, ffe-kA$, and $fC-bA$, for A, C, D , respectively; and the rule will appear.

Ex. 2.

Let $xx + 5x - 3yy = 0$,

and $3xx - 2yx + 4 = 0$,

to exterminate x .

By Rule 2, $a=1, b=5, c=-3yy, f=3, g=-2y, b=4$, and $A=15+2y, B=20-6y^2, D=-9y^2-4$.

$$\text{Then } AB + DD = \overline{15+2y} \times \overline{20+6y^2} + \overline{-9y^2-4}^2 = 300 + 40y - 90y^2 - 12y^4 + 81y^4 + 72y^2 + 16 = 0.$$

Ex.

Ex. 3.

Suppose $y^3 - xy - 3x = 0$.and $y^2 + xy - xx + 3 = 0$.to expunge y .Here by Rule 3, $a=1$, $b=-x$, $c=0$, $d=-3x$.and $f=1$, $g=x$, $h=-xx+3$. $A=-x-x=-2x$, $D=xx-3$.

$$fD - gA = xx - 3 + 2xx = 3xx - 3$$

$$bD - fdg = -x^4 + 6x^2 - 9 + 3x^2 = -x^4 + 9x^2 - 9.$$

$$dff - bA = -3x - 2x^3 + 6x = 3x - 2x^3.$$

$$\text{Then } 3xx - 3 \times -x^4 + 9x^2 - 9 + \frac{3x - 2x^3}{12x^4 + 4x^6} = 0.$$

$$\text{Or, } -3x^6 + 27x^4 - 27x^2 + 3x^4 - 27x^2 + 27 + 9x^2 - 12x^4 + 4x^6 = 0.$$

$$\text{And reduced } x^6 + 18x^4 - 45x^2 + 27 = 0.$$

Ex. 4.

Let $y^4 - 3x^2y + 3 = 0$,
 and $2y^3 + xy^2 - 4x^3 = 0$, } to expunge y .

By Rule 6, $a=1$, $b=0$, $c=0$, $d=-3x^2$, $e=3$,
 $f=2$, $g=x$, $h=0$, $k=-4x^3$.

Then $A=-x$, $C=-2x^3$, $D=0$. Whence

$$E = xx, F = 12 - 4x^4, G = -4x^3. \text{ And}$$

$$P = \frac{-2x^5 + 12x - 4x^5 \times xx + 4x^4 \times -4x^3}{12x^3 - 6x^7 - 16x^7} = \frac{12x^3 - 22x^7}{12x^3 - 22x^7}.$$

$$Q = \frac{-2x^5 + 12x - 4x^5 \times 12 - 4x^4 \times 24x^5}{12x - 6x^5 \times 12 - 4x^4 \times 24x^5} \\ = \frac{144x - 96x^5 + 24x^9}{144x - 96x^5 + 24x^9}.$$

$$R = 6x^4 + 4x^4 \times 12 - 4x^4 = 54x^4 - 16x^8.$$

Whence

Whence

$$\begin{aligned}
 PQ + RR &= \frac{12x^3 - 22x^7 \times 144x - 96x^5 + 24x^9}{+ 54x^4 - 16x^8} = 1728x^4 - 4320x^8 + \\
 &2400x^{12} - 528x^{16} + 2916x^8 - 1728x^{12} + 256x^{16} = 0, \\
 \text{reduced, } &68x^{12} - 168x^8 + 351x^4 - 432 = 0.
 \end{aligned}$$

S C H O L I U M.

In the solution of determined problems, you will often have three or more equations, involving as many unknown quantities. Then these must be exterminated one after another, by degrees, by repeating the foregoing rules; till at last there remains only one unknown quantity contained in one final equation. But a person used to these sorts of computations, will often find shorter methods than by these particular rules, but the finding those, is only to be attained by constant practice.

P R O B L E M L V.

To designate or denote any affections of literal quantities, as, sums, products, &c.

R U L E.

The original quantities being written down; any affections of them, as sums, differences, products, quotients, &c. are got by the rules of algebraic addition, subtraction, multiplication, division, &c. before laid down.

Ex.

Ex. 1.

There are two quantities, a the greater, and e the lesser, to find the sum, difference, product, &c. as follows.

The sum	— — — — —	$a + e$
difference	— — — — —	$a - e$
product	— — — — —	ae
greater divided by the less		$\frac{a}{e}$
less divided by the greater		$\frac{e}{a}$
sum of their squares		$aa + ee$
difference of their squares		$aa - ee$
sum of their sum and diff.		$2a$
diff. of their sum and diff.		$2e$
prod. of the sum and diff.		$a + e \times a - e$, Or $aa - ee$
square of the sum		$aa + 2ae + ee$
square of the difference		$aa - 2ae + ee$
sum of the squares of the sum and difference	}	$2aa + 2ee$
difference of the squares of the sum and diff.	}	$4ae$
square of the product		$aaee$
cube of the greater		a^3
cube of the lesser		e^3
cube of the sum		$a^3 + 3a^2e + 3ae^2 + e^3$
cube of the difference		$a^3 - 3a^2e + 3ae^2 - e^3$

Ex. 2.

There are two quantities, whose sum is b , and the greater is a ; what is the lesser, the difference, &c.

Lesser	_____	_____	_____	$b - a$
difference	_____	_____	_____	$2a - b$
product	_____	_____	_____	$ab - aa$
greater \div by the lesser	_____	_____	_____	$\frac{a}{b - a}$
sum of their squares	_____	_____	_____	$2aa + bb - 2ba$
difference of their squares	_____	_____	_____	$2ba - bb$
sum of the sum and difference	_____	_____	_____	$2a$
difference of the sum and difference	_____	_____	_____	$2b - 2a$
product of the sum and difference	_____	_____	_____	$2ab - bb$
square of the difference	_____	_____	_____	$4aa - 4ab + bb$
difference of the squares of the sum and difference	_____	_____	_____	$4ab - 4aa$
square of the product	_____	_____	_____	$a^2b^2 - 2ba^3 + a^4$

Ex.

Ex. 3.

There are two quantities, the greater is a , and the greater is to the lesser as r to s , what is the lesser, &c.

$$\text{The lesser } (r : s :: a :) \quad \text{---} \quad \frac{sa}{r}$$

$$\text{the sum} \quad \text{---} \quad \text{---} \quad a + \frac{sa}{r}$$

$$\text{difference} \quad \text{---} \quad \text{---} \quad a - \frac{sa}{r}$$

$$\text{product} \quad \text{---} \quad \text{---} \quad \frac{saa}{r}$$

$$\text{sum of the squares} \quad \text{---} \quad aa + \frac{ssaa}{rr}$$

$$\text{difference of the squares} \quad \text{---} \quad aa - \frac{ssaa}{rr}$$

$$\text{greater divided by the lesser} \quad \frac{r}{s}$$

$$\text{product of the sum and differ.} \quad aa - \frac{ssaa}{rr}$$

$$\text{sum of the squares of the sum} \quad \left. \begin{array}{l} \text{and difference} \end{array} \right\} 2aa + \frac{2ssaa}{rr}$$

$$\text{difference of the squares of the sum} \quad \left. \begin{array}{l} \text{and difference} \end{array} \right\} \frac{4saa}{r}$$

$$\text{the sum divided by the greater} \quad 1 + \frac{s}{r}$$

$$\text{the difference divided by the lesser} \quad \frac{r}{s} - 1$$

Ex. 4.

The product of two quantities is p , and the lesser is e , what is the greater, &c.

Greater	$\frac{p}{e}$
sum	$\frac{p}{e} + e$
difference	$\frac{p}{e} - e$
lesser \div by the greater	$\frac{ee}{p}$
sum of their squares	$\frac{pp}{ee} + ee$
difference of their squares	$\frac{pp}{ee} - ee$
sum of the sum and difference	$\frac{2p}{e}$
diff. of their sum and diff.	$2e$
square of the sum	$\frac{pp}{ee} + 2p + ee$
square of the difference	$\frac{pp}{ee} - 2p + ee$
diff. squares of the sum and diff.	$4p$
the sum \div by the difference	$\frac{p+ee}{p-ee}$

PROBLEM LVI.

To keep a short account of the steps in any operation.

In long and tedious operations, it is necessary to shew, how one step is produced from another, or

one equation derived from other foregoing ones; which to explain in words would take up a great deal of room. Therefore the method of tracing the several steps, will be best done by registering them in the margin.

R U L E.

Against every step write the numbers 1, 2, 3, &c. in order, and set down, in the margin on the left hand, the step or steps in figures, that each step is produced from; with the signs + - ×, &c. according to the several operations, used; by which means one may see at one view how any equation comes, or is produced; and when an absolute number is registered, it must be put in a parenthesis (); and if any quantity is added, subtracted, &c. it must be put down.

Example.

Let	{	1	$a + e = b.$
		2	$a - e = c.$
		<hr/>	
1 + 2		3	$2a = b + c$
1 - 2		4	$2e = b - c$
1 × 2		5	$aa - ee = bc$
1 ÷ 2		6	$\frac{a+e}{a-e} = \frac{b}{c}$
1 lw 2p		7	$\sqrt{a+e} = \sqrt{b}$
4 © 2p		8	$4ee = bb - 2bc + cc$
3 + 7		9	$2a + \sqrt{a+e} = b + c + \sqrt{b}$
4 × 5		10	$2aae - 2e^2 = bbc - bcc$
3 + (4)		11	$2a + 4 = b + c + 4$
4 ÷ (4)		12	$\frac{e}{2} = \frac{b-c}{4}$
9 - $\sqrt{a+e}$		13	$2a = b + c + \sqrt{b} - \sqrt{a+e}$
3 = 13		14	$b + c = b + c + \sqrt{b} - \sqrt{a+e}$
			&c.

Sect. V. TRACING THE STEPS. 131

EXPLANATION.

1+2 signifies that the third step is found by adding the first and second steps together. 1—2 signifies, the fourth step is got by subtracting the second from the first. Likewise, the fifth step (1×2) is had by multiplying the first and second: the sixth step, by dividing the first by the second: the seventh, by extracting the square root of the first: the eighth ($4 \odot 2p$) is had by squaring the fourth: the ninth (3+7), by adding the third and seventh steps: the tenth (4×5), by multiplying the fourth and fifth steps: the eleventh (3+(4)), is had by adding the number 4 to the third step: the twelfth ($4 \div (4)$), shews that it is gained by dividing the fourth step by the number 4: and the thirteenth ($9 - \sqrt{a+e}$), is had by subtracting $\sqrt{a+e}$ from the ninth: the fourteenth ($3 = 13$) is got by making the third and thirteenth equations equal; and so for others.



S E C T. VI.

Infinite Series.

AN infinite series is formed, either by actually dividing any fractional quantity having a compound denominator; or by extracting the root of a surd, and such series being continued will run on *ad infinitum*, in the manner of a decimal fraction. And in many cases the law of the progression of the terms will be evident, by obtaining a few of the foremost; and consequently may be continued without actually performing the whole operation.

P R O B L E M LVII.

To find the value of a fraction or surd, to be designated by an infinite series.

I R U L E.

Proceed in the same manner as is taught in Prob. iv. Rule 2. for division; or in Prob. vi. Rule 2 and 3, continuing on, the operation at pleasure.

Ex. 1.

Let $\frac{ax}{a-x}$ be given.

$$\begin{array}{r}
 a-x)ax \\
 \underline{ax-xx} \\
 +xx \\
 \underline{+xx - \frac{x^3}{a}} \\
 + \frac{x^3}{a} \\
 \underline{+ \frac{x^3}{a} - \frac{x^4}{aa}} \\
 + \frac{x^4}{aa}, \&c.
 \end{array}$$

Therefore

$$\frac{ax}{a-x} = x + \frac{xx}{a} + \frac{x^3}{aa} + \frac{x^4}{a^3} + \frac{x^5}{a^4} + \frac{x^6}{a^5}, \&c.$$

ad infinitum.

Ex. 2.

Let the fraction $\frac{aa}{b+x}$ be proposed.

$$b+x)aa+o\left(\frac{aa}{b} - \frac{aax}{bb} + \frac{aaxx}{b^3} - \frac{aax^3}{b^4} + \frac{aax^4}{b^5} \text{ \&c.}\right.$$

$$aa + \frac{aax}{b}$$

Answer.

$$- \frac{aax}{b} + o$$

$$- \frac{aax}{b} - \frac{aaxx}{bb}$$

$$+ \frac{aaxx}{bb}$$

$$+ \frac{aaxx}{bb} + \frac{a^2x^3}{b^3}$$

$$- \frac{a^2x^3}{b^3} \text{ \&c.}$$

Or thus,

$$x+b)aa+o\left(\frac{aa}{x} - \frac{aab}{x^2} + \frac{a^2b^2}{x^3} - \frac{a^2b^3}{x^4} \text{ \&c.}\right.$$

$$aa + \frac{baa}{x}$$

$$- \frac{baa}{x}$$

$$- \frac{aab}{x} - \frac{aabb}{xx}$$

$$+ \frac{aab^2}{xx}, \text{ \&c.}$$

Ex.

Ex. 3.

Suppose

$$\frac{1}{1+xx}$$

$$\frac{1+xx}{1+xx} = 1 + 0(1-xx+x^4-x^6+x^8-\dots) \&c.$$

$$\begin{array}{r} \hline -xx \\ -xx-x^4 \\ \hline +x^4 \\ +x^4+x^6 \\ \hline -x^6 \\ -x^6-x^8 \\ \hline +x^8 \&c. \end{array}$$

Ex. 4.

Let the fraction be $\frac{2x^{\frac{3}{2}} - x^{\frac{9}{2}}}{1+x^{\frac{1}{2}}-3x}$.

$$\frac{1+x^{\frac{1}{2}}-3x}{1+x^{\frac{1}{2}}-3x} = 2x^{\frac{3}{2}} - x^{\frac{9}{2}} + 0(2x^{\frac{1}{2}} - 2x + 7x^{\frac{3}{2}} - 13x^{\frac{5}{2}} + 34x^{\frac{7}{2}} \&c.)$$

$$\begin{array}{r} \hline -2x + 5x^{\frac{3}{2}} \\ -2x - 2x^{\frac{3}{2}} + 6x^2 \\ \hline +7x^{\frac{3}{2}} - 6x^2 \\ +7x^{\frac{3}{2}} + 7x^2 - 21x \\ \hline -13x^2 + 21x^{\frac{5}{2}} \\ -13x^2 - 13x^{\frac{5}{2}} \\ \hline +34x^{\frac{7}{2}} \&c. \end{array}$$

K 4

Ex.

Ex. 6.

Extract the cube root of $1-x^3$.

From $1-x^3$
 take 1 $\left(1 - \frac{x^3}{3} - \frac{x^6}{9} - \frac{5x^9}{81} \right. \&c.$

$$\begin{array}{r} \hline 3 \) -x^3 \\ \hline \end{array}$$

From $1-x^3$

take $1-x^3 + \frac{x^6}{3} - \frac{x^9}{27} = 1 - \frac{x^3}{3} \Bigg)^3$

$$\begin{array}{r} \hline 3 \) -\frac{x^6}{3} + \frac{x^9}{27} \\ \hline \end{array}$$

From $1-x^3$

take $1-x^3 + \frac{5x^9}{27} \&c. = 1 - \frac{x^3}{3} - \frac{x^6}{9} \Bigg)^3$

$$\begin{array}{r} \hline 3 \) -\frac{5x^9}{27} \&c. \\ \hline \end{array}$$

2 R U L E.

Assume a series with unknown coefficients, to represent it. Which series being multiplied, or involved, &c. according as the question requires; the quantities of the same dimension must be put equal to each other; from which equations, the coefficients will be determined.

Ex. 7.

Let $\frac{1}{a-x}$ be given.

Suppose $\frac{1}{a-x} = A + Bx + Cx^2 + Dx^3 + Ex^4 \&c.$
 the series required. Multiply by $a-x$.

Then

Then

$$1 = aA + aBx + aCx^2 + aDx^3 + aEx^4, \&c.$$

$$- Ax - Bx^2 - Cx^3 - Dx^4, \&c.$$

Whence equating the coefficients of the same powers of x , we have $aA = 1$, $aB - A = 0$, $aC - B = 0$, $aD - C = 0$, $aE - D = 0$, &c. Therefore $A = \frac{1}{a}$,

$$B = \frac{A}{a} = \frac{1}{aa}, \quad C = \frac{B}{a} = \frac{1}{a^3}, \quad D = \frac{C}{a} = \frac{1}{a^4}$$

$E = \frac{D}{a} = \frac{1}{a^5}$, &c. by reduction. Therefore the

series is $\frac{1}{a} + \frac{x}{aa} + \frac{x^2}{a^3} \&c.$ or $\frac{1}{a-x} = \frac{1}{a} +$
 $\frac{x}{aa} + \frac{x^2}{a^3} + \frac{x^3}{a^4} + \frac{x^4}{a^5} + \&c.$

Ex. 8.

Let $\frac{cc}{cc+2cy-yy}$ be given.

Suppose it $= A + By + Cy^2 + Dy^3, \&c.$ Multi-
 ply by $cc+2cy-yy$.

$$\text{Then } cc = ccA + ccBy + ccCy^2 + ccDy^3, \&c.$$

$$+ 2cAy + 2cBy^2 + 2cCy^3$$

$$- Ay^2 - By^3$$

And equating the homologous terms, $cc = ccA$,
 $ccB + 2cA = 0$, $ccC + 2cB - A = 0$,
 $ccD + 2cC - B = 0$, &c. and by reduction,

$$A = 1. \quad B = -\frac{2A}{c} = -\frac{2}{c}. \quad C = \frac{A - 2cB}{cc} =$$

$$\frac{1+4}{cc} = \frac{5}{cc}. \quad D = \frac{B - 2cC}{cc} = \frac{-2-10}{c^3} = -\frac{12}{c^3},$$

&c. Whence $\frac{cc}{cc+2cy-yy} = 1 - \frac{2y}{c} + \frac{5y^2}{cc} -$

$$\frac{12y^3}{c^3} \&c.$$

Ex.

Ex. 9.

What is $\sqrt{aa-xx}$.Let $\sqrt{aa-xx} = A + Bx^2 + Cx^4 + Dx^6, \&c.$ which being squared,

$$aa-xx = A^2 + 2ABx^2 + B^2x^4 + 2ADx^6, \&c. \\ + 2ACx^4 + 2BCx^6$$

Here $A^2 = aa$, $2AB = -1$, $BB + 2AC = 0$, $2AD + 2BC = 0$, &c. Whence $A = a$,

$$B = -\frac{1}{2A} = -\frac{1}{2a}, \quad C = -\frac{BB}{2A} = -\frac{1}{8a^3},$$

$$D = -\frac{BC}{A} = -\frac{1}{16a^5}, \&c. \text{ Therefore } \sqrt{aa-xx}$$

$$= a - \frac{xx}{2a} - \frac{x^4}{8a^3} - \frac{x^6}{16a^5} \&c.$$

PROBLEM LVIII.

To reduce any binomial surd to an infinite series, or to extract any root of a binomial.

R U L E.

This is done by substituting the particular letters or quantities, instead of these in the following general form, duly observing the signs.

$$P + PQ^{\frac{m}{n}} = P^{\frac{m}{n}} + \frac{m}{n} A Q + \frac{m-2n}{2n} C$$

$$BQ + \frac{m-2n}{3n} CQ + \frac{m-3n}{4n} DQ + \&c.$$

Where P is the first term, Q the second term divided by the first, $\frac{m}{n}$ the index of the power or root, A, B, C, D, &c. the foregoing terms with their signs.

Ex.

Ex. 1.

Extract the square root of $rr - xx$.

Here $P = rr$, $Q = \frac{-xx}{rr}$, $\frac{m}{n} = \frac{1}{2}$. Therefore $\sqrt{rr - xx} = r + \frac{1}{2} A \times \frac{-xx}{rr} - \frac{1}{4} B \times \frac{-xx}{rr} - \frac{3}{6} C \times \frac{-xx}{rr} - \frac{5}{8} D \times \frac{-xx}{rr} - \&c. = r - \frac{x}{2rr} A + \frac{xx}{4rr} B + \frac{3xx}{6rr} C + \frac{5x^3}{8rr} D + \&c.$ that is, restoring the values of A, B, C, &c. $\sqrt{rr - xx} = r - \frac{xx}{2r} - \frac{x^4}{8r^3} - \frac{x^6}{16rs} - \frac{5x^8}{128r^7} - \&c.$

Ex. 2.

What is the value of $\frac{rr}{r+x}$.

Here $\frac{rr}{r+x} = rr \times r+x^{-1}$, and $P = r$, $Q = \frac{x}{r}$, $\frac{m}{n} = -1$, or $m = -1$, $n = 1$. Therefore $r+x^{-1} = r - 1A \times \frac{x}{r} - 1B \times \frac{x}{r} - 1C \times \frac{x}{r} - 1D \times \frac{x}{r}, \&c. = \frac{r}{r} - \frac{x}{r} A - \frac{x}{r} B - \frac{x}{r} C \&c.$ And $rr \times r+x^{-1} = rr \times \left(\frac{r}{r} - \frac{x}{r} - \frac{x}{r^2} + \frac{xx}{r^3} - \frac{x^3}{r^4} \&c. \right)$ that is, $\frac{rr}{r+x} = r - x + \frac{xx}{r} - \frac{x^3}{r^2} + \frac{x^4}{r^3} - \&c.$

Ex.

Ex. 3.

To find the value of $\frac{1}{\sqrt{2rx-xx}}$.

$$\frac{1}{\sqrt{2rx-xx}} = \frac{1}{2rx-xx}^{-\frac{1}{2}}, \text{ and } P = 2rx,$$

$$Q = -\frac{x}{2r}, \quad m = -1, \quad n = 2. \text{ Then}$$

$$\frac{1}{2rx-xx}^{-\frac{1}{2}} = \frac{1}{2rx}^{-\frac{1}{2}} - \frac{1}{2} A \times \frac{-x}{2r} - \frac{3}{4} B \times \frac{-x}{2r}$$

$$- \frac{5}{6} C \times \frac{-x}{2r} - \frac{7}{8} D \times \frac{-x}{2r} \text{ \&c.} = \frac{1}{\sqrt{2rx}}$$

$$+ \frac{x}{4r} A + \frac{3x}{8r} B + \frac{5x}{12r} C + \frac{7x}{16r} D + \text{\&c.}$$

$$= \frac{1}{\sqrt{2rx}} + \frac{x}{4r\sqrt{2rx}} + \frac{3xx}{32rr\sqrt{2rx}} \text{ \&c.} = \frac{1}{\sqrt{2rx}}$$

$$\times: 1 + \frac{x}{4r} + \frac{3x^2}{32r^2} + \frac{3 \cdot 5x^3}{4 \cdot 8 \cdot 12r^3} + \frac{3 \cdot 5 \cdot 7x^4}{4 \cdot 8 \cdot 12 \cdot 16r^4} + \text{\&c.}$$

Ex. 4.

What is the cube root of $1-x^3$.

Here $P=1$, $Q=-x^3$, $m=1$, $n=3$. Whence

$$\frac{1}{1-x^3}^{\frac{1}{3}} = 1 + \frac{1}{3} A \times -x^3 - \frac{2}{6} B \times -x^3 -$$

$$\frac{5}{9} C \times -x^3 - \frac{8}{12} D \times -x^3 - \frac{11}{15} E \times -x^3 \text{ \&c.}$$

$$= 1 - \frac{x^3}{3} A + \frac{x^3}{3} B + \frac{5x^3}{9} C + \frac{2x^3}{3} D + \frac{11x^3}{15} E$$

$$\text{\&c. that is, } \sqrt[3]{1-x^3} = 1 - \frac{x^3}{3} - \frac{x^6}{9} - \frac{5x^9}{81}$$

$$- \frac{10x^{12}}{243} - \frac{22x^{15}}{729} \text{ \&c.}$$

Ex.

Ex. 5.

What is $\sqrt[3]{\frac{aa}{aa+xx}}$ in an infinite series.

This reduced is $a^{\frac{2}{3}} \times \overline{aa+xx}^{-\frac{2}{3}}$. Here $P=aa$,
 $Q = \frac{xx}{aa}$, $m = -2$, $n = 3$. And $\overline{aa+xx}^{-\frac{2}{3}}$
 $= aa^{-\frac{2}{3}} - \frac{2}{3} A \frac{xx}{aa} - \frac{5}{6} B \frac{xx}{aa} - \frac{8}{9} C \frac{xx}{aa} -$
 $\frac{11}{12} D \frac{xx}{aa} \&c. = \frac{1}{a^{\frac{1}{3}}} - \frac{2xx}{3a^{\frac{4}{3}}} + \frac{5x^4}{9a^{\frac{7}{3}}} - \frac{40x^6}{81a^{\frac{10}{3}}}$
 $+ \frac{110x^8}{243a^{\frac{13}{3}}} \&c. = \frac{1}{a^{\frac{1}{3}}} \times : \frac{1}{a} - \frac{2x^2}{3a^3} - \frac{5x^4}{9a^5}$
 $- \frac{40x^6}{81a^7} + \frac{110x^8}{243a^9} - \&c:$ therefore $\frac{a^2}{aa+xx}^{\frac{2}{3}}$
 $= \frac{1}{\sqrt[3]{aa}} \times : 1 - \frac{2x^2}{3a^2} + \frac{5x^4}{9a^4} - \frac{40x^6}{81a^6} + \&c.$

Ex. 6.

What is the value of $\sqrt[5]{aa-xx}$

$\sqrt[5]{aa-xx} = \overline{aa-xx}^{\frac{1}{5}}$. Here $P = a a$,
 $Q = \frac{-xx}{aa}$, $m = 1$, $n = 5$. Therefore $\overline{aa-xx}^{\frac{1}{5}}$
 $= \overline{aa}^{\frac{1}{5}} + \frac{1}{5} A \times \frac{-xx}{aa} - \frac{4}{10} B \times \frac{-xx}{aa} -$
 $\frac{9}{15} C \times \frac{-xx}{aa} - \frac{14}{20} D \times \frac{-xx}{aa} \&c. = a^{\frac{2}{5}} - \frac{xx}{5aa}$
 $A + \frac{2xx}{5aa} B + \frac{3xx}{5aa} C + \frac{7xx}{10aa} D \&c. = a^{\frac{2}{5}} \times :$
 $1 - \frac{xx}{5aa} - \frac{2x^4}{25a^4} - \frac{6x^6}{125a^6} - \frac{21x^8}{625a^8} \&c.$

Ex.

Ex. 7.

To reduce $\sqrt{a+x} \times \sqrt[4]{a-x}$ to a series.

$$\sqrt[4]{a-x} = \overline{a-x}^{\frac{1}{4}} \text{ Where } P=a, Q = \frac{-x}{a},$$

$$m=1, n=4. \text{ Then } \overline{a-x}^{\frac{1}{4}} = a^{\frac{1}{4}} + \frac{1}{4} A \times \frac{-x}{a} \\ - \frac{3}{8} B \times \frac{-x}{a} - \frac{7}{12} C \times \frac{-x}{a} \&c. = a^{\frac{1}{4}} - \frac{x}{4a} A \\ + \frac{3x^2}{8a^2} B + \frac{7x^3}{12a^3} C \&c. = a^{\frac{1}{4}} - \frac{x}{4a^{\frac{3}{4}}} - \frac{3x^2}{32a^{\frac{7}{4}}} \\ - \frac{7x^3}{128a^{\frac{11}{4}}} \&c.$$

Multiply by $a+x$

$$\text{Then } \frac{a^{\frac{5}{4}}}{a^{\frac{3}{4}}} - \frac{a^{\frac{1}{4}}x}{4} - \frac{3x^2}{32a^{\frac{3}{4}}} - \frac{7x^3}{128a^{\frac{7}{4}}} \&c. \\ + a^{\frac{1}{4}}x - \frac{x^2}{4a^{\frac{3}{4}}} - \frac{3x^3}{32a^{\frac{7}{4}}} \&c.$$

$$\sqrt{a+x} \times \sqrt[4]{a-x} = a^{\frac{5}{4}} + \frac{3a^{\frac{1}{4}}x}{4} - \frac{11x^2}{32a^{\frac{3}{4}}} - \frac{19x^3}{128a^{\frac{7}{4}}} \&c. \\ = \sqrt[4]{a} \times a + \frac{3}{4} x - \frac{11x^2}{32a} - \frac{19x^3}{128a^{\frac{3}{2}}} \&c.$$

Ex.

Ex. 8.

To designate $\sqrt{\frac{aa+xx}{aa-xx}}$ by a series.

$$\sqrt{aa+xx} = \overline{aa+xx}^{\frac{1}{2}}. \quad \text{Where } P = aa,$$

$$Q = \frac{xx}{aa}, \quad m=1, \quad n=2, \quad \text{and } \overline{aa+xx}^{\frac{1}{2}} = a +$$

$$\frac{1}{2} A \frac{xx}{aa} - \frac{1}{4} B \frac{xx}{aa} - \frac{3}{6} C \frac{xx}{aa} \quad \&c. = a +$$

$$\frac{xx}{2a} - \frac{x^4}{8a^3} + \frac{x^6}{16a^5} \quad \&c.$$

$$\text{Again, } \frac{1}{\sqrt{aa-xx}} = \overline{aa-xx}^{-\frac{1}{2}}. \quad \text{Here } P = aa,$$

$$Q = \frac{-xx}{aa}, \quad m=-1, \quad n=2. \quad \text{And } \overline{aa-xx}^{-\frac{1}{2}}$$

$$= \frac{1}{a} - \frac{1}{2} A \times \frac{-xx}{aa} - \frac{3}{4} B \times \frac{-xx}{aa} -$$

$$\frac{5}{6} C \times \frac{-xx}{aa} \quad \&c. = \frac{1}{a} + \frac{xx}{2a^3} +$$

$$\frac{3x^4}{8a^5} + \frac{5x^6}{16a^7} \quad \&c. \quad \text{Whence } \sqrt{\frac{aa+xx}{aa-xx}}$$

$$\text{or } \frac{\sqrt{aa+xx}}{\sqrt{aa-xx}} = a + \frac{xx}{2a} - \frac{x^4}{8a^3} + \frac{x^6}{16a^5} \quad \&c.$$

$$\times \frac{1}{a} + \frac{xx}{2a^3} + \frac{3x^4}{8a^5} + \frac{5x^6}{16a^7} \quad \&c. = 1 + \frac{xx}{aa} -$$

$$\frac{x^4}{2a^4} + \frac{x^6}{2a^6} \quad \&c. \quad \text{by multiplication.}$$

Ex. 9.

What is the value of $\frac{ax}{aa-ax+xx}$.

This may be treated as a binomial. Put $y=ax-xx$. Then $aa-ax+xx=aa-y$. And

$$\frac{ax}{aa-ax+xx} = ax \times \frac{1}{aa-y}^{-1}. \text{ Here } P=aa,$$

$$Q = \frac{-y}{aa}, \text{ } m = -1, \text{ } n = 1.$$

$$\text{And } \frac{1}{aa-y}^{-1} + \frac{1}{aa} - 1 A \times \frac{-y}{aa} - 1 B \times \frac{-y}{aa}$$

$$- 1 C \times \frac{-y}{aa}, - 1 D \times \frac{-y}{aa} \&c. = \frac{1}{aa} + \frac{y}{aa} A + \frac{y}{aa} B$$

$$+ \frac{y}{aa} C + \frac{y}{aa} D \&c. = \frac{1}{aa} + \frac{y}{a^2} + \frac{y^2}{a^3} + \frac{y^3}{a^4}$$

$$+ \frac{y^4}{a^5} \&c. = (\text{by restitution}) \frac{1}{aa} + \frac{ax-xx}{a^2}$$

$$+ \frac{ax-xx^2}{a^3} + \frac{ax-xx^3}{a^4} + \frac{ax-xx^4}{a^5} \&c. \text{ which in-}$$

olved and reduced into order will be

$$\frac{1}{aa} + \frac{x}{a^3} - \frac{xx}{a^4}$$

$$+ \frac{xx}{a^4} - \frac{2xx^2}{a^5} + \frac{x^4}{a^3}$$

$$+ \frac{x^3}{a^5} - \frac{3x^4}{a^6}, \&c.$$

$$+ \frac{x^4}{a^6}, \&c.$$

$$\frac{1}{aa} + \frac{x}{a^3} - \frac{xx}{a^4} - \frac{x^2}{a^5} - \frac{x^4}{a^3} \&c.$$

$$\text{and } \frac{ax}{aa-ax+xx} = \frac{x}{a} + \frac{x^2}{a^2} - \frac{x^4}{a^4} - \frac{x^5}{a^5} \&c.$$

The truth of this rule will appear by induction. For if any of these series be involved according to the index of the root, it will produce the original quantity. Thus if $r - \frac{rx}{2r} - \frac{x^2}{8r^2}$ &c. be squared, it produces $rr - xx$, as in Examp. 1. If $1 - \frac{x^2}{3} - \frac{x^6}{9}$ &c. be cubed it produces $1 - x^2$, Ex. 4. If $a^{\frac{2}{3}} \times : 1 - \frac{xx}{5aa} - \frac{2x^4}{25a^4}$ &c. be involved to the 5th power, it gives $aa - xx$, Ex. 5. and the like of others.

$$\text{Cor. 1. } \overline{P+PQ}^{\frac{m}{n}} = P^{\frac{m}{n}} \times : 1 + \frac{m}{n} Q + \frac{m}{n} \times \frac{m-n}{2n} Q^2 + \frac{m}{n} \times \frac{m-n}{2n} \times \frac{m-2n}{3n} Q^3 + \frac{m}{n} \times \frac{m-n}{2n} \times \frac{m-2n}{3n} \times \frac{m-3n}{4n} Q^4 \text{ \&c.}$$

$$\text{Cor. 2. } a+x = a^n + \frac{nx}{a+x} A + \frac{n+1}{2} \times \frac{x^2}{a+x} B + \frac{n+2}{3} \times \frac{x^3}{a+x} C + \frac{n+3}{4} \times \frac{x^4}{a+x} D + \text{\&c.}$$

where n is any index; A, B, C, &c. the foregoing terms with their signs.

For put $y = \frac{x}{a+x}$, then $x = \frac{ay}{1-y}$, and

$$a+x = \frac{a}{1-y}. \text{ Therefore } \overline{a+x}^n = \left(\frac{a}{1-y} \right)^n = a^n \times \overline{1-y}^{-n}.$$

Here $P=1$, $Q=-y$, $m=-n$, $n=1$ (see Prob. xlix); then by this problem,

$$\overline{1-y}^{-n} = 1 - \frac{n}{1} A \times -y - \frac{n+1}{2} B \times -y - \frac{n+2}{3} C \times -y - \frac{n+3}{4} D \times -y \text{ \&c.} = 1 + nyA$$

$$1 + nyA + \frac{n+1}{2} By + \frac{n+2}{3} Cy + \frac{n+3}{4} Dy$$

&c. and $a^n \times \sqrt[n]{1-y} = a^n \times (1 + nyA + \frac{n+1}{2} By + \frac{n+2}{3} Cy \&c.) =$ (restoring the value

of y) $a^n \times (1 + \frac{nx}{a+x} A + \frac{n+1}{2} \times \frac{x}{a+x} B + \frac{n+2}{3} \times \frac{x}{a+x} C + \frac{n+3}{4} \times \frac{x}{a+x} D \&c.)$

PROBLEM LIX.

To involve the series $ax + bx^2 + cx^3 + dx^4 + ez^4 \&c.$ to any power m , whole or fractional.

R U L E.

Substitute the particular letters or numbers in the given series, instead of these in the following general form.

$$\begin{aligned} & z \times \sqrt[m]{ax + bx^2 + cx^3 + dx^4 + ez^4 \&c.}^m \\ & = z^m \times \text{into} : a^m \\ & + \frac{mbA}{a} x \\ & + \frac{2mcA + m-1 \cdot bB}{2a} x^2 \\ & + \frac{3mdA + 2m-1 \cdot cB + m-2 \cdot bC}{3a} x^3 \\ & + \frac{4meA + m-1 \cdot dB + 2m-2 \cdot cC + m-3 \cdot bD}{4a} x^4 \\ & + \frac{5mfA + m-1 \cdot eB + 3m-2 \cdot dC + 2m-3 \cdot cD}{5a} x^5 \\ & + \frac{m-4 \cdot bE}{x^5} \end{aligned}$$

L 2 + 6mgA

$$\begin{aligned}
 & + \frac{6mgA + \overline{5m-1}.fB + \overline{4m-2}.eC + \overline{3m-3}.dD}{6a} \\
 & + \frac{\overline{2m-4}.cE + \overline{m-5}.bF}{x^5} \\
 & + \frac{7mbA + \overline{6m-1}.gB + \overline{5m-2}.fC + \overline{4m-3}.eD}{7a} \\
 & + \overline{3m-4}.dE + \overline{2m-5}.cF + \overline{m-6}.bG, \text{ \&c.}
 \end{aligned}$$

Where A, B, C, D, &c. are the coefficients of the terms immediately preceding those wherein they first appear. And the law of progression is evident.

Ex. 1.

What is the square of $1+x+x^2+x^3+x^4+\text{\&c.}$

Here $z=1$, $a=1$, $b=1$, $c=1$, $d=1$, &c.
 And $m=2$, then $\overline{1+x+x^2+x^3+\text{\&c.}}^2 =$

$$\begin{aligned}
 & 1 + \frac{2A}{1}x + \frac{4A+B}{2}x^2 + \frac{6A+3B+0}{3}x^3 + \\
 & + \frac{8A+5B+2C-D}{4}x^4 \text{ \&c.} \\
 & \quad \quad \quad A \quad B \quad C \quad D \quad E \\
 & = 1 + 2x + 3x^2 + 4x^3 + 5x^4 \text{ \&c.}
 \end{aligned}$$

Ex. 2.

What is the square root of $1+x+xx+x^3 \text{ \&c.}$

Here $z=1$, $a=1$, $b=1$, $c=1$, $d=1$, &c.
 and $m = \frac{1}{2}$. Whence $\overline{1+x+x^2+x^3+\text{\&c.}}^{\frac{1}{2}} =$

$$\begin{aligned}
 & = 1 + \frac{1}{2}Ax + \frac{A-\frac{1}{2}B}{2}x^2 + \frac{A+0-\frac{1}{2}C}{3}x^3 \\
 & + \frac{2A+\frac{1}{2}B-C-\frac{1}{2}D}{4}x^4, \text{ \&c.} = 1 + \frac{1}{2}x + \frac{3}{8}x^2 \\
 & + \frac{5}{16}x^3 + \frac{35}{128}x^4 \text{ \&c.}
 \end{aligned}$$

Ex.

Ex. 3.

Find the cube of $1+x+x^2+x^3 \&c.$

Here $z=1, a=1, b=1, c=1, d=1, e=1$
 &c. $m=3$. Then $1+x+x^2+x^3 \&c.^3 = 1 +$
 $\frac{3Ax}{1} + \frac{6A+2B}{2} x^2 + \frac{9A+5B+C}{3} x^3 +$
 $\frac{12A+8B+4C}{4} x^4 \&c. = 1+3x+6x^2+10x^3+$
 $15x^4 + \&c.$

Ex. 4.

What is the value of

$$\frac{1}{rr - \frac{1}{2}yy + \frac{y^4}{4r^2} - \frac{y^6}{8r^4} + \frac{y^8}{16r^6} \&c.}$$

Here $z=1, x=yy, a=rr, b=-\frac{1}{2}, c=\frac{1}{4rr^2}$
 $d=-\frac{1}{8r^4}, e=\frac{1}{16r^6} \&c. m=-1$. Then
 $rr - \frac{1}{2}yy + \frac{y^4}{4r^2} - \frac{y^6}{8r^4} \&c.]^{-1} = \frac{1}{rr} + \frac{A}{2rr} x +$
 $\frac{\frac{1}{2rr} A+B}{2rr} xx + \frac{\frac{3}{8r^4} A - \frac{3}{4rr} B + \frac{3}{2} C}{3rr} x^3 \&c.$
 $= \frac{1}{rr} + \frac{1}{2r^2} x + 0x^2 + 0x^3 \&c. = \frac{1}{rr} +$
 $\frac{1}{2r^2} x = \frac{1}{rr} + \frac{1}{2r^2} yy.$

Ex. 5.

To square the series $y-y^3+y^5-y^7+y^9 \&c.$

This is equal to $y:1-y^2+y^4-y^6+y^8 \&c.$
 Here $z=y, x=yy, a=1, b=-1, c=1, d=-1,$
 $e=1,$

$e=1$, &c. and $m=2$. Then $1-y^2+y^4 \&c.$ =
 $1-2Ax + \frac{4A-B}{2} x^2 + \frac{-6A+3B}{3} x^3 \&c.$ =
 $1-2x+3x^2-4x^3 \&c.$ and $y-y^3+y^5 \&c.$ =
 $y^2 \times 1-2x+3x^2 \&c.$ = $yy-2y^4+3y^6-4y^8 +$
 $\&c.$

Ex. 6.

To square the series

$$\sqrt{2r}x : v^{\frac{1}{2}} + \frac{v^{\frac{3}{2}}}{2 \cdot 2 \cdot 3r} + \frac{3v^{\frac{5}{2}}}{4 \cdot 2 \cdot 4 \cdot 5r^2} + \frac{3 \cdot 5 \cdot v^{\frac{7}{2}}}{8 \cdot 2 \cdot 4 \cdot 6 \cdot 7r^3} \&c.$$

The series is $2rv^{\frac{1}{2}} \times : 1 + \frac{v}{12r} + \frac{3v^2}{160r^2} +$
 $\frac{5v^3}{896r^3} \&c.$ Here $z = \sqrt{2rv}$, $a=1$, $b = \frac{1}{12r}$,
 $c = \frac{3}{160rr}$, $d = \frac{5}{896r^3} \&c.$ $m=2$. Then
 $2rv^{\frac{1}{2}} \times : 1 + \frac{v}{12r} + \frac{3v^2}{160r^2} \&c. = 2rv \times : 1 +$
 $+\frac{1}{6r} Ax + \frac{3}{40rr} A + \frac{1}{12r} B$
 $\frac{130}{896r^3} A + \frac{9}{160rr} B$
 $\frac{x}{6r} + \frac{2x^2}{45rr} + \frac{57x^3}{4480r^3} \&c.$

Ex. 7.

Find the m power of

$$ax^r + b x^{r+n} + c x^{r+2n} + d x^{r+3n} \&c.$$

This reduced is $x^r \times : a + b x^n + c x^{2n} +$
 $d x^{3n} \&c.$ Here $z = x^r$, $x = x^n$, $m=2$, $\&c.$
 Then

Then $x^r \times : a + bx^n + cx^{2n} \&c. \overset{m}{=} x^{rm} \times : a^m$
 $+ \frac{mb}{a} Ax^n + \frac{2mcA + m-1.bB}{2a} x^{2n} +$
 $\frac{3mdA + 2m-1.cB + m-2.bC}{3a} x^{3n} +$
 $\frac{4mcA + 3m-1.dB + 2m-2.cC + m-3.bD}{4a} x^{4n}$
 &c.

2 R U L E.

Substitute each letter in the given series, instead of the correspondent one, in the following general form.

$$z^m \times a + bx + cx^2 + dx^3 + ex^4 \&c. \overset{m}{=} z^m \times \text{into}$$

$$a^m + mba^{m-1}x + m \cdot \frac{m-1}{2} a^{m-2}bb \left. \begin{array}{l} \\ + ma^{m-1}c \end{array} \right\} x^2$$

$$+ m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} a^{m-3}b^3 \left. \begin{array}{l} \\ + m \cdot \frac{m-1}{2} \cdot 2a^{m-2}bc \\ + ma^{m-1}d \end{array} \right\} x^3$$

$$+ m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} a^{m-4}b^4 \left. \begin{array}{l} \\ + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot 3a^{m-3}bbc \\ + m \cdot \frac{m-1}{2} \cdot a^{m-2} \times \left\{ \begin{array}{l} 2bd \\ + cc \end{array} \right\} \\ + ma^{m-1}e \end{array} \right\} x^4$$

$$\begin{aligned}
 &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot \frac{m-4}{5} a^{m-5} b^5 \\
 &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot 4a^{m-4} b^3 c \\
 &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot 3a^{m-3} \left\{ \begin{array}{l} bcc \\ +bbd \end{array} \right\} \\
 &+m \cdot \frac{m-1}{2} \cdot 2a^{m-2} \left\{ \begin{array}{l} cd \\ be \end{array} \right\} \\
 &+ma^{m-1} f
 \end{aligned}
 \quad \left. \vphantom{\begin{aligned} &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot \frac{m-4}{5} a^{m-5} b^5 \\ &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot 4a^{m-4} b^3 c \\ &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot 3a^{m-3} \left\{ \begin{array}{l} bcc \\ +bbd \end{array} \right\} \\ &+m \cdot \frac{m-1}{2} \cdot 2a^{m-2} \left\{ \begin{array}{l} cd \\ be \end{array} \right\} \\ &+ma^{m-1} f \end{aligned}} \right\} x^5$$

$$\begin{aligned}
 &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot \frac{m-4}{5} \cdot \frac{m-5}{6} a^{m-6} b^6 \\
 &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot \frac{m-4}{5} \cdot 5a^{m-5} b^4 c \\
 &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot a^{m-4} \left\{ \begin{array}{l} 6bbcc \\ 4b^2 d \end{array} \right\} \\
 &+m \cdot \frac{m-2}{2} \cdot \frac{m-2}{3} \cdot a^{m-3} \left\{ \begin{array}{l} 3bbe \\ 6bcd \\ c^3 \end{array} \right\} \\
 &+m \cdot \frac{m-1}{2} a^{m-2} \left\{ \begin{array}{l} 2bf \\ 2ce \\ dd \end{array} \right\} \\
 &+ma^{m-1} g \\
 &\&c.
 \end{aligned}
 \quad \left. \vphantom{\begin{aligned} &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot \frac{m-4}{5} \cdot \frac{m-5}{6} a^{m-6} b^6 \\ &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot \frac{m-4}{5} \cdot 5a^{m-5} b^4 c \\ &+m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \cdot a^{m-4} \left\{ \begin{array}{l} 6bbcc \\ 4b^2 d \end{array} \right\} \\ &+m \cdot \frac{m-2}{2} \cdot \frac{m-2}{3} \cdot a^{m-3} \left\{ \begin{array}{l} 3bbe \\ 6bcd \\ c^3 \end{array} \right\} \\ &+m \cdot \frac{m-1}{2} a^{m-2} \left\{ \begin{array}{l} 2bf \\ 2ce \\ dd \end{array} \right\} \\ &+ma^{m-1} g \\ &\&c. \end{aligned}} \right\} x^5$$

For let $y = bx + cx^2 + dx^3 \ \&c.$ $p = \frac{m-1}{2} m,$

$$q = \frac{m-2}{3} p, r = \frac{m-3}{4} q, s = \frac{m-4}{5} r, \ \&c. \ \text{Then}$$

$$\overline{a + bx + cx^2 + dx^3}^m \ \&c. = a + y^m = a^m + ma^{m-1} y$$

$$+ pa^{m-2} y^2 + qa^{m-3} y^3 + ra^{m-4} y^4 \ \&c. \ \text{But}$$

$$y = bx$$

$$\begin{aligned}
 y &= bx + cx^2 + dx^3 + ex^4 \ \&c. \\
 yy &= bbxx + 2bcx^3 + 2bdx^4 + 2bex^5 \ \&c. \\
 &\quad + ccx^4 + 2cdx^5 \ \&c. \\
 y^3 &= b^3x^3 + 3bbcx^4 + 3bbdx^5 + \ \&c. \\
 &\quad + 3bccx^5 \\
 y^4 &= b^4x^4 + 4b^3cx^5 + \ \&c. \\
 y^5 &= b^5x^5 \\
 &\ \&c.
 \end{aligned}$$

Then the power $a^m + ma^{m-1}y + pa^{m-2}yy$
 $qa^{m-3}y^3 \ \&c.$ becomes

$$\begin{aligned}
 &a^m \\
 + ma^{m-1} \times &: bx + cx^2 + dx^3 + ex^4 + fx^5 \ \&c. \\
 + pa^{m-2} \times &: \quad bbx^2 + 2bcx^3 + 2bdx^4 + 2bex^5 \\
 &\quad + cc + 2cd \\
 + qa^{m-3} \times &: \quad b^3x^3 + 3b^2cx^4 + 3bbdx^5 \\
 &\quad + 3bcc \\
 + ra^{m-4} \times &: \quad b^4x^4 + 4b^3cx^5 \\
 + sa^{m-5} \times &: \quad b^5x^5 \\
 &\ \&c.
 \end{aligned}$$

These being actually multiplied, and the coefficients of each power of x collected; will give the several terms as in the form above.

And the first Rule is in effect the same as this.

For let $a + bx + cxx + dx^3 \ \&c.^m = A + Bx^2 + Cx^3$

$+ D^4 \ \&c.$ Then by Rule 1, $A = a^m$, as in Rule

2d. Also $B = \frac{mbA}{a} = mba^{m-1}$, as in Rule 2d.

Likewise $C = \frac{mcA}{a} + \frac{m-1}{2a} \cdot bB = mca^{m-1} +$

$\frac{m-1}{2} \cdot mbba^{m-2}$, as in Rule 2d.

Again

$$\begin{aligned}
 \text{Again } D &= \frac{mdA}{a} + \frac{2m-1 \cdot cB}{3a} + \frac{m-2 \cdot bC}{3a} \\
 &= mda^{m-1} + \frac{2m-1}{3} \times mcb a^{m-2} + \frac{m-2}{3} \cdot b \times \\
 &\quad \frac{mca^{m-2} + \frac{m-1}{2} \cdot mbba^{m-3}}{3} = mda^{m-1} + \\
 &\quad \frac{2m-1}{3} + \frac{m-2}{3} \times mbca^{m-2} + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} b^2 \\
 &\quad a^{m-3} = mda^{m-1} + m \cdot \frac{m-1}{2} \cdot 2bca^{m-2} + \\
 &\quad + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} b^2 a^{m-3}, \text{ as in Rule 2d, and} \\
 &\text{so for the rest. In using this last rule, it will} \\
 &\text{be the easiest way to divide all by the first term,} \\
 &\text{that } a \text{ may be 1.}
 \end{aligned}$$

Ex. 8.

What is the fourth power of $1+x+x^2+x^3$ &c.
 Here $z=1$, $a=1$, $b=1$, $c=1$, $d=1$, &c.
 $m=4$. Then $1+x+x^2+x^3$ &c.⁴

$$\begin{aligned}
 &= 1 + 4bx + 6b^2x^2 + 4b^3x^3 + b^4x^4 \text{ &c.} \\
 &\quad + 4c \quad + 12bc \quad + 12b^2c \\
 &\quad \quad + 4d \quad + 6cc \\
 &\quad \quad \quad + 12bd \\
 &\quad \quad \quad + 4e \\
 &= 1 + 4x + 10x^2 + 20x^3 + 35x^4 \text{ &c.}
 \end{aligned}$$

Ex. 9.

What is the square of $\frac{1}{x} + \frac{1}{xx} + \frac{1}{x^3} + \frac{1}{x^4}$ &c.

In this Example, $z = \frac{1}{x}$, $x = \frac{1}{x}$, $a=1$, $b=1$,
 $c=1$, $d=1$, &c. $m=2$.

Then

Then $\overbrace{\frac{1}{x} + \frac{1}{xx} + \frac{1}{x^3} \&c.}^2 = \frac{1}{xx} \times :$

$$1 + 2b \times \frac{1}{x} + bb \times \frac{1}{xx} + 2bc \times \frac{1}{x^3} + 2bd \times \frac{1}{x^4} \&c.$$

$$+ 2c \quad + 2d \quad + 2e$$

$$= \frac{1}{xx} \times : 1 + \frac{2}{x} + \frac{3}{xx} + \frac{4}{x^3} + \frac{5}{x^4} \&c. = \frac{1}{xx}$$

$$+ \frac{2}{x^2} + \frac{3}{x^3} + \frac{4}{x^4} + \frac{5}{x^5} \&c.$$

Ex. 10.

To square the series $y - y^3 + y^5 - y^7 + \&c.$

Here $z = y$. $a = 1, b = 0, c = -1, d = 0, e = 1,$
 $f = 0, g = -1 \&c.$ and $m = 2$. Whence

$$\overbrace{y - y^3 + y^5 \&c.}^2 = y^2 \times :$$

$$1 + 0x - 2ax^2 - 0x^3 + ccx^4$$

$$+ 2e$$

$$+ 0x^5 - 2ccx^6 \&c. = y^2 \times : 1 - 2y^2 + 3y^4 - 4y^6 \&c.$$

$$- 2g$$

$$= y^2 - 2y^4 + 3y^6 - 4y^8 \&c.$$

Or thus,

$z = y, x = yy, a = 1, b = -1, c = 1, d = -1, \&c.$

and $m = 2$. Then $\overbrace{y - y^3 + y^5 \&c.}^2 = yy \times :$

$$1 + 2bx + bbx^2 + 2bcx^3 + 2bdx^4 \&c.$$

$$+ 2c \quad + 2d \quad + 2e$$

$$+ 2c$$

$$= y^2 \times : 1 - 2y^2 + 3y^4 - 4y^6 + 5y^8 \&c. = y^2 - 2y^4$$

$$+ 3y^6 - 4y^8 + 5y^{10} \&c.$$

Ex.

Ex. II.

What is the square root of $rr - zz + \frac{z^4}{3r^2} - \frac{2z^6}{45r^4}$
 $+ \frac{z^8}{315r^6} - \&c.$

Here $z=1$, $x=zz$, $a=rr$, $b=-1$, $c=\frac{1}{3r^2}$,
 $d=\frac{-2}{45r^4}$, $e=\frac{1}{315r^6}$ &c. and $m=\frac{1}{2}$,
 $\frac{m-1}{2} = -\frac{1}{4}$, $\frac{m-2}{3} = -\frac{1}{2}$, $\frac{m-3}{4} = -\frac{5}{8}$
 &c. Then $rr - zz + \frac{z^4}{3rr}$ &c. $\left. \right)^{\frac{1}{2}} = r + \frac{1}{2} \times \frac{-1}{r} x$
 $-\frac{1}{8r^3} x^2 + \frac{1}{6r^3} x^2$ &c. $= r - \frac{zz}{2r} + \frac{z^4}{24r^3}$ &c.

Rather thus,

The quantity reduced is $rr \times : 1 - \frac{zz}{rr} + \frac{z^4}{3r^4}$
 $-\frac{2z^6}{45r^6}$ &c. Here $z=rr$, $a=1$, $b=-\frac{1}{rr}$,
 $c=\frac{1}{3r^4}$, $d=\frac{-2}{45r^6}$ &c. Whence
 $rr - zz + \frac{z^4}{3rr}$ &c. $\left. \right)^{\frac{1}{2}} = r \times : 1 - \frac{x}{2rr}$
 $\left. \begin{array}{l} + \frac{1}{8r^4} \\ + \frac{1}{6r^4} \end{array} \right\} x^2 - \frac{1}{16r^6}$
 $\left. \begin{array}{l} + \frac{1}{12r^6} \\ - \frac{1}{45r^6} \end{array} \right\} x^3$ &c. $= r \times : 1 - \frac{x}{2rr}$
 $+ \frac{1}{24r^4} x^2 - \frac{1}{720r^6} x^3 + \&c.$

$= r - \frac{zz}{1.2r} + \frac{z^4}{1.2.3.4r^3} - \frac{z^6}{1.2.3.4.5.6r^5} + \&c.$

Ex.

Ex. 12.

What is the square root of

$$\frac{1}{rr} - \frac{zz}{2} + \frac{z^4}{4rr} - \frac{z^6}{6r^4} + \frac{z^8}{8r^6} \text{ \&c.}$$

The quantity reduced is

$$\frac{1}{rr} \times \frac{1}{1 - \frac{zz}{2rr} + \frac{z^4}{4r^4} - \frac{z^6}{6r^6} \text{ \&c.}}$$

Where $z = \frac{1}{rr}$, $x = zz$, $a = 1$, $b = -\frac{1}{2rr}$,

$c = \frac{1}{4r^4}$, $d = -\frac{1}{6r^6}$, &c. and $m = -\frac{1}{2}$,

$\frac{m-1}{2} = -\frac{3}{4}$, $\frac{m-2}{3} = -\frac{5}{6}$, $\frac{m-3}{4} = -\frac{7}{8}$ &c.

And $\sqrt{\frac{1}{rr - \frac{zz}{2} + \frac{z^4}{4rr} \text{ \&c.}}} = \frac{1}{r} \times :$

$$1 + \frac{x}{4rr} + \frac{2x^2}{32r^4} + \frac{5}{128r^6} x^4 \text{ \&c.}$$

$$- \frac{1}{8r^4} - \frac{3}{32r^6}$$

$$+ \frac{1}{12r^6}$$

$$= \frac{1}{r} + \frac{x}{4r^3} - \frac{x^2}{32r^5} + \frac{11x^3}{384r^7} - \text{\&c.}$$

SCHOLIUM.

From this problem the powers of a compound quantity are deduced as follows, which will be serviceable upon particular occasions.

If

If $y = A + B + C + D \text{ \&c.}$ Then

$$y = A + B + C + D + E + F + G + H$$

$$y^2 = A^2 + 2AB + 2AC + 2AD + 2AE + 2AF + 2AG + 2AH, \text{ \&c.}$$

$$+ BB + 2BC + 2BD + 2BE + 2BF + 2BG$$

$$+ CC + 2CD + 2CE + 2CF$$

$$+ DD + 2DE$$

$$y^3 = A^3 + 3A^2B + 3A^2C + 3A^2D + 3A^2E + 3A^2F + 3A^2G, \text{ \&c.}$$

$$+ 3ABB + 6ABC + 6ABD + 6ABE + 6ABF$$

$$+ B^3 + 3ACC + 6ACD + 6ACE$$

$$+ 3BBC + 3BBD + 6BCD$$

$$+ 3BCC + 3ADD$$

$$+ 3BBE$$

$$+ C^3$$

$$y^4 = A^4 + 4A^3B + 6A^2B^2 + 4AB^3 + 6A^2C^2 + 4B^3C, \text{ \&c.}$$

$$+ 4A^3C + 12A^2BC + 12AB^2C + 12ABC^2$$

$$+ 4A^3D + 12A^2BD + 12AB^2D$$

$$+ 4A^3E + 12A^2CD$$

$$+ B^4 + 12A^2BE$$

$$+ 4A^3F$$

$$y^5 = A^5 + 5A^4B + 10A^3B^2 + 10A^2B^3 + 5AB^4 + B^5, \text{ \&c.}$$

$$+ 5A^4C + 20A^3BC + 30A^2B^2C + 20AB^3C$$

$$+ 5A^4D + 20A^3BD + 30A^2BC^2$$

$$+ 10A^3CC + 30A^2B^2D$$

$$+ 5A^4E + 20A^3CD$$

$$+ 20A^3BE$$

$$+ 5A^4F$$

$$y^6 = A^6 + 6A^5B + 15A^4B^2 + 20A^3B^3 + 15A^2B^4, \text{ \&c.}$$

$$+ 6A^5C + 30A^4BC + 60A^3B^2C$$

$$+ 6A^5D + 30A^4BD$$

$$+ 15A^4CC$$

$$+ 6A^5E$$

$$y^7 = A^7 + 7A^6B + 21A^5B^2 + 35A^4B^3 + 35A^3B^4, \text{ \&c.}$$

$$+ 7A^6C + 42A^5BC + 105A^4B^2C$$

$$+ 7A^6D + 42A^5BD$$

$$+ 21A^5CC$$

$$+ 7A^6E$$

$$y^8 = A^8 + 8A^7B + 28A^6B^2 + 56A^5B^3 + 70A^4B^4, \text{ \&c.}$$

$$+ 8A^7C + 56A^6BC + 168A^5B^2C$$

$$+ 8A^7D + 56A^6BD$$

$$+ 28A^6CC$$

$$+ 8A^7E$$

$$y^9 = A^9$$

Sect. VI. INFINITE SERIES. 159

$$y^9 = A^9 + 9A^8B + 36A^7B^2 + 84A^6B^3 + 126A^5B^4 \text{ \&c.}$$

$$+ 9A^8C + 72A^7BC + 252A^6B^2C$$

$$+ 9A^8D + 72A^7BD$$

$$+ 36A^7CC$$

$$+ 8A^8E$$

$$y^{10} = A^{10} + 10A^9B + 45A^8B^2 + 120A^7B^3 + 210A^6B^4 \text{ \&c.}$$

$$+ 10A^9C + 90A^8BC + 360A^7B^2C$$

$$\text{\&c.} + 10A^9D + 90A^8BD$$

$$+ 45A^8CC$$

$$+ 10A^9E$$

In making use of any of these forms, the terms of the given series must be ranged in order (Prob. xlviii.), and the whole terms thereof substituted one by one, in the room of the quantities A, B, C, D, &c. (Prob. xlix).

Ex. 1.

Let $a + bx + cx^2 + dx^3 + ex^4$ &c. be cubed.

$$A + B + C + D + E \text{ \&c.}$$

$$= a + bx + cx^2 + dx^3 + ex^4 \text{ \&c.}$$

that is $A = a$, $B = bx$, &c. Then

$$(y^3) A^3 + 3A^2B + 3A^2C \text{ \&c.} =$$

$$+ 3AB^2$$

$$a^3 + 3a^2bx + 3aacx^2 + 3aadx^3 + 3aaex^4 \text{ \&c.}$$

$$+ 3abbx^2 + 6abcx^3 + 6abdx^4$$

$$+ b^3x^3 + 3accx^4$$

$$+ 3bbcx^4$$

Ex. 2.

What is the fourth power of

$$x - \frac{2}{x} + \frac{p}{x^3} - \frac{2cd}{x^5}$$

$$A + B + C + D$$

$$= x - \frac{2}{x} + \frac{p}{x^3} - \frac{2cd}{x^5}$$

Then

$$\begin{aligned} \text{Then } y^4 = x^4 - 4x^3 \frac{2}{x} + 6xx \times \frac{4}{xx} - 4x \times \frac{8}{x^3} \&c. = \\ &+ 4x^3 \times \frac{p}{x^3} - 12xx \times \frac{2p}{x} \\ &\quad - 4x^3 \times \frac{2cd}{x^5} \\ x^4 - 8x^2 + 24 + 4p - \frac{32 + 24p + 8cd}{xx} \&c. \end{aligned}$$

Ex. 3.

Involve $2x^{\frac{1}{2}} + 3x^{\frac{5}{2}} - 4x^{\frac{7}{2}} + 5x^{\frac{9}{2}} - 6x^{\frac{11}{2}} \&c.$ to the 5th power.

$$\begin{aligned} A + B + C + D + E \&c. = y. \\ = 2x^{\frac{1}{2}} + 3x^{\frac{5}{2}} - 4x^{\frac{7}{2}} + 5x^{\frac{9}{2}} - 6x^{\frac{11}{2}} \&c. = y. \\ y^5 = 32x^{\frac{5}{2}} + 80x^{\frac{4}{2}} \times 3x^{\frac{5}{2}} + 80x^{\frac{3}{2}} \times 9x^{\frac{10}{2}} \\ &+ 80x^{\frac{4}{2}} \times 4x^{\frac{7}{2}} \\ - 160x^{\frac{3}{2}} \times 12x^{\frac{12}{2}} - \\ + 80x^{\frac{4}{2}} \times 5x^{\frac{9}{2}} \&c. \\ 80x^{\frac{4}{2}} \times 6x^{\frac{11}{2}} \&c. = 32x^{\frac{5}{2}} + 240x^{\frac{9}{2}} + 720x^{\frac{13}{2}} - \\ - 320x^{\frac{11}{2}} + \\ 1920x^{\frac{15}{2}} - 480x^{\frac{15}{2}} \&c. \text{ that is} \\ 420x^{\frac{13}{2}} \&c. \end{aligned}$$

$$\begin{aligned} y^5 = 32x^{\frac{5}{2}} + 240x^{\frac{9}{2}} - 320x^{\frac{11}{2}} + 720x^{\frac{13}{2}} - 1920x^{\frac{15}{2}} \\ + 400x^{\frac{13}{2}} - 480x^{\frac{15}{2}} \\ \&c. \end{aligned}$$

Or $y^5 = 32x^{\frac{5}{2}} + 240x^{\frac{9}{2}} - 320x^{\frac{11}{2}} + 1120x^{\frac{13}{2}} - 2400x^{\frac{15}{2}} \&c.$ Here I omit all these terms, where I see the index of x exceeds $\frac{15}{2}$.

Or.

Or thus,

$$A + B + C + D + E + F \text{ \&c.}$$

$$= 2x^{\frac{1}{2}} + 0 + 3x^{\frac{3}{2}} - 4x^{\frac{5}{2}} + 5x^{\frac{7}{2}} - 6x^{\frac{9}{2}} \text{ \&c. then}$$

$$y^5 = 32x^{\frac{5}{2}} + 0 + 80x^{\frac{7}{2}} - 80x^{\frac{4}{2}} \times 4x^{\frac{7}{2}} + 80x^{\frac{3}{2}} \times 9x^{\frac{10}{2}} + 80x^{\frac{4}{2}} \times 5x^{\frac{9}{2}}$$

$$- 160x^{\frac{3}{2}} \times 12x^{\frac{12}{2}} \text{ \&c.} = 32x^{\frac{5}{2}} + 240x^{\frac{7}{2}} - 320x^{\frac{11}{2}}$$

$$- 80x^{\frac{4}{2}} \times 6x^{\frac{11}{2}} + 1120x^{\frac{11}{2}} - 2400x^{\frac{13}{2}} \text{ \&c.}$$

Ex. 4.

If $y = 1 + x^3 - 2x$, what is y^8 .

$$A + B + C + D + E + F \text{ \&c.}$$

$$= 1 - 2x + x^3 + 0 + 0 \text{ \&c.}$$

$$y^8 = 1 - 16x + 28 \times 4x^2 - 56 \times 8x^3 + 70 \times 16x^4 \text{ \&c.} + 8x^3 - 56 \times 2x^4 + 168 \times 4x^5 + 28x^6$$

$$= 1 - 16x + 112x^2 - 448x^3 + 1120x^4 \text{ \&c.} + 8x^3 - 112x^4 + 672x^5 + 28x^6$$

$$y^8 = 1 - 16x + 112x^2 + 8x^3 - 112x^4 \text{ \&c. that is,} - 448x^3 + 1120x^4$$

$y^8 = 1 - 16x + 112x^2 - 440x^3 - 1008x^4 \text{ \&c.}$ This is supposing x to be very small; but when x is very great, then x^3 must begin the series;

Thus,

$$A + B + C + D + E + F \text{ \&c.}$$

$$= x^3 + 0 - 2x + 1 + 0 \text{ \&c. Then}$$

$$y^8 = x^{24} + 0 - 8x^{21} \times 2x + 8x^{21} \times 1 + 28x^{18} \times 4xx$$

$$\text{or } y^8 = x^{24} - 16x^{22} + 8x^{21} + 112x^{20} \text{ \&c.}$$

P R O B L E M L X.

To abridge an infinite series, or denote it in a short manner for working.

When a series consists of terms very much compounded, or having a great many factors; it is very laborious to reduce them into numbers. And when several factors in any term are contained in the succeeding terms; the work may be shortened, by making use of the preceding term or some part of it, instead of such factors as are equivalent to it, in the following terms; as follows.

R U L E.

Put A, B, C, D, &c. for the first, second, third, fourth, &c. terms of the given series. Then to get the coefficients thereof, divide every term by the preceding one, gives the coefficient of that term. Whence you will have a new series equal to the former, and shorter designated.

Ex. 1.

$$\text{If } z + \frac{z^3}{2a^2} + \frac{3z^5}{2.4a^4} + \frac{3.5z^7}{2.4.6a^6} + \frac{3.5.7z^9}{2.4.6.8a^8} \&c. = y.$$

$$\text{Then } z \left(\frac{z^3}{2a^2} \right) \left(= \frac{z^4}{2aa} = \text{coefficient of B} = \frac{B}{A} \right).$$

$$\frac{z^3}{2a^2} \left(\frac{3z^5}{2.4a^4} \right) \left(\frac{3z^2}{4a^2} = \text{coefficient of C} = \frac{C}{B} \right).$$

$$\frac{3z^5}{2.4a^4} \left(\frac{3.5z^7}{2.4.6a^6} \right) \left(\frac{5z^2}{6a^2} = \text{coefficient of D} = \frac{D}{C} \right) \&c.$$

Hence the series becomes

$$z + \frac{z^4}{2aa} A + \frac{3z^2}{4aa} B + \frac{5z^2}{6aa} C + \frac{7z^2}{8aa} D \&c. = y.$$

Ex.

Ex. 2.

Suppose $1 + \frac{v}{1.3} + \frac{v^2}{1.3.5} + \frac{v^3}{3.5.7} + \frac{v^4}{5.7.9} + \frac{v^5}{7.9.11} \&c. = y.$

Here $\frac{B}{A} = \frac{v}{3}, \frac{C}{B} = \frac{v}{5}, \frac{D}{C} = \frac{v}{7}, \frac{E}{D} = \frac{3v}{9}, \frac{F}{E} = \frac{5v}{11}, \&c.$

Then the series is,

$1 + \frac{v}{3} A + \frac{v}{5} B + \frac{v}{7} C + \frac{3v}{9} D + \frac{5v}{11} E + \frac{7v}{13} F, \&c. = y.$

Ex. 3.

Let $x = \frac{3x^2}{1.2} - \frac{5x^3}{1.2.3.4} - \frac{7x^4}{1.2.3.4.5.6} \&c.$
be given.

$x \left(\frac{3x^2}{1.2} \left(\frac{-3x}{1.2} - \frac{-3x^2}{1.2} \right) - \frac{5x^3}{1.2.3.4} \left(\frac{5x}{3.3.4} - \frac{-5x^3}{1.2.3.4} \right) - \frac{7x^4}{1.2.3.4.5.6} \left(\frac{7x}{5.5.6} \&c. \right) \right)$

And the series is

$x - \frac{3x}{2} A + \frac{5x}{3.3.4} B + \frac{7x}{5.5.6} C \&c.$

Or thus,

$\left(\frac{-3x^2}{2} - \frac{-5x^3}{2.3.4} \left(\frac{5x}{3.4} - \frac{-5x^3}{2.3.4} \right) - \frac{7x^4}{2.3.4.5.6} \left(\frac{7x}{5.6} \&c. \right) \right)$

And the series

$$= x - \frac{3x}{2} A + \frac{\frac{3x}{2}}{3 \cdot 4} B + \frac{\frac{3x}{2}}{5 \cdot 6} C \text{ \&c.}$$

Where A, B, C, \&c. are the foregoing terms with their signs.

Ex. 4.

$$\text{Suppose } bx - \frac{bx^3}{2 \cdot 3aa} - \frac{bx^5}{5 \cdot 2 \cdot 4a^4} - \frac{bx^7}{7 \cdot 2 \cdot 4 \cdot 6a^6} - \frac{bx^9}{9 \cdot 2 \cdot 4 \cdot 6 \cdot 8a^8} \text{ \&c.} = d.$$

$$\text{Then } bx \left(\frac{-bx^3}{2 \cdot 3aa} \right) \left(\frac{-2x}{2 \cdot 3aa} \right) = \text{coefficient of B.}$$

$$\left(\frac{-bx^3}{2 \cdot 3aa} \right) \left(\frac{-bx^5}{5 \cdot 2 \cdot 4a^4} \right) \left(\frac{3x}{4 \cdot 5aa} \right) = \text{coefficient of C.}$$

$$\left(\frac{-bx^5}{5 \cdot 2 \cdot 4a^4} \right) \left(\frac{-bx^7}{7 \cdot 2 \cdot 4 \cdot 6a^6} \right) \left(\frac{5x}{6 \cdot 7aa} \right) = \text{coef. of D, \&c.}$$

And the series is

$$bx - \frac{2x}{2 \cdot 3aa} A + \frac{3x^2}{4 \cdot 5aa} B + \frac{5x^2}{6 \cdot 7a^2} C + \frac{7x^2}{8 \cdot 9aa} D \text{ \&c.}$$

2 R U L E.

If there be some single factor or factors, which are not in all the terms; set them aside at present. Then put A, B, C, D, \&c. for the remaining terms; and proceed as before. And at last restore these single factors into their proper terms.

Ex.

Ex. 5.

$$\text{If } x - \frac{3x^2}{1.2} - \frac{5x^3}{1.2.3.4} - \frac{7x^4}{1.2.3.4.5.6} - \frac{9x^5}{1.2.3.4.5.6.7.8} \&c. = y.$$

Here the factors 3, 5, 7, 9, &c. are not in all the terms, and being left out, the series is

$$x - \frac{x^2}{1.2} - \frac{x^3}{1.2.3.4} - \frac{x^4}{1.2.3.4.5.6} \&c.$$

abridged to $x - \frac{x}{1.2} A + \frac{x}{3.4} B + \frac{x}{5.6} C + \frac{x}{7.8} D$ &c. and the factors restored, the series becomes

$$x - \frac{x}{1.2} A \times 3 + \frac{x}{3.4} B \times 5 + \frac{x}{5.6} C \times 7 + \frac{x}{7.8} D \times 9 \&c. = y.$$

Where A, B, C, &c. are the several terms with their proper signs; without the numbers, 3, 5, 7, &c.

Ex. 6.

$$\text{If } bz - \frac{bz^3}{3.2aa} + \frac{bz^5}{5.2.4a^4} - \frac{bz^7}{7.2.4.6a^6} + \frac{bz^9}{9.2.4.6.8a^8} \&c. = y.$$

Then the factors 3, 5, 7, 9 &c. not being common to all the terms, are left out, and the series is

$$bz - \frac{bz^3}{2aa} + \frac{bz^5}{2.4a^4} - \frac{bz^7}{2.4.6a^6} \&c.$$

$$= bz - \frac{zz}{2aa} A - \frac{zz}{4aa} B - \frac{zz}{6aa} C - \frac{zz}{8aa} D \&c.$$

And restoring the numbers, the series will then be

$$bz \quad \frac{zx^2}{2aa} A \quad \frac{zx^4}{4aa} B \quad \frac{zx^6}{6aa} C \quad \frac{zx^8}{8aa} D$$

$$\frac{1}{3} \quad \frac{5}{7} \quad \frac{9}{9}$$

$\&c. = y$. Where A, B, C, $\&c.$ are the foregoing numerators, with their proper signs.

Ex. 7.

There is given

$$x - \frac{ax^3}{3 \cdot 2} + \frac{bx^5}{5 \cdot 2 \cdot 4} - \frac{cx^7}{7 \cdot 2 \cdot 4 \cdot 6} + \frac{dx^9}{9 \cdot 2 \cdot 4 \cdot 6 \cdot 8} \&c.$$

curtailed, $x - \frac{x^3}{2} + \frac{x^5}{2 \cdot 4} - \frac{x^7}{2 \cdot 4 \cdot 6} + \frac{x^9}{2 \cdot 4 \cdot 6 \cdot 8} \&c.$

or shortened, $x - \frac{xx}{2} A - \frac{xx}{4} B - \frac{xx}{6} C - \frac{xx}{8} D \&c.$

compleat, $x - \frac{xx}{2} A \times \frac{a}{3} - \frac{xx}{4} B \times \frac{b}{5} -$
 $\frac{xx}{6} C \times \frac{c}{7} - \frac{xx}{8} D \times \frac{d}{9} \&c.$

Where A, B, C, $\&c.$ are the foregoing terms, exclusive of the following quantities.

Cor. 1. If the first term of any transformed series be multiplied by any number or quantity; the whole series is multiplied thereby. For the first term is virtually contained in all the following terms. This is made plain by Ex. 4.

Cor. 2. In like manner A, B, C may be made to stand only for the coefficients, or otherwise, as any one pleases.

PROBLEM LXI.

To find the finite value of an infinite series, or what surd it is involved from.

R U L E.

Divide all the terms by the first; then the first term will be 1. Then compare three terms of this series with three terms of the series Rule 2, Prob. lix. each with each, supposing a to be 1, and c , &c. 0; which two equations will find the index, and the second term, if it is a binomial. If this does not succeed, compare four terms with four, for a trinomial; or five terms with five, for a quadrinomial; making $d=0$, or $e=0$, &c.

Ex. I.

Suppose this series $1 - \frac{y}{a} + \frac{y^2}{aa} - \frac{y^3}{a^3} + \frac{y^4}{a^4}$ &c.

Compare this with $\dots 1 + mbx + m \cdot \frac{m-1}{2} b^2 x^2$.

Then $mbx = -\frac{y}{a}$, and $m \cdot \frac{m-1}{2} b^2 x^2 = \frac{yy}{aa}$,

and dividing the last by the first, $\frac{m-1}{2} b^2 x^2 = -\frac{y}{a}$

$= mbx$; therefore $\frac{m-1}{2} = m$, and $2m = m-1$,

whence $m = -1$. Therefore $mbx = -bx = -\frac{y}{a}$,

or $bx = \frac{y}{a}$. Whence the index is -1 , and

the second term of the binomial (if it is one) is

$\frac{y}{a}$. And the binomial $1 + \frac{y}{a}$, or $1 + \frac{y}{a}$

that is $\frac{a}{a+y}$ the root required; which succeeds.

Ex. 2.

Suppose $a + \frac{xx}{2a} - \frac{x^4}{8a^3} + \frac{x^6}{16a^5} \&c.$

Reduced $a \times : 1 + \frac{xx}{2aa} - \frac{x^4}{8a^4} \&c.$

Rule $1 + mbx + m \cdot \frac{m-1}{2} bbxx.$

Here $mbx = \frac{xx}{2aa}$, and $m \cdot \frac{m-1}{2} bbxx = -\frac{x^4}{8a^4}$,

and by division, $\frac{m-1}{2} bx = -\frac{xx}{4aa}$. Then

$xx = mbx \times 2aa = -4aa \times \frac{m-1}{2} bx$; whence

$-m = m-1$, and $2m = 1$, or $m = \frac{1}{2}$ the index.

And $mbx = \frac{1}{2} bx = \frac{xx}{2aa}$, or $bx = \frac{xx}{aa}$ the se-

cond term. And the surd is $a \times 1 + \frac{xx}{aa}^{\frac{1}{2}}$ or $\overline{aa + xx^{\frac{1}{2}}}$.

Ex. 3.

Let $x \sqrt[4]{8} - \frac{5aa}{8x} \sqrt[4]{8} - \frac{75a^4}{512x^3} \sqrt[4]{8} - \&c.$
be given.

Reduced $x \sqrt[4]{8} \times : 1 - \frac{5aa}{8xx} - \frac{75a^4}{512x^4} \&c.$

Rule $1 + my + m \cdot \frac{m-1}{2} yy$, put-
ting $bx = y$.

Here $my = \frac{-5aa}{8xx}$, and $m \cdot \frac{m-1}{2} yy = \frac{-75a^4}{512x^4}$,

and

and by dividing $\frac{m-1}{2} y = \frac{15aa}{64xx}$; then $y = -\frac{5aa}{8xxx} = \frac{15aa}{32xx \cdot m-1}$, and $-\frac{1}{m} = \frac{3}{4 \times m-1}$, or $-4m+4=3m$, and $7m=4$, whence $m = \frac{4}{7}$ the index. Also $y = \frac{-5aa}{8xxx} = -\frac{5aa}{\frac{32}{7}xx} = -\frac{35aa}{32xx}$

the second term. And the binomial surd is

$$x \sqrt[4]{8 \times 1 - \frac{35aa}{32xx}}$$

Ex. 4.

Let the series

$$a^{\frac{1}{2}} - \frac{y}{4a^{\frac{1}{2}}} + \frac{5y^2}{96a^{\frac{3}{2}}} + \frac{5y^3}{384a^{\frac{5}{2}}} + \frac{35y^4}{18432a^{\frac{7}{2}}} \text{ \&c. be}$$

proposed.

This example resolved like the foregoing, gives $m = -\frac{3}{2}$, and $\frac{y}{6a}$ for the second term of the binomial. But $a^{\frac{1}{2}} \times 1 + \frac{y}{6a}$ does not produce the given series. Whence we may conclude it has not a binomial root.

For a trinomial root; for brevity's sake put $1, z, v$ for a, bx, cxx in the Rule, Prob. lix.

which rule then becomes $1+z+v =$

$$1 + mz + m \cdot \frac{m-1}{2} zz + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} z^3$$

$$+ mv + m \cdot \frac{m-1}{2} \cdot 2zv$$

and

and $a^{\frac{1}{2}} \times : 1 - \frac{y}{4a} + \frac{5y^2}{96aa} + \frac{5y^3}{384a^3}$ is the given series reduced. Then we have these three equations, $mz = \frac{-y}{4a}$, $m \cdot \frac{m-1}{2} zz + mv = \frac{5yy}{96aa}$,

and $m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} z^3 + m \cdot \frac{m-1}{2} \cdot 2zv = \frac{5y^3}{384a^3}$.

Divide the third by the first, and there comes out $\frac{m-1}{2} \cdot \frac{m-2}{3} z z + \overline{m-1} \cdot v = \frac{-5yy}{96aa}$; add this to

the second, and we have $m \cdot \frac{m-1}{2} z z + \frac{m-1}{2} \cdot \frac{m-2}{3} z z + \overline{2m-1} \cdot v = 0$, or $\frac{m-1}{2} \times \frac{4m-2}{3}$

$z z + \overline{2m-1} \cdot v = 0$. And squaring the first, $mmzz = \frac{yy}{16aa}$, and $\frac{yy}{aa} = 16mmzz$. Also

$mv = \frac{5yy}{96aa} - m \cdot \frac{m-1}{2} z z = \frac{5}{96} \times 16mmzz -$

$m \cdot \frac{m-1}{2} z z$. And $v = \frac{5}{6} mzz - \frac{m-1}{2} z z =$

$\frac{2m+3}{6} z z$. Therefore $\frac{m-1}{2} \cdot \frac{4m-2}{3} z z +$

$\overline{2m-1} \cdot v = \frac{m-1}{3} \cdot \overline{2m-1} \cdot z z + \overline{2m-1} \cdot$

$\frac{2m+3}{6} z z = 0$; or $\frac{2m-2+2m+3}{6} \times \overline{2m-1} = 0$,

that is $\frac{4m+1}{6} \times \overline{2m-1} = 0$, and $\overline{4m+1} \times$

$\overline{2m-1} = 0$. Which equation has two roots, $m = -\frac{1}{4}$, and $m = \frac{1}{2}$. If $m = -\frac{1}{4}$, then

$z = \frac{-y}{4am} = \frac{-v}{-a} = \frac{y}{a}$, and $v = \frac{2m+3}{6} z z$

$= 3^{-\frac{1}{2}} \times \frac{yy}{aa} = \frac{5yy}{12aa}$. And the surd root is $a^{\frac{1}{2}} \times$
 $: 1 + \frac{y}{a} + \frac{5yy}{12aa}$ $^{-\frac{1}{2}}$, which involved produces
 four terms of the series, but not the last.

And if $m = \frac{1}{2}$. Then $z = -\frac{7}{4am} = \frac{-y}{2a}$,
 and $v = \frac{2m+3}{0} zz = \frac{4}{6} \times \frac{yy}{4aa} = \frac{yy}{6aa}$. And
 then the surd is $a^{\frac{1}{2}} \times : 1 - \frac{y}{2a} + \frac{yy}{6aa}$ $^{\frac{1}{2}}$, which
 involved, produces all the terms of the given se-
 ries; and therefore is the root required.

P R O B L E M LXII.

To revert an infinite series; or to find the root of such a series.

R U L E.

If the series consists of all the powers of z , as $Az + Bz^2 + Cz^3 + Dz^4 + Ez^5$ &c. $= y$; then substitute the values of the coefficients, A, B, C, D, E , &c. into the following form, for the root.

$$\begin{aligned} z &= \frac{1}{A}y - \frac{B}{A^2}y^2 + \frac{2BB-AC}{A^3}y^3 \\ &+ \frac{5ABC-A^2D-5B^2}{A^4}y^4 \\ &+ \frac{14B^2-21AB^2C+6A^2BD+3A^2C^2-A^2E}{A^5}y^5 \\ &+ \frac{-42B^3+84AB^2C-28A^2B^2D-28A^2BC^2}{A^6}y^6 \\ &+ \frac{7A^2BE+7A^2DC-A^2F}{A^7}y^7, \text{ \&c.} \end{aligned}$$

For

For put $z = ay + by^2 + cy^3 + dy^4 \&c.$ Then

$$zz = aay^2 + 2aby^3 + bby^4 \&c.$$

$$z^3 = a^3y^3 + 3a^2by^4 \&c.$$

$$z^4 = a^4y^4 \&c.$$

$\&c.$ Whence

$$\begin{array}{r} Az = Aay + Aby^2 + Acy^3 + Ady^4 \&c. \\ + Bz^2 = Ba^2y^2 + 2Bab y^3 + Bbby^4 \&c. \\ + Cz^3 = Ca^3y^3 + 3Ca^2by^4 \&c. \\ + Dz^4 = Da^4y^4 \&c. \\ \&c. \end{array} \left. \vphantom{\begin{array}{r} Az \\ + Bz^2 \\ + Cz^3 \\ + Dz^4 \\ \&c. \end{array}} \right\} = y.$$

Then making the homologous powers equal, $Aay = y$,
and $a = \frac{1}{A}$. And $Ab + Ba^2 = 0$, or $b = \frac{-B}{A^2}$.

Likewise $Ac + 2Bab + Ca^3 = 0$, and $c = \frac{2BB - AC}{A^5}$.

In like manner $Ad + Bbb + 2Bac + 3Ca^2b + Da^4 = 0$,
whence $D = \frac{5ABC - A^2D - 5B^3}{A^7}$; and so on.

Ex. 1.

Suppose $x - xx + x^3 - x^4 + x^5 \&c. = y$, to find the
value of x in terms of y .

Here $z = x$, $A = 1$, $B = -1$, $C = 1$, $D = -1$, $\&c.$

Whence $x = \frac{y}{1} + \frac{1}{1}y^2 + \frac{2-1}{1}y^3 + \frac{-5+1+5}{1}y^4 \&c. = y + y^2 + y^3 + y^4 + y^5 \&c.$

Ex. 2.

Let $z = x + \frac{xx}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \frac{x^5}{5} \&c.$ is
find x in a series of z .

Here $z = x$, $y = z$, $A = 1$, $B = \frac{1}{2}$, $C = \frac{1}{3}$,
 $D =$

Sect. VI. INFINITE SERIES. 173

$$D = \frac{1}{4}, E = \frac{1}{5}, \&c. \text{ and } x = \frac{1}{1}z - \frac{1}{2}z^2$$

$$+ \frac{\frac{1}{2} - \frac{1}{3}}{1} z^3 + \frac{5 \times \frac{1}{6} - \frac{1}{4} - 5 \times \frac{1}{8}}{1} z^4 \&c.$$

$$= z - \frac{1}{2}z^2 + \frac{1}{6}z^3 - \frac{1}{24}z^4 + \frac{1}{120}z^5 \&c.$$

$$= z - \frac{z^2}{1.2} + \frac{z^3}{2.3} - \frac{z^4}{2.3.4} + \frac{z^5}{2.3.4.5} \&c.$$

$$\text{that is, } x = z - \frac{z}{2}A - \frac{z}{3}B - \frac{z}{4}C - \frac{z}{5}D \&c.$$

where A, B, C, &c. are the foregoing terms, with their signs.

Ex. 3.

$$\text{Suppose } r - \frac{aa}{2r} + \frac{a^4}{24r^3} - \frac{a^6}{720r^5} + \frac{a^8}{40320r^7} - \&c. = c, \text{ to find } a.$$

$$\text{Put } r - c = v. \text{ Then } \frac{aa}{2r} - \frac{a^4}{24r^3} + \frac{a^6}{720r^5} - \frac{a^8}{40320r^7} \&c. = v. \text{ Here } x = aa, y = v,$$

$$A = \frac{1}{2r}, B = \frac{-1}{24r^3}, C = \frac{1}{720r^5}, D = \frac{-1}{40320r^7} \&c. \text{ Whence}$$

$$aa = 2rv - \frac{1}{24r^3}vv + \frac{1}{288r^5} - \frac{1}{1440r^7}v^3 \&c.$$

$$= 2rv + \frac{1}{3}vv + \frac{4}{45r}v^3 + \frac{1}{35r^2}v^4 \&c. \text{ And}$$

$$\text{extracting the root, } a = \sqrt{2rv} \times \left(1 + \frac{v}{12r} + \right.$$

$$\left. \frac{v^2}{100r^2} + \frac{5v^3}{896r^3} + \&c. \right)$$

2 R U L E.

If the series consists of the odd powers of z , as
 $Az + Bz^3 + Cz^5 + Dz^7 \&c. = y$. Substitute the va-
 lues of the coefficients $A, B, C, \&c.$ into the fol-
 lowing form; which will give the root.

$$z = \frac{1}{A} y - \frac{B}{A^4} y^3 + \frac{3BB-AC}{A^7} y^5 +$$

$$\frac{8ABC - A^2D - 12B^3}{A^{10}} y^7$$

$$+ \frac{55B^4 - 55AB^2C + 10A^2BD + 5A^2C^2 - A^3E}{A^{13}} y^9$$

&c.

For put $z = ay + by^3 + cy^5 + dy^7 \&c.$

$$\text{Then } z^3 = a^3y^3 + 3a^2by^5 + 3a^2cy^7$$

$$+ 3abb$$

$$z^5 = a^5y^5 + 5a^4by^7$$

$$z^7 = a^7y^7$$

&c.

$$\text{And } Az = Aay + Aby^3 + Acy^5 + Ady^7 \&c. \left. \vphantom{Az} \right\} = y$$

$$+ Bz^3 = +Ba^3y^3 + 3Ba^2by^5 + 3Ba^2cy^7$$

$$+ 3Babb$$

$$+ Cz^5 = + Ca^5y^5 + 5Ca^4by^7$$

$$+ Dz^7 = + Da^7y^7 \left. \vphantom{Dz^7} \right\}$$

Then equating the coefficients of like terms;
 $Aa=1, Ab + Ba^3=0, Ac + 3Ba^2b + Ca^5=0,$
 $Ad + 3Ba^2c + 3Babb + 5Ca^4b + Da^7=0, \&c.$ whence

$$a = \frac{1}{A}, b = -\frac{Ba^3}{A} = -\frac{B}{A^4} \quad \text{Likewise}$$

$$c = \frac{3BB-AC}{A^7}, d = \frac{8ABC - A^2D - 12B^3}{A^{10}} \quad \&c.$$

Ex. 4.

Let $a = \frac{a^3}{2 \cdot 3 \cdot d d} + \frac{a^5}{2 \cdot 3 \cdot 4 \cdot 5 d^4} - \frac{a^7}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 d^6} + \&c.$
 $= y$; to find a .

Here $z = a$, $y = y$, $A = 1$, $B = -\frac{1}{2 \cdot 3 \cdot d d}$,
 $C = \frac{1}{2 \cdot 3 \cdot 4 \cdot 5 d^4}$, $D = -\frac{1}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 d^6} \&c.$

Whence $a = y + \frac{1}{2 \cdot 3 \cdot d d} y^3 + \frac{1}{3 \cdot 4 d^4} - \frac{1}{2 \cdot 3 \cdot 4 \cdot 5 d^4}$
 $\times y^5 + -\frac{1}{2 \cdot 3 \cdot 3 \cdot 5 d^4} + \frac{1}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 d^6} + \frac{1}{2 \cdot 3 \cdot 3 d^6} \times y^7$
 $\&c. = y + \frac{1}{2 \cdot 3 \cdot d d} y^3 + \frac{3}{2 \cdot 4 \cdot 5 d^4} y^5 + \frac{3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 7 d^6} y^7$
 $+ \&c.$

Ex. 5.

Suppose $y + \frac{y^3}{2 \cdot 3 \cdot d d} + \frac{3y^5}{2 \cdot 4 \cdot 5 d^4} + \frac{3 \cdot 5 y^7}{2 \cdot 4 \cdot 6 \cdot 7 d^6} +$
 $\&c. = a$, to find y .

Here $z = y$, $y = a$, $A = 1$, $B = \frac{1}{2 \cdot 3 \cdot d d}$, $C = \frac{3}{2 \cdot 4 \cdot 5 d^4}$,
 $D = \frac{3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 7 d^6} \&c.$ Then $y = a - \frac{1}{2 \cdot 3 \cdot d d} a^3 +$
 $\frac{1}{2 \cdot 2 \cdot 3 d^4} - \frac{3}{2 \cdot 4 \cdot 5 d^4} \times a^5 + \frac{1}{2 \cdot 5} - \frac{5}{2 \cdot 4 \cdot 2 \cdot 7} - \frac{1}{2 \cdot 3 \cdot 3} \times$
 $\frac{1}{d^6} \&c. = a - \frac{1}{2 \cdot 3 \cdot d d} a^3 + \frac{1}{2 \cdot 3 \cdot 4 \cdot 5 d^4} a^5 -$
 $\frac{1}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 d^6} a^7 + \&c.$

Ex.

Ex. 6.

Given $bx - \frac{bx^3}{6aa} - \frac{bx^5}{40a^4} - \frac{bx^7}{336a^6} - \frac{bx^9}{3456a^8}$
 &c. = d , to find z .

Dividing by b , $z - \frac{z^3}{6aa} - \frac{z^5}{40a^4} - \frac{z^7}{336a^6}$ &c.
 = $\frac{d}{b} = n$. Then $y = n$, $A = 1$, $B = -\frac{1}{6aa}$,

$C = -\frac{1}{40a^4}$, $D = -\frac{1}{336a^6}$, &c. then will

$$z = n + \frac{n^3}{6aa} + \frac{1}{12a^4} + \frac{1}{40a^4} \times n^5 +$$

$$\frac{8}{6.40} + \frac{1}{336} + \frac{12}{6.6.6} \times \frac{n^7}{a^6} + \text{\&c.} = n + \frac{n^3}{6aa}$$

$$+ \frac{13}{120a^4} n^5 + \frac{463}{840} n^7 + \text{\&c.}$$

3 R U L E.

When the series consists of any powers of z denoted by m and n , as $Az^m + Bz^{m+n} + Cz^{m+2n} + Dz^{m+3n} + Ez^{m+4n}$ &c. = y . Then substitute the values of the coefficients, A , B , C , &c. into this form, for the root or value of z .

Put $v = \frac{y}{A}$. Then

$$z = v^{\frac{1}{m}} - \frac{B}{mA} v^{\frac{1+n}{m}}$$

$$+ \frac{m+1+2n.BB-2mAC}{2mmAA} v^{\frac{1+2n}{m}}$$

$$\left. \begin{aligned} & - \frac{2mm + 9mn + 9nn + 3m + 6n + 1}{6m^3 A^3} B^3 \\ & + \frac{m + 3n + 1}{mm A^2} BC \\ & - \frac{D}{mA} \\ & \&c. \end{aligned} \right\} v^{\frac{1+3n}{m}}$$

For, put $z = v^{\frac{1}{m}} + bv^{\frac{1+n}{m}} + cv^{\frac{1+2n}{m}} + dv^{\frac{1+3n}{m}} + \&c.$

Then dividing the given series by A, we have

$$z^m + \frac{B}{A} z^{m+n} + \frac{C}{A} z^{m+2n} \&c. = \frac{y}{A} = v.$$

Whence by involution,

$$\left. \begin{aligned} z^m &= v + mbv^{\frac{m+n}{m}} + mcv^{\frac{m+2n}{m}} \&c. \\ &+ m \cdot \frac{m-1}{2} bb \\ \frac{B}{A} z^{m+n} &= \frac{B}{A} \times v^{\frac{m+n}{m}} + m+n \cdot bv^{\frac{m+2n}{m}} \&c \\ \frac{C}{A} z^{m+2n} &= \frac{C}{A} \times v^{\frac{m+2n}{m}} \&c. \\ \frac{D}{A} z^{m+3n} &= \&c. \end{aligned} \right\} = v.$$

Then equating the coefficients, $mb + \frac{B}{A} = 0$, and

$$b = \frac{-B}{mA}. \text{ And } mc + m \cdot \frac{m-1}{2} bb + m+n \cdot \frac{bB}{A} + \frac{C}{A}$$

$$= 0, \text{ and } c = \frac{m+1+2n \cdot BB - 2mAC}{2m^2 A^2} \&c.$$

Note, In all these rules, I have only pursued these series to a few terms; to have gone farther would have taken up too much room: but the method is visible.

Ex. 7.

Suppose $\frac{1}{2}x + \frac{1}{3}x^2 + \frac{1}{4}x^3 + \frac{1}{5}x^4$ &c. = y .

Here $z = x$, $v = 2y$, $A = \frac{1}{2}$, $B = \frac{1}{3}$, $C = \frac{1}{4}$, $D = \frac{1}{5}$, &c. and $m = 2$, $n = 1$. Whence,

$$x = v^{\frac{1}{2}} - \frac{1}{3}v^{\frac{3}{2}} + \frac{5BB - 4AC}{8AA}v^{\frac{5}{2}} \text{ \&c.} = v^{\frac{1}{2}} - \frac{1}{3}v + \frac{1}{2 \cdot 18}v^{\frac{3}{2}} + \frac{1}{270}v^2 \text{ \&c.}$$

Ex. 8.

Let $x = \frac{a^2}{2x} + \frac{a^4}{6x^3} - \frac{a^6}{24x^5} + \text{\&c.} = y = v$.

Here $z = x$, $m = 1$, $n = -2$, $A = 1$, $B = -\frac{aa}{2}$, $C = \frac{a^4}{6}$, $D = -\frac{a^6}{24}$ &c. and $x = y^{\frac{1}{2}} + \frac{aa}{2}y^{-\frac{3}{2}} + \frac{-2BB - 2AC}{2A^2}y^{-\frac{5}{2}}$, &c. = $y + \frac{aa}{2y} - \frac{5a^4}{12y^3} + \frac{5a^6}{8y^5}$ &c.

Ex. 9.

Let $x^{-\frac{1}{2}} - \frac{1}{2}x^{\frac{1}{2}} - \frac{1}{8}x^{\frac{3}{2}} - \frac{1}{16}x^{\frac{5}{2}} - \frac{5}{128}x^{\frac{7}{2}}$ &c. = z , to find x .

In this Ex. $z = x$, $v = z$, $m = -\frac{1}{2}$, $n = 1$, $A = 1$, $B = -\frac{1}{2}$, $C = -\frac{1}{8}$, $D = -\frac{1}{16}$, $E = -\frac{5}{128}$ &c.

Whence

Sect. VI. INFINITE SERIES. 179

Whence $x = z^{-2} - z^{-4} + \frac{2^1 BB + C}{\frac{1}{2}} z^{-6} +$
 $: \frac{14B^3 + 14BC + 2D}{\frac{1}{2}} : \times z^{-8} \&c.$ that is
 $x = \frac{1}{2z} - \frac{1}{2z^3} + \frac{1}{2z^5} - \frac{1}{2z^7} \&c.$

Cor. 1. If you would find any power of y ; find y in a series of z , and then involve that series to the power required, or else put $s = y$; then find s (y) from such a series as this,

$$As^{\frac{m}{r}} + Bs^{\frac{m+n}{r}} + Cs^{\frac{m+2n}{r}} \&c. = y,$$

by the last rule.

Cor. 2. The reverted series is of the same form as the given series; for otherwise they are not convertible into one another.

PROBLEM LXIII.

To extract the root of a series containing all the powers of two letters.

R U L E.

If the series consists of all the single powers of z and y , as $az + bz^2 + cz^3 + dz^4 \&c. = gy + by^2 + jy^3 + ky^4 \&c.$ substitute the values of the coefficients in the following form, for the root.

$$z = \frac{g}{a} y + \frac{b - bA^2}{a} y^2 + \frac{j - 2bAB - cA^3}{a} y^3$$

$$+ \frac{k - bB^2 - 2b^2AC - 3cA^2B - dA^4}{a} y^4$$

$$+ \frac{l - 2bBC - 2bAD - 3cAB^2 - 3cA^2C - 4dA^3B - eA^5}{a} y^5$$

$$+ \frac{m - 2bBD - bC^2 - 2bAE - cB^3 - 6cABC - 3cA^2D - 6dA^2B^2 - 4dA^3C - 5eA^4B - fA^6}{a} y^6 \&c.$$

Where A, B, C, &c. are the coefficients of the first, second, third, &c. terms.

Let $z = Ay + By^2 + Cy^3 + Dy^4$ &c. Then

$$\begin{aligned} az &= aAy + aBy^2 + aCy^3 + aDy^4 \text{ \&c.} \\ + bz^2 &= + bA^2y^2 + 2bABy^3 + bBBy^4 \\ &\quad + 2bAC \\ + cz^3 &= + cA^3y^3 + 3cA^2By^4 \\ + dz^4 &= + dA^4y^4 \\ \text{\&c.} & \end{aligned} \quad \left. \vphantom{\begin{aligned} az \\ + bz^2 \\ + cz^3 \\ + dz^4 \\ \text{\&c.} \end{aligned}} \right\} \\ = gy + by^2 + jy^3 + ky^4 \text{ \&c.}$$

And equating the coefficients, $aA = g$, and $A = \frac{g}{a}$.

Also $aB + bA^2 = b$, and $B = \frac{b - bAA}{a}$. Also

$aC + 2bAB + cA^3 = j$, and $C = \frac{j - 2bAB - cA^3}{a}$.

Again $aD + bB^2 + 2bAC + 3cA^2B + dA^4 = k$, and
 $D = \frac{k - bB^2 - 2bAC - 3cA^2B - dA^4}{a}$ &c.

Ex. I.

$$\text{Let } x = \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} \text{ \&c.} = \frac{1}{2}y + \frac{1}{3}y^2 + \frac{1}{4}y^3 + \frac{1}{5}y^4 \text{ \&c. to find } x.$$

$$\begin{aligned} \text{Here } z = x, y = y, a = 1, b = -\frac{1}{2}, c = \frac{1}{6}, \\ d = -\frac{1}{24} \text{ \&c. and } g = \frac{1}{2}, b = \frac{1}{3}, j = \frac{1}{4}, \\ k = \frac{1}{5} \text{ \&c.} \end{aligned}$$

Then

Then

$$x = \frac{1}{2}y + \frac{\frac{1}{3} + \frac{1}{8}}{1}y^2 + \frac{\frac{1}{4} + \frac{11}{48} - \frac{1}{48}}{1}y^3 \&c. \text{ or}$$

$$x = \frac{1}{2}y + \frac{11}{24}y^2 + \frac{11}{24}y^3 + \frac{1381}{2880}y^4 \&c.$$

Ex. 2.

Suppose $z + \frac{z^3}{6dd} + \frac{3z^5}{40d^4} + \frac{5z^7}{112d^6} \&c. = ny + \frac{ny^3}{6dd}$
 $+ \frac{3ny^5}{40d^4} + \frac{5ny^7}{112d^6} \&c. \text{ to find } z.$

Comparing this with the rule, and we have

$$a=1, b=0, c = \frac{1}{6dd}, d=0, e = \frac{3}{40d^4}, f=0, \&c.$$

$$g=n, h=0, j = \frac{n}{6dd}, k=0, l = \frac{3n}{40d^4}, m=0, \&c.$$

Whence

$$z = \frac{n}{1}y + 0y^2 + \frac{\frac{n}{6dd} - \frac{A^3}{6dd}}{1}y^3 + 0y^4 +$$

$$\frac{\frac{3n}{40d^4} - \frac{AAC}{2dd} - \frac{3A^5}{40d^4}}{1}y^5 \&c. \text{ where } B, D, \&c.$$

$$= 0; \text{ that is } z = ny + \frac{n-n^3}{6dd}y^3 +$$

$$\frac{\frac{3n}{40d^4} - \frac{n^3}{12d^4} + \frac{n^5}{120d^4}}{1}y^5 \&c. = ny + \frac{n-n^3}{6dd}y^3 +$$

$$\frac{9n-10n^3+n^5}{120d^4}y^5 = ny + \frac{n}{1} \times \frac{1-nn}{6dd}y^3 + \frac{n-n^3}{6dd}$$

$$\times \frac{9-nn}{20dd}y^5 \&c.$$

Or $z = ny + \frac{1-nn}{2.3dd}yyA + \frac{9-nn}{4.5dd}yyB \&c. \text{ where}$
 $A, B, \&c. \text{ are the foregoing terms.}$

2 R U L E.

In two series consisting of the powers and products of x and y ; as

$$az + bx^2 + cz^3 + dx^4 \&c. + fy + gzy + bx^2y + jz^3y \&c. \\ + ly^2 + my^2z + ny^2z^2 \&c. + py^3 + qy^3z \&c. + sy^4 \&c. \\ = 0.$$

Then substitute the values of the coefficients, into the following form;

$$z = -\frac{f}{a}y - \frac{l+gA+bA^2}{a}y^2 \\ - \frac{2bAB+cA^3+p+gB+mA+bA^2}{a}y^3 \\ - \frac{2bAC+bBB+3cA^2B+dA^4+s+gC+mB+qA}{a} \\ + \frac{2bAB+nA^2+jA^3}{a}y^4 \&c.$$

Where $A, B, C, \&c.$ are the coefficients of the first, second, third, $\&c.$ terms.

For put $z = Ay + By^2 + Cy^3 + Dy^4 \&c.$

Then

$$\left. \begin{aligned} az &= aAy + aBy^2 + aCy^3 \&c. \\ + lz^2 &= bA^2y^2 + 2bABy^3 \\ + cz^3 &= cA^3y^3 \\ + fy &= fy \\ + ly^2 &= ly^2 \\ + py^3 &= py^3 \\ + gyz &= gAy^2 + gBy^3 \\ + my^2z &= mAy^3 \\ + byz^2 &= bA^2y^3 \\ \&c. &= \end{aligned} \right\} = 0.$$

Then equating the coefficients, $aA+f=0$, $aB+bA^2+l+gA=0$, $\&c.$ whence $A = -\frac{f}{a}$, $B = -\frac{2bAB+cA^3+p+gB+mA}{a}$, $C = -\frac{2bAC+bBB+3cA^2B+dA^4+s+gC+mB+qA}{a} + \frac{2bAB+nA^2+jA^3}{a}$, $\&c.$

Ex. 3.

Suppose $2y + \frac{1}{12}y^3 + \frac{1}{2}x - \frac{1}{4}xx + xy - \frac{3}{8}xyy$
 $+ x^2y^2 = 0$; to find y .

Here $z = y$, $y = x$, $a = 2$, $c = \frac{1}{12}$, $f = \frac{1}{2}$,

$g = 1$, $b = -\frac{3}{8}$, $l = -\frac{1}{4}$, $n = 1$; $b, d, j, m, p,$
 $q, s = 0$. Therefore

$$y = -\frac{1}{4}x - \frac{-\frac{1}{2} + A}{2}x^2 - \frac{\frac{1}{2}A^2 + B - 3AA}{2}x^3 -$$

$$\frac{\frac{1}{2}A^3 + C - 3AB + A^2}{2}x^4 \&c. = -\frac{1}{4}x + \frac{1}{4}x^2 -$$

$$\frac{173}{768}x^3 + \frac{43}{384}x^4 \&c.$$

PROBLEM LXIV.

To extract the root of an adjoined equation, by a series.

R U L E.

If the equation consists of terms which contain the powers of x and y ; and you want the value of y , in a series of x . Make the equation $= 0$, and assume an indetermined series for the root, as $y = Ax^n + Bx^{n+r} + Cx^{n+s} + Dx^{n+t} \&c.$ wherein the indices $n+r, n+s, \&c.$ continually increase if x be very small; but they decrease if x be great; the first is an *ascending* series, and the latter a *descending* one. By this means the series will converge; every following term growing still less, till they vanish or become of no moment.

For y and its powers in the given equation, substitute the first term Ax^n and its powers. Then to determine n , put the two least indices equal to

each other, for an ascending series; or the two greatest, for a descending one. And if it appears not at first sight, which is the two least, or two greatest; it will be known, by comparing every two of the indexes.

Then to determine r , s , t , &c. substitute its value for n , in all these indices, and having taken the least for an ascending series, or the greatest for a descending one; subtract it from each of the rest. Then take these remainders, and add them to themselves and to one another, all possible ways; and these remainders, and the sums resulting, taken in order, will be the values of r , s , t , &c. which will be affirmative, in an ascending series; but negative in a descending one. Then put these values in the series, $Ax^n + Bx^{n+r} + Cx^{n+2r}$ &c.

Then to find the coefficients A , B , C , D , &c. substitute the last series for the powers of y , in the equation; and put the coefficient of each power of x , successively $= 0$; and A , B , C , &c. will be gradually found from these equations.

Ex. 1.

Let $a^4x^2 - a^4xy + x^6 = ay^5$, to find y .

By reduction $a^4x^2 - a^4xy + x^6 - ay^5 = 0$. Put $y = Ax^n + Bx^{n+r} + Cx^{n+s} + Dx^{n+t}$ &c. substitute Ax^n for y , in the equation, and we have $a^4x^2 - a^4Ax^{n+1} + x^6 - aA^5x^{5n} = 0$. Then equating the indices, $n+1=2$, for the least, or $5n=6$ for the greatest indices.

For an ascending series.

Here $n+1=2$, and $n=1$. Then the indices 2, $n+1$, 6, $5n$, become 2, 2, 6, 5. Subtract 2 from

from the rest, and you have 3, 4; out of which is composed this series 3, 4, 6, 7, 8, 9, 10, &c. for the values of $r, s, t, \&c.$ whence the form of the series will be $y = Ax + Bx^4 + Cx^6 + Dx^7 + Ex^8 \&c.$ This series substituted for y in the given equation, will be as follows:

$$\begin{array}{r}
 a^4x^2 = a^4x^2 \\
 -a^4xy = -a^4Ax^2 - a^4Bx^5 - a^4Cx^6 - a^4Dx^7 \&c. \\
 +x^6 = * \quad * \quad + \quad x^6 \\
 -ay^5 = * - aA^5x^5 \quad * - 5aA^4Bx^8 \&c. \\
 \hline
 0 = 0 \quad 0 \quad 0 \quad 0 \quad 0
 \end{array}$$

Then equating the homologous terms $a^4 - a^4A = 0$, and $A = 1$. Also $-a^4B - aA^5 = 0$, and $B = \frac{-A^5}{a^4} = -\frac{1}{a^4}$. Again, $-a^4C + 1 = 0$, and $C = \frac{1}{a^4}$. Likewise $-a^4D - 5aA^4B = 0$. Whence $D = +\frac{5}{a^6}$, &c. Then the series or root required is

$$y = x - \frac{x^4}{a^4} + \frac{x^5}{a^4} + \frac{5x^7}{a^6} \&c.$$

For a descending series.

Here $5n = 6$, the greatest indices, and $n = 1\frac{1}{5}$, and substituting this value of n , the indices 2, $n+1$, 6, $5n$ become 2, $2\frac{1}{5}$, 6, 6, and the remainders $-3\frac{1}{5}$, -4 ; and $r, s, t, \&c.$ will be $-3\frac{1}{5}$, -4 , $-7\frac{1}{5}$, $-7\frac{1}{5}$, -8 , &c. and the series becomes $y = Ax^{1\frac{1}{5}} + Bx^{-2\frac{1}{5}} + Cx^{-2\frac{1}{5}} + Dx^{-6\frac{1}{5}} \&c.$ which substituted in the given equation, will be

$$a^4x^2$$

$$\begin{aligned}
 a^4x^2 &= * & * & + a^4x^2 \\
 -a^4xy &= * & -a^4Ax^{2\frac{1}{2}} & * & -a^4Bx^{-1\frac{1}{2}} \\
 +x^6 &= +x^6 & & & \&c. \\
 -ay^5 &= -aA^5x^6 - 5aA^4Bx^{2\frac{1}{2}} - 5aA^4Cx^2 - 5aA^4Dx^{-1\frac{1}{2}} \\
 & & & & -10aA^3B^2 \\
 &= 0. & & & \&c.
 \end{aligned}$$

Then equating the coefficients of like terms,
 $1 - aA^5 = 0$, and $A = \frac{1}{a^{\frac{1}{5}}}$. Likewise $-a^4A - 5aA^4B = 0$, and $B = -\frac{1}{5}a^{\frac{3}{5}}$, also $a^4 - 5aA^4C = 0$, and $C = \frac{1}{5}a^{\frac{3}{5}}$. Also $-a^4B - 5aA^4D - 10aA^3B^2 = 0$, and $D = -\frac{1}{25}a^{\frac{7}{5}}$ &c. Whence the root is

$$y = \frac{x^{1\frac{1}{5}}}{a^{\frac{1}{5}}} - \frac{a^{\frac{3}{5}}}{5x^{2\frac{1}{5}}} + \frac{a^{\frac{3}{5}}}{5x^{2\frac{1}{5}}} - \frac{a^{\frac{7}{5}}}{25x^{6\frac{1}{5}}} \&c.$$

If you put $n+1=6$, the indices will be, 2, 6, 6, 25; but 6 is neither the greatest nor the least, therefore this succeeds not.

If you put $5n=2$, the indices will be 2, $1\frac{1}{5}$, 6, 2; but here also 2 is neither the greatest nor the least. Therefore this will not succeed.

If we put $n+1=5n$, the indices will be 2, $1\frac{1}{4}$, 6, $1\frac{1}{4}$; and $1\frac{1}{4}$ being the least, this will do for an ascending series; and the form of it will be
 $y = Ax^{\frac{1}{4}} + Bx + Cx^{1\frac{1}{4}} + Dx^{2\frac{1}{4}} \&c.$

Ex. 2.

Let $a^4x + ax^3 - a^4y - y^4 = 0$, be proposed.

Putting Ax^n for y , the equation becomes
 $a^4x + ax^3 - a^4Ax^n - A^4x^{4n} = 0$. Then put $n = \frac{1}{4}$
for

And

$$y = a^{\frac{1}{2}} x^{\frac{3}{2}} + \frac{a^{\frac{2}{4}}}{4} x^{-1\frac{1}{4}} - \frac{a^{\frac{2\frac{1}{2}}{4}}}{4} x^{-1\frac{1}{2}} - \frac{3a^{\frac{4\frac{1}{2}}{4}}}{32} x^{-3\frac{1}{4}} \&c.$$

Ex. 3.

Suppose $y^3 + aay + axy - x^3 - 2a^3 = 0$, to find y .

Put Ax^n for y , and the equation becomes
 $A^3x^{3n} + aaAx^n + aAx^{n+1} - x^3 - 2a^3x^0 = 0$.

For an ascending series.

Put the least indices $n=0$, and the indices become 0, 0, 1, 3 0; and the differences 1, 3; and $r, s, t, \&c. = 1, 2, 3, 4, 5, \&c.$ and the series $y = A + Bx + Cx^2 + Dx^3 \&c.$ Then

$$\begin{aligned} y^3 &= A^3 + 3A^2Bx + 3AB^2x^2 + 3A^2Dx^3 \&c. \\ &\quad + 3A^2C + B^3 \\ + a^2y &= a^2A + aaBx + aaCx^2 + aaDx^3 \\ + axy &= \quad + aAx + aBx^2 + aCx^3 \\ - x^3 &= \quad \quad \quad - x^3 \\ - 2a^3 &= - 2a^3 \end{aligned}$$

Then equating the coefficients, $A^3 + aaA - 2a^3 = 0$, and extracting the root, $A = a$; also $3A^2B + aaB +$

$aA = 0$, and $B = -\frac{1}{4}$. In like manner $C = \frac{1}{64a}$,

and $D = \frac{131}{512aa}$, &c. Whence $y = a - \frac{x}{4} +$

$$\frac{xx}{64a} + \frac{131x^3}{512aa} \&c.$$

Ex.

Ex. 4.

Let $y^3 + y^2 + y - x^3 = 0$, to find y in a descending series.

Putting Ax^n for y , the equation becomes $A^3x^{3n} + A^2y^{2n} + Ay^n - x^3 = 0$. Put $3n=3$, for the greatest indices. Then $n=1$, and all the indexes are 3, 2, 1, 3; and the differences $-1, -2$; and the series $-1, -2, -3, -4, \&c.$ and $y = Ax + B + Cx^{-1} + Dx^{-2} \&c.$ Then

$$\begin{array}{l|l}
 y^3 & = A^3x^3 + 3A^2Bx^2 + 3A^2Cx + 3A^2D \\
 & \qquad \qquad \qquad + 3ABB + 6ABC \} \&c. \\
 + y^2 & \qquad \qquad \qquad + A^2x^2 + 2ABx + BB \} \&c. \\
 & \qquad \qquad \qquad \qquad \qquad \qquad + 2AC \} \\
 + y & \dots\dots\dots + Ax \dots + B \quad \&c. \\
 - x^3 & - x^3
 \end{array}$$

Then equating the coefficients, $A^3=1$, and $A=1$.

Likewise $B = -\frac{1}{3}$, $C = -\frac{2}{9}$, $D = \frac{7}{81} \&c.$ and

therefore $y = x - \frac{1}{3} - \frac{2}{9x} + \frac{7}{81xx} \&c.$

2 R U L E.

Assume $y = Ax^n + Bx^{n+r} + Cx^{n+2r} + Dx^{n+3r}$ &c. and having found n , and put its value into the indices, as in Rule 1; set them down in order, and subtract each of them from the next greater; and you will have a series of differences. Then find the greatest number, which will measure all these differences; and this is the value of r , which must be affirmative in an ascending series, or when x is small; and negative, in a descending one,

one, when x is great. Then the values of n and r must be substituted in the assumed series.

The process must then go on as in Rule 1; and if there be any superfluous terms, which will be known by some of the coefficients A, B, C coming out $=0$; these terms must be thrown out of the series, and the operation begun anew.

Ex. 5.

Let $y^3 - axy + x^3 = 0$, be given.

Put $Ax^n +$ for y , and the equation becomes $A^3x^{3n} - aAx^{n+1} + x^3 = 0$. Let $n+1=3$, and $n=2$; and the indices are 6, 3, 3; that is, 3, 6. Then $6-3=3$, then $r=3$; and the least indices being compared, the series will be an ascending one, which is this $y = Ax^2 + Bx^5 + Cx^8 + Dx^{11}$ &c. which substituted in the given equation will be as follows:

$$\begin{array}{l} y^3 \\ -axy \\ +x^3 \end{array} \left| \begin{array}{l} +A^3x^6 + 3A^2Bx^9 + 3A^2Cx^{12} \text{ \&c.} \\ -aAx^3 - aBx^6 - aCx^9 - aDx^{12} \\ +x^3 \end{array} \right.$$

Then $aA=1$, and $A = \frac{1}{a}$, $B = \frac{1}{a^4}$, $C = \frac{3}{a^7}$,
 $D = \frac{12}{a^{10}}$ &c. Whence $y = \frac{x^2}{a} + \frac{x^5}{a^4} + \frac{3x^8}{a^7}$
 $+ \frac{12x^{11}}{a^{10}}$ &c.

Ex. 6.

Let $y^5 - by^2 + 9bx^2 - x^3 = 0$.

Substitute Ax^n for y , and the equation is $A^5x^{5n} - bA^2x^{2n} + 9bx^2 - x^3 = 0$. Put $2n=2$; whence

Sect. VI. INFINITE SERIES. 191

whence $n=1$, and the indices are 5, 2, 2, 3; and the differences 1, 2. Whence $r=1$. Therefore $y = Ax + Bx^2 + Cx^3 + Dx^4$ &c. Then

$$\begin{array}{l} -by^5 \\ +9bx^5 \\ -x^5 \end{array} \left| \begin{array}{l} -bA^2x^2 - 2bABx^3 - 2bACx^4 - 2bADx^5 \\ +9bx^2 \\ -x^3 \end{array} \right. \begin{array}{l} +A^5x^5 \\ -bBB \\ -2bBC \end{array}$$

Here $bA^2 = 9b$; and $A = 3$: also $B = -\frac{1}{6b}$

$$C = -\frac{1}{216bb}, D = \frac{81}{2b} - \frac{1}{3888b^3}. \text{ Whence}$$

$$y = 3x - \frac{x^2}{6b} - \frac{x^3}{216bb} + \frac{81}{2b} - \frac{1}{3888b^3} \times x^4$$

&c.

3 R U L E.

If the equation determining A, be an adaffected equation, which has several equal roots or values of A, then you must divide the least remainder, found by Rule 1, by the number of equal roots, one of which you take for A; and take this quotient for another remainder. Or else divide r found by Rule 2, by that number, and make use of the quotient, instead of r .

Ex. 7.

Let $y^9 - xy^3 + 2x^2y^2 - x^3y - x^4 = 0$, to find y :

Put Ax^n for y , and the equation becomes

$$A^9x^9 - A^3x^{3n+1} + 2A^2x^{2n+2} - Ax^{n+3} - x^4 = 0.$$

Let $3n+1 = 2n+2$; whence $n=1$, then the indices are 9, 4, 4, 4, 14. But the sum of the

coefficients for the least index 4, is $-A^3 + 2A^2 - A = 0$, or $A^3 - 2A + 1 = 0$, which equation has

two

two equal roots $A=1$, and $A=1$. Now the difference of the indexes will be 5, 10; therefore divide 5 by 2, gives $\frac{5}{2}$, and we have $\frac{5}{2}$, 5, 10 for the differences. Therefore $r, s, t, \&c.$ will be $\frac{5}{2}, 5, 7\frac{1}{2}, 10, \&c.$ Or (Rule 2) $r=5$; therefore $\frac{r}{2} = \frac{5}{2}$ to be taken for r ; whence the series will be

$$y = Ax + Bx^{3\frac{1}{2}} + Cx^6 + Dx^{8\frac{1}{2}} \&c. \text{ Then}$$

y^9		$+ A^9 x^9$
$-xy^3$	$-A^3 x^4 - 3A^2 Bx^{6\frac{1}{2}}$	$-3A^2 Cx^9 - 3AB^2$
$+2x^2 y^2$	$+2A^2 x^4 + 4ABx^{6\frac{1}{2}}$	$+4ACx^9 + 2BB$
$-x^3 y$	$-Ax^4 - Bx^{6\frac{1}{2}}$	$-Cx^9$
$-x^{14}$		$- \&c.$

Hence $-A^3 + 2A^2 - A = 0$, and $A=1$; $4B - 4B = 0$, and B may be taken at pleasure. Suppose $B=-1$, then $1 - 3C - 3 + 4C + 2 - C = 0$, or $4C = 4C$, and C may be taken at pleasure. Let $C=1$; then $y = x - x^{3\frac{1}{2}} - x^6 \&c.$

Or thus; In the second equation, $4B = 4B$; which concludes nothing; also $1 - 3C - 3BB + 4C + 2BB - C = 0$; that is, $1 - BB = 0$, and $B=1$ or $-1, \&c.$

Ex. 8.

$$\text{Let } A^4 y^2 - 2a^4 xy + a^4 x^2 + x^4 y^2 = 0.$$

Put Ax^n for y , and the indices become $2n, n+1, 2, 2n+4$. Let $2n=2$, or $n=1$, and the indices

tion; then find the root in an ascending series of the new letter. Or if the quantity (x) be very great, and the series for y is to ascend by x^2 's. Take some quantity nearly equal to x , and substitute the sum of that and a new letter for x .

Ex. 9.

Let $y^3 + aay - x^3 = 0$, where $x = \frac{2}{3}a$, nearly.

Put $\frac{2}{3}a - v = x$. Then $x^3 = \frac{8}{27}a^3 - \frac{4}{3}aav + 2av^2 - v^3$, then $y^3 + aay - \frac{8}{27}a^3 + \frac{4}{3}aav - 2av^2 + v^3$

$= 0$. Let $y = Av^n$ then the indices are $3n, n, 0, 1, 2, 3$. Let $n = 0$, then the indices become $0, 0, 0, 1, 2, 3$; and $y = A + Bv + Cv^2 + Dv^3$ &c. Then

$$\begin{array}{r|l}
 y^3 & A^3 + 3A^2Bv + 3A^2Cv^2 \text{ \&c.} \\
 + aay & + 3ABB \\
 - \frac{8}{27}a^3 & aaA + aaBv + aaCv^2 \\
 + \frac{4}{3}a^2v & - \frac{8}{27}a^3 \\
 - 2av^2 & \phantom{- \frac{8}{27}a^3} + \frac{4}{3}a^2v \\
 + v^3 & \phantom{- \frac{8}{27}a^3} - 2av^2 \\
 & \phantom{- \frac{8}{27}a^3} + \text{\&c.}
 \end{array}$$

Then $A^3 + 2\frac{1}{3}aaA - \frac{8}{27}a^3 = 0$, let $A = r$. Also

$B = -\frac{4}{3}$; and $3A^2C + 3ABB + aaC = 2a$. Whence

$$C = \frac{2a - 5\frac{1}{3}r}{3rr + aa}, \text{ \&c.}$$

Ex.

Ex. 10.

Let $y^4 - x^2y^2 + xy^2 + 2y^2 - 2y + 1 = 0$, where $x = 2$ very near.

Let $x = 2 + z$, which substituted for x , there arises $y^4 - z^2y^2 - 3zy^2 - 2y + 1 = 0$.

Let $y = Az^n$, then the equation is $A^4z^{4n} - A^2z^{2n+2} - 3A^2z^{2n+1} - 2Az^n + 1 = 0$. Let $n = 0$, and the indices are 0, 2, 1, 0, 0; and the differences 1, 2, 3, 4, &c. whence $y = A + Bz + Cz^2 + Dz^3$ &c. Then

y^4	A^4	$+ 4A^3Bz$	$+ 4A^2Cz^2$	$\&c.$
$- z^2y^2$			$+ 6A^2BB$	
$- 3zy^2$		$- 3A^2z$	$- 6ABz^2$	
$- 2y$	$- 2A$	$- 2Bz$	$- 2Cz^2$	
$+ 1$	$+ 1$			

Here $A^4 - 2A + 1 = 0$, and $A = 1$; also $4B - 2B = 3$, and $B = \frac{3}{2}$; and $4C + 6BB - 1 - 6B - 2C = 0$, and $C = -\frac{7}{4}$, &c. and $y = 1 + \frac{3}{2}z - \frac{7}{4}z^2 + z^3$ &c.

Cor. 1. In all these cases of extracting roots, the series must be made to converge, or else they are of no use. For in a converging series, the terms grow continually less and less, and so approach nearer and nearer to the true root, till the difference is as small as you will. But a diverging series always runs farther from the root, and therefore gives a false value thereof.

Cor. 2. If y be denoted by a series of x ascending; the lesser x is, the faster the series converges. And

And in a series of x descending; the greater x is, the faster likewise it converges. Therefore we are so to contrive the series, that we may have the least quantity in the numerators, or the greatest in the denominators.

Cor. 3. If the equation for finding the first term A , be an adsefected equation; as many roots or different values of A , as that equation has, so many different series will arise. For the first term A being different in each, the coefficients B, C, D , depending thereon, will also be different. Likewise, if two roots are equal, the second term will vanish, and the coefficient B will be found in the third, which will be a quadratic equation. And if there be three equal values of A , the second and third equations vanish, and the fourth contains a cubic equation of B , &c.

Cor. 4. An equation will also admit of several different series for the roots, according to the different values assumed for n . Also there are other equations that are impossible, and will admit of no roots.

Cor. 5. When the first equation, or that for determining A , has several equal roots; then the values of r, s, t , &c. must be divided by that number. Or, which is the same thing, the indices of x (r, s, t) found by Rule 1, must have others interposed between them, according to the number of equal roots. As for two equal roots, the series $Ax^n + Bx^{n+r} + Cx^{n+s}$ &c. must be reduced to this, $Ax^n + Bx^{n+\frac{r}{2}} + Cx^{n+r} + Dx^{n+\frac{r}{2}} + Ex^{n+s}$ &c. If this be not done, the second term B will be infinite, and all the following ones.

Cor. 6. If the series $A + \overline{B+C} \times z + \overline{D+E+F} \times z^2 + \overline{G+H+I+K} \times z^3$ &c. $= 0$, z being an indetermined quantity; then whatever value is put upon z , it will be $A=0, B+C=0, D+E+F=0, G+H+I+K=0$, &c. For

Sect. VI. INFINITE SERIES. 197

For this being a general equation, where z may be of any value; therefore put $z=0$, and then will $A=0$, and $B+C \times z + D+E+F \times z \&c. = 0$, divide by z , then $B+C + D+E+F \times z \&c. = 0$. Again, put $z=0$, and then $B+C=0$; whence $D+E+F \times z + G+H+I+K \times z^2 \&c. = 0$. Divide again by z , and $D+E+F + G+H+I+K \times z = 0$. Again, put $z=0$, then $D+E+F=0$, and $G+H+I+K \times z = 0$, and $G+H+I+K = 0$, &c.

Reversion of series, and the extracting the roots of all infinite series, depends upon this. For the coefficients of the several powers of the indetermined quantity, must be put $=0$, or else the whole equation cannot vanish, as it ought to do. And this being done, the several assumed coefficients A, B, C are determined as in the problems above.

SCHOLIUM.

To find y in the series $ay^\mu + by^{\mu+\nu} + cy^{\mu+2\nu} \&c. = fx^\pi + gx^{\pi+\rho} + bx^{\pi+2\rho} \&c.$ Assume $y = Ax^n + Bx^{n+r} + Cx^{n+s} \&c.$ Then by substitution we get, $aA^\mu x^{\mu n} + bA^{\mu+n} x^{\mu n + \nu n} \&c. = fx^\pi + gx^{\pi+\rho} \&c.$ Whence making the least indices equal, $\mu n = \pi$; then $n = \frac{\pi}{\mu}$, and the differences will be $\frac{\pi\nu}{\mu}, \rho$; &c. Then find q the greatest common divisor of $\frac{\pi\nu}{\mu}$ and ρ ; and the form of the series will be

$$y = Ax^{\frac{\pi}{\mu}} + Bx^{\frac{\pi}{\mu} + q} + Cx^{\frac{\pi}{\mu} + 2q} + Dx^{\frac{\pi}{\mu} + 3q} \&c.$$

in which the coefficients will be determined as before.

S E C T. VII.

*Some general and fundamental Problems, useful
and necessary in algebraical calculations.*

P R O B L E M LXV.

*The sum and difference of two quantities being given;
to find the quantities.*

L E T s = the sum
 d = the difference
 a = greater quantity
 e = the lesser.

then $a + e = s$, by the problem.
 and $a - e = d$.

then $2a = s + d$ by addition
 and $2e = s - d$ by subtracting.

Whence $a = \frac{s+d}{2}$, and $e = \frac{s-d}{2}$

or $a = \frac{1}{2}s + \frac{1}{2}d$, and $e = \frac{1}{2}s - \frac{1}{2}d$.

Cor. 1. *Half the sum added to half the difference
of two quantities, is equal to the greater.*

Cor. 2. *Half the difference of two quantities, taken
from half the sum, gives the lesser quantity.*

P R O B L E M LXVI.

*To find out the least common dividend, or the least
quantity, that can be divided by several given
quantities.*

R U L E.

Resolve each of the quantities into all the simple
divisors contained therein, by first dividing by
the

Sect. VII. FUNDAMENTAL PROBLEMS. 199

the least, and then by the next, and so on, till they are all exhausted; and collect these divisors together for each quantity. Then if there be any divisors in the second quantity which is not in the first, multiply the first by such divisors. Likewise, if there be any divisors in the third quantity which is not in this last, multiply it thereby, or put them into that quantity. Likewise such divisors as are in the fourth quantity and are not in this last, must be put into it, and so on. And lastly, all these divisors, in this last quantity must be multiplied together for the least common dividend.

Or shorter thus,

Divide the product of any two of the quantities by their greatest common divisor, (found by Prob. x. Sect. II.) take this quotient and a third quantity, and divide their product, by their greatest common divisor. Take this quotient and another quantity, and proceed as before; and so on to the last quantity. And the last quotient will be the least common dividend.

Ex. I.

What is the greatest common dividend of a^2bc , and $2ab^2d$.

The divisors of a^2bc are a, a, b, c
of $2abbd$ are $2, a, b, b, d$,

Here $2, b, d$ are in the last but not in the first; therefore $a \times a \times b \times c \times 2bd$, or $2aabbcd$ is the least common dividend.

Or thus,

The greatest common measure is ab , then the product is $2a^2b^2cd$. $ab)2a^2b^2cd(2aabbcd$ the least dividend.

Ex. 2.

Let $ab+cd$ and $ac+bd$ be proposed.

These have no divisors but 1. Therefore $\overline{ab+cd} \times \overline{ac+bd}$ or $aabc+abbd+accd+bcdd$, is the dividend required.

Ex. 3.

Let $3a^2b$, a^3+a^2b , and $aa-bb$, be given.

The greatest common divisor of $3a^2b$ and a^3+a^2b is aa . Then

$$aa) 3a^2b + 3a^2bb \ (3a^2b + 3aabb)$$

Then the greatest common divisor of $3a^2b + 3aabb$ and $aa-bb$, is $a+b$; then $\overline{3a^2b + 3aabb} \times \overline{aa-bb}$ divided by $a+b$ is

$(a+b) 3a^2b + 3a^2bb - 3a^2b^1 - 3a^2b^1 (3a^2b - 3a^2b^1)$
the least common dividend.

Ex. 4.

Let the given quantities be a^4-b^4 , $aa+ab$, $a^2+a^2b^2$, and $a+b$.

These quantities resolved into their divisors are $\overline{aa+bb} \times \overline{a+b} \times \overline{a-b}$, $\overline{a} \times \overline{a+b}$, $\overline{a} \times \overline{a} \times \overline{aa+bb}$, and $\overline{a+b}$. Now because there is one factor a in the second which is not in the first, put it in the first, which becomes $\overline{aa+bb} \times \overline{a+b} \times \overline{a-b} \times \overline{a}$, the least dividend for the first two quantities.

Likewise, there is a , one factor in the third, which is not in this last; let it be inserted, and it becomes $\overline{aa+bb} \times \overline{a+b} \times \overline{a-b} \times \overline{aa}$, the least dividend for three quantities.

Lastly, Since $a+b$ the last given quantity is in the last dividend; it will be the dividend for all four; that is, $\overline{aa+bb} \times \overline{a+b} \times \overline{a-b} \times \overline{aa}$, or a^6-b^6 .

$a^6 - a^4bb$ is the least common dividend for the four given quantities.

SCHOLIUM:

All the simple divisors of a quantity, are found the same way, as in Prob. 6, 7. Chap. iv. B. II. Arithmetic.

PROBLEM LXVII.

The sum and difference of two quantities being given; to find the difference of their squares.

Let s = sum, d = difference, A = greater quantity, E = the lesser. Then $A = \frac{s+d}{2}$, and $E = \frac{s-d}{2}$ (Prob. lxv). Whence

$$AA = \frac{ss + 2sd + dd}{4}$$

$$\text{and } EE = \frac{ss - 2sd + dd}{4}$$

$$\text{and } AA - EE = \frac{4sd}{4} = sd.$$

Cor. *The product of the sum and difference of two quantities, is equal to the difference of their squares.*

PROBLEM LXVIII.

Two quantities being given to find the square of the sum.

Let a be the greater quantity, e the lesser; then the sum is $a + e$; and $a + e$ being squared is $aa + 2ae + ee$.

Cor. 1. *Hence the square of the sum of two quantities is equal to the sum of the squares of the quantities, increased by double their product.*

Cor.

Cor. 2. *The square of the sum of any number of quantities, $a+b+c$ &c. is equal to the sum of all the squares, together with twice the sum of all the products of every two.*

For by this prob. $\overline{a+b+c}^2 = \overline{a+b}^2 + 2 \times \overline{a+b} \times c + cc$; that is $aa+2ab+bb+2ac+2bc+cc$, and so for more quantities.

Schol. By the same way, theorems may be found for the cube of the sum of two or more quantities.

PROBLEM LXIX.

Two quantities being given to find the square of their difference.

Let a be the greater, e the lesser; then the difference is $a-e$, which being squared, produces $aa-2ae+ee$.

Cor. Hence the square of the difference of two quantities, is equal to the sum of their squares abating twice their product.

Schol. By the same method a rule may be found for the cube of the difference of two quantities.

PROBLEM LXX.

The sum and difference of two quantities being given; to find their rectangle.

Let s =sum, d =difference, A the greater, E the lesser. Then $A+E=s$, and $A-E=d$; and adding these equations $2A=s+d$; and subtracting, $2E=s-d$. Then $2A \times 2E$ or $4AE = \overline{s+d} \times \overline{s-d} = ss-dd$, and $AE = \frac{ss-dd}{4}$.

Cor. *The square of the sum, less the square of the difference of two quantities, is equal to four times their rectangle.*

PRO-

PROBLEM LXXI.

Given the n^{th} power of the binomial $a+b$; to find the difference between the square of the sum of the odd terms, and the square of the sum of the even terms.

The n^{th} power of $a+b$, that is $\overline{a+b^n} = \overline{a^n + na^{n-1}b + n \cdot \frac{n-1}{2} a^{n-2}bb + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} a^{n-3}b^3$
 &c. Put A, B, C, D, E, &c. for the first, second, third, fourth, &c. terms. Then $A+C+E$ &c. = sum of the odd terms; and $B+D+F$ &c. = sum of the even terms. But $\overline{A+C+E}$ &c.

$$\overline{B+D+F}^2 = \overline{A+B+C+D+E}$$
 &c. \times

$$\overline{A-B+C-D+E}$$
 &c. =

$$\overline{a^n + na^{n-1}b + n \cdot \frac{n-1}{2} a^{n-2}bb + \&c.}$$
 \times

$$\overline{a^n - na^{n-1}b + n \cdot \frac{n-1}{2} a^{n-2}bb - \&c.}$$

$$= \overline{a+b^n} \times \overline{a-b^n} = \overline{aa-bb^n}.$$

Cor. 1. Hence $\overline{aa-bb^n} =$

$$\overline{a^n + \frac{n \cdot n-1}{2} a^{n-2}bb + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \frac{n-3}{4} \cdot a^{n-4}b^4 + \&c.}$$

$$\overline{-na^{n-1}b + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} a^{n-3}b^3 + n \cdot \frac{n-1}{2} \cdot}$$

$$\overline{\frac{n-2}{3} \cdot \frac{n-3}{4} \cdot \frac{n-4}{5} a^{n-5}b^5 + \&c.}$$

Cor. 2. $\overline{aa-bb^2} = \overline{aa+bb^2} - \overline{2ab^2}.$

Cor. 3. $\overline{aa-bb^3} = \overline{a^3+3abb^2} - \overline{3aab+b^3}.$

Cor.

$$\text{Cor. 4. } \overline{aa-bb}^4 = \overline{a^4 + 6aabb + b^4} - \overline{4a^3b + 4ab^3}^2.$$

$$\text{Cor. 5. } \overline{aa-bb}^5 = \overline{a^5 + 10a^3bb + 5ab^4} - \overline{5a^2b + 10a^2b^3 + b^5}^2 \&c.$$

PROBLEM LXXII.

To find the n^{th} root of the binomial surd $\overline{A+B}$, where either A or B is a surd square root, the other rational.

Suppose $x+v = \sqrt[n]{\overline{A+B}}$, then by involution $\overline{A+B} = x + nx^{n-1}v + n \cdot \frac{n-1}{2} x^{n-2}vv + n \cdot \frac{n-1}{2}$.

$\frac{n-2}{3} x^{n-3}v^3 \&c.$ Suppose v a surd square root, and put the odd terms of the series $=A$, and the

even ones $=B$; that is $x^n + n \cdot \frac{n-1}{2} x^{n-2}vv \&c.$

$=A$, and $nx^{n-2}v + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} x^{n-3}v^3 \&c.$

$=B.$ Then $x^n + n \cdot \frac{n-1}{2} x^{n-2}vv \&c.$

$= nx^{n-1}v + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} x^{n-3}v^3 \&c. = A^2 -$

$B^2 = D$ by substitution. That is (by Cor. 1. Prob.

lxxi.), $\overline{xx-vv}^n = D$, and $xx-vv = D^{\frac{1}{n}}$. There-

fore $vv = xx - D^{\frac{1}{n}}$. Whence this equation,

$x^n + n \cdot \frac{n-1}{2} x^{n-2} \times \overline{xx - D^{\frac{1}{n}}} + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}$

$\frac{n-3}{4} x^{n-4} \times \overline{xx - D^{\frac{1}{n}}}^2 + \&c. = A.$ Which ad-

fect equation, by a few trials, will give x , and then

then v will be had by the equation,

$$v = \pm \sqrt[n]{xx - \sqrt{D}}; \text{ and } x \pm v = \sqrt[n]{A \pm B}, \text{ as required.}$$

Cor. Hence if $x \pm \sqrt{y} = \sqrt[n]{A \pm B}$, and $A^2 - B^2 = D$; then $x^n + n \cdot \frac{n-1}{2} \times \frac{xx - D^{\frac{1}{n}}}{xx} P + \frac{n-2}{3} \cdot \frac{n-3}{4} \times \frac{xx - D^{\frac{1}{n}}}{xx} Q + \frac{n-4}{5} \cdot \frac{n-5}{6} \times \frac{xx - D^{\frac{1}{n}}}{xx} R \&c. = A$; and $\sqrt{y} = \sqrt[n]{xx - \sqrt{D}}$. Where P, Q, R, are the foregoing terms.

Or thus,

$$\text{Since } x^n + n \cdot \frac{n-1}{2} x^{n-2} v v \&c. = A.$$

$$\text{and } nx^{n-1}v + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} x^{n-3} v^3 \&c. = B.$$

by adding,

$$x^n + nx^{n-1}v + n \cdot \frac{n-1}{2} x^{n-2} v^2 \&c. = A + B.$$

by subtracting,

$$x^n - nx^{n-1}v + n \cdot \frac{n-1}{2} x^{n-2} v^2 - \&c. = A - B.$$

Therefore by extraction,

$$x + v = \sqrt[n]{A + B}.$$

$$\text{and } x - v = \sqrt[n]{A - B}.$$

$$\text{and } 2x = \sqrt[n]{A + B} + \sqrt[n]{A - B}.$$

$$2v = \sqrt[n]{A + B} - \sqrt[n]{A - B}.$$

whence

$$\text{whence } x = \frac{\sqrt[n]{A+B} + \sqrt[n]{A-B}}{2}$$

$$v = \frac{\sqrt[n]{A+B} - \sqrt[n]{A-B}}{2}$$

Therefore $x \pm v$ will be had, at least in decimals.

$$\text{Cor. Hence, } \sqrt[n]{A+B} = \frac{\sqrt[n]{A+B} + \sqrt[n]{A-B}}{2} \\ \pm \frac{\sqrt[n]{A+B} - \sqrt[n]{A-B}}{2}$$

Ex. 1.

Extract the square root of $11+6\sqrt{2}$.

Here $A=11$, $B=6\sqrt{2}$, $AA-BB=49=D$,
and $\sqrt{D}=7$. Therefore $x^2 + \frac{xx-7}{xx}x^2 = 11$,
or $2xx-7=11$, and $xx=9$, or $x=3$. Likewise
 $\sqrt{y}=+\sqrt{9-7}=+\sqrt{2}$, and $x+\sqrt{y}=3+\sqrt{2}$
the root.

Or thus,

$A+B=11+6\sqrt{2}=19.484$, and $A-B=2.516$.

Whence $x = \frac{\sqrt{19.484} + \sqrt{2.516}}{2} =$
 $\frac{4.414 + 1.586}{2} = 3$, and $v = \frac{4.414 - 1.586}{2} =$
 $1.414 = \sqrt{2}$, and $x+v=3+\sqrt{2}$, the root.

Ex. 2.

Suppose $37-20\sqrt{3}$ be given, to extract the square root.

Here $A=37$, $B=20\sqrt{3}$, $AA-BB=169=D$,
and $\sqrt{D}=+13$. Therefore $2xx-13=37$, and
 $xx=25$, or $x=5$.

Also

Also $\sqrt{y} = -\sqrt{25-13} = -\sqrt{12}$, and
 $x + \sqrt{y} = 5 - \sqrt{12}$. Or the root may be $\sqrt{12} - 5$,
 Putting $\sqrt{D} = -13$.

Or thus,

$$x = \frac{\sqrt{71.64} + \sqrt{2.36}}{2} = \frac{10.00}{2} = 5.$$

$$v = \frac{\sqrt{2.36} - \sqrt{71.64}}{2} = -\frac{6.92}{2} = -3.46$$

$= -2\sqrt{3}$, and $x + v = 5 - 2\sqrt{3}$ the root.

Ex. 3.

Let $7 - 5\sqrt{2}$ be given, to find the cube root.

Here $A = 7$, $B = 5\sqrt{2}$, $A^2 - B^2 = -1 = D$, and
 $D^{\frac{1}{2}} = -1$.

Then $x^3 + 3 \times \frac{xx+1}{xx} x^2 = 7$, or $4x^3 + 3x = 7$, and
 the root of this equation is $x = 1$.

Also $\sqrt{y} = -\sqrt{xx+1} = -\sqrt{2}$, and
 $x + \sqrt{y} = 1 - \sqrt{2}$.

Or thus,

$$\sqrt[3]{A+B} = \sqrt[3]{14.07} = 2.414, \text{ and}$$

$$\sqrt[3]{A-B} = \sqrt[3]{-.07} = -.412;$$

$$\text{Then } x = \frac{-.414 + 2.414}{2} = 1.$$

$$\text{And } v = \frac{-.414 - 2.414}{2} = -1.414 = -\sqrt{2}$$

for here B is negative; therefore $x + v = 1 - \sqrt{2}$.

Ex. 4.

What is the cube root of $25 + \sqrt{968}$.

Here $A = 25$, $B = \sqrt{968}$, $AA - BB = -343 = D$, and $\sqrt[3]{D} = -7$.

Then $x^3 + 3x^2 + 21x = 25$, and $x = 1$.

And $\sqrt{y} = \sqrt{8}$, and the root, $1 + \sqrt{8}$.

Ex.

Ex. 5:

Extract the cube root of $-10 + \sqrt{-243}$.

Here $A^2 - B^2 = 100 + 243 = 343 = D$, and $D^{\frac{1}{3}} = 7$. Therefore $x^3 + 3x \times \sqrt{-7} = -10$, or $4x^3 - 21x = -10$, and the root is $x = 2$, whence v or $\sqrt{y} = \sqrt{4 - 7} = \sqrt{-3}$, and $x + \sqrt{y} = 2 + \sqrt{-3}$, as required.

In like manner the cube root of $-10 - \sqrt{-243}$ is $2 - \sqrt{-3}$.

Ex. 6.

Extract the 5th root of $843 - 589\sqrt{2}$.

Here $AA - BB = 16807$, and $D^{\frac{1}{5}} = 7$.

And $16x^5 - 140x^3 + 245x = 843$, and the root is $x = 3$; and $\sqrt{y} = -\sqrt{9 - 7} = -\sqrt{2}$, and $x + \sqrt{y} = 3 - \sqrt{2}$ the root required.

Ex. 7.

What is the 7th root of $568 + 328\sqrt{3}$.

Here $A^2 - B^2 = -128 = D$, and $\sqrt[7]{D} = -2$. Then $A + B = 1136.112$, $A - B = -.112$, and

$$x = \frac{\sqrt[7]{1136.112} - \sqrt[7]{-.112}}{2} = \frac{2.732 - .732}{2} = 1,$$

$$\text{And } v = \frac{\sqrt[7]{1136.112} + \sqrt[7]{-.112}}{2} =$$

$$\frac{2.732 + .732}{2} = 1.732 = \sqrt{3}, \text{ and } x + v = 1 + \sqrt{3} \\ = \text{the root.}$$

SCHOLIUM.

In the former method, if $\sqrt[n]{D}$ is not rational, neither member of the root will be rational, and in

in the second, if neither the sum nor difference of $\sqrt[n]{A+B}$ and $\sqrt[n]{A-B}$, is rational; neither member of the root will be so: and in these cases the rules are of no use. Logarithms will be useful here in finding these roots, being exact enough in finding whether any of the quantities be rational or not. When none of these quantities are rational, multiply the given equation by some number, till $\sqrt[n]{D}$, or $\sqrt[n]{A+B} \pm \sqrt[n]{A-B}$, comes out rational; then extract the root as before. But remember to divide the values of x , v , at last, by the root of that number. Thus $22 + \sqrt{486}$ has not such a cube root; but multiply by 2, and then $\frac{44 + \sqrt{1944}}{2}$, will have a cube root, for the numerator.

PROBLEM LXXIII.

To explain the several properties of (o) nothing, and infinity.

It is plain, nothing added to, or subtracted from, any quantity, makes it neither bigger nor less.

Likewise, if any quantity is multiplied by o, that is, taken no times at all; the product will be nothing.

Let $\frac{b}{a} = q$; that is, let the quotient, of b divided by a , be q . Then if b remains the same, it is plain the less a is, the greater the quotient q will be. Let a be indefinitely small beyond all bounds, then q will be indefinitely great beyond all bounds. Therefore when a is nothing, the quotient q will be infinite. Whence

Also since $\frac{b}{o} = \text{infinity}$, therefore $b = \text{nothing} \times \text{infinity}$.

Let there be several geometrical proportionals, $x, x^2, x^3, x^4, x^5, \&c.$ If this series be continued backwards, it will be $x, 1, \frac{1}{x}, \frac{1}{xx}$; that is, x^1, x^0, x^{-1}, x^{-2} , the indices continually decreasing by 1. Then its plane x^0 is equal to 1, whatever x be; for it may stand univervally for any thing: Therefore $o^0 = 1$.

Let x be an indefinitely small quantity, beyond all conception; then in the series $x, x^2, x^3, \&c.$ each term will be indefinitely greater than the following one. And when x is 0, then in the series $\frac{1}{o}, o^0, o^1, o^2, \&c.$ $\frac{1}{o}$ is infinite, and 0 is nothing, by what goes before. Therefore the mean o^0 is a finite quantity. Suppose $=b$, whence $\frac{1}{o} \times o = bb$, that is $bb = \frac{1 \times o}{o} = 1$, and $b = 1$, whence it is plain again, that $(b) o^0 = 1$.

Let $\frac{a}{1-1}$ or its equal $\frac{a}{-1+1}$ be an infinite quantity, then by actually dividing, $\frac{a}{1-1} = a+a+a + \frac{a}{1-1}$, and $\frac{a}{-1+1} = -a-a-a + \frac{a}{-1+1}$. Therefore $\frac{a}{1-1} + a+a+a \&c. = \frac{a}{1-1} - a-a-a - a \&c.$ that is, an infinite quantity is neither increased nor decreased by finite quantities.

Cor. 1. If 0 multiply any finite quantity, the product will be nothing.

Cor.

Cor. 2. If 0 multiply an infinite quantity, the product is a finite quantity. Or a finite quantity is a mean proportional between nothing and infinity.

For $0 \times \text{infinity} = b$.

Cor. 3. If a finite quantity is divided by 0, the quotient is infinite ($\frac{b}{0} = \text{inf.}$).

Cor. 4. If 0 be divided by 0, the quotient is a finite quantity of some sort.

For (Cor. 1.) $b \times 0 = 0$, and therefore $\frac{0}{0} = b$, a finite quantity, or nothing.

Cor. 5. Hence also $0^0 = 1$, or the infinitely small power, of an infinitely small quantity, is infinitely near 1.

Cor. 6. Adding or subtracting any finite quantities to or from an infinite quantity, makes no alteration.

Cor. 7. Therefore in any equation, where are some quantities infinitely less than others; they may be thrown out of the equation.

Cor. 8. An infinite quantity may be considered either as affirmative or negative.

For infinity = $\frac{b}{+0}$ or $\frac{b}{-0}$.

SCHOLIUM.

There is something extremely subtle, and hard to conceive, in the doctrine of *infinites* and *nothings*. Yet although the objects themselves are beyond our comprehension; yet we cannot resist the force of demonstration, concerning their powers, properties, and effects; which properties, under such and such conditions, I think, I have truly explained in this proposition. Any metaphysical notions, that go beyond these mathematical operations, are

not the business of a mathematician. But thus much may be observed, that 0, in a mathematical sense, never signifies absolute nothing; but always nothing in relation to the object under consideration. For illustration thereof, suppose we are considering the area contained between the base of a parallelogram and a line drawn parallel to the base. As this line draws nearer the base, the area diminishes; till at last, when the line coincides with the base, the area becomes nothing. So the area here degenerates into a line; which is nothing, or no part of the area. But it is a line still, and may be compared with other lines.

P R O B L E M LXXIV.

To find the value of a fraction, when the numerator and denominator, is each of them nothing.

I R U L E.

Consider, from the nature of the question proposed, what quantities are infinitely greater than others, when they are all taken infinitely small. Then throw out of the equation, all those terms that are infinitely less than others; retaining only those that are infinitely greater than the rest; by which expunge one of the unknown quantities, and the value of the fraction will be known.

Ex. 1.

Let $x^3 + y^3 = axy$, and y infinitely greater than x , when they vanish; to find the value of $\frac{yy}{x}$, when x and y are $= 0$.

Here x^3 is infinitely less than axy or y^3 , whence $y^3 = axy$, or $yy = ax$. Then $\frac{yy}{x} = \frac{ax}{x} = a$, the value of the fraction proposed.

Ex.

Ex. 2.

If $2ax + xx = yy$, what is the value of $\frac{x}{yy}$, when x and $y = 0$, and y infinitely greater than x .

Here reject xx being infinitely less than the rest; then $yy = 2ax$, and $\frac{x}{yy} = \frac{1}{2a}$.

Ex. 3.

What is the value of $\frac{y}{x}$, when $2ay + yy = rx$; y, x being $= 0$.

Here yy is infinitely less than $2ay$. Whence $2ay = rx$, and $\frac{y}{x} = \frac{r}{2a} = \frac{0}{0}$.

2 R U L E.

Observe what the unknown quantity is equal to, when the numerator, &c. vanishes; put the unknown quantity $=$ that value $\pm e$, where e is supposed infinitely small. Which being substituted for that unknown quantity, and the roots of all surds, extracted to a sufficient number of places of e ; at last you will have some terms in both the numerator and denominator, which will determine the value of the fraction.

Ex. 4.

What is the value of $\frac{a\sqrt{ax-xx}}{a-\sqrt{ax}}$, when $x = a$.

Put $x = a + e$, then expunging x ; $\frac{a\sqrt{ax-xx}}{a-\sqrt{ax}}$

$$\begin{aligned}
 &= \frac{a \times aa + ae^{\frac{1}{2}} - a + e^2}{a - aa + ae^{\frac{1}{2}}} = \\
 & a \times : a + \frac{1}{2}e \text{ \&c.} - aa - 2ae \text{ \&c.} \\
 & \frac{a - a - \frac{1}{2}e \text{ \&c.}}{aa + \frac{1}{2}ae \text{ \&c.} - aa - 2ae \text{ \&c.}} = \frac{-\frac{1}{2}ae}{-\frac{1}{2}e} = \frac{3ae}{e} \\
 &= 3a, \text{ the value of the fraction.}
 \end{aligned}$$

Ex. 5.

What is the value of $\frac{\sqrt{2a^3x - x^4} - a\sqrt{aax^3}}{a - \sqrt{ax^3}}$, when

$x = a$.

Let the fraction $= y$, and put $x = a - e$, then

$$y = \frac{\sqrt{2a^3 \times a - e - a - e^4} - a\sqrt{a^3 - aae}}{a - \sqrt{a \times a - e^3}}. \text{ But}$$

$$\sqrt{2a^3 \times a - e - a - e^4} = \sqrt{2a^4 - 2a^3e - a^2 + 4a^2e} =$$

$$a^2 + 2a^2e^{\frac{1}{2}} = aa + ae \text{ \&c.} \text{ Also } a\sqrt{a^3 - aae} =$$

$$a \times a - \frac{1}{2}ae, \text{ \&c.} \text{ And } \sqrt{a \times a - e^3} =$$

$$a^2 - 3a^2e^{\frac{1}{2}} = a - \frac{1}{2}e \text{ \&c.} \text{ Whence}$$

$$y = \frac{aa + ae \text{ \&c.} - aa + \frac{1}{2}ae \text{ \&c.}}{a - a + \frac{1}{2}e \text{ \&c.}} = \frac{\frac{3}{2}ae}{\frac{1}{2}e} = \frac{16a}{9}$$

Ex. 6.

Let $\frac{a\sqrt{4a^3 + 4x^3} - ax - aa}{\sqrt{2aa + 2ax - x - a}} = y$, what is its value when $x = a$.

Let $a - e = x$. And expunging x ,

$$\frac{a\sqrt{4a^3 + 4 \times a - e^3} - aa + ae - aa}{\sqrt{2aa + 2 \times a - e^2} - x - a} = y. \text{ But}$$

$a\sqrt{\quad}$

$$a\sqrt[3]{4a^3+4xa-e} = a\sqrt[3]{8a^3-12aae+12ace} = 2aa - ae + \frac{1}{2}ee \&c. \text{ And } \sqrt{2aa+2xx} = 4aa-4ae+2ee^{\frac{1}{2}}$$

$$= 2a-e + \frac{ee}{4a} \&c. \text{ Whence}$$

$$y = \frac{2aa-ae+\frac{1}{2}ee\&c.-2aa+ae}{2a-e+\frac{ee}{4a}\&c.-2a+e} = \frac{+\frac{1}{2}ee}{+\frac{ee}{4a}} =$$

$$\frac{2ace}{ee} = 2a.$$

Here, if I had gone no farther than the first power of e , it is evident by inspection, that all the terms would have vanished; by which nothing could have been concluded.

SCHOLIUM.

If e remains at last in the numerator, the value of the fraction is 0, and if e remains in the denominator, the fraction is infinite. But if all the terms vanish out of both numerator and denominator, the series must then be carried to more places, to have a solution.

PROBLEM LXXV.

To find two whole numbers x, y , in the equation $ax=by+c$, being in its least terms: a, b, c , being given numbers.

R U L E.

Let wb . stand for the words *a whole number*.

Reduce the equation, then $x = \frac{by+c}{a} = wb$. By an abridged fraction, I mean the fraction resulting by throwing all whole numbers out of it, till the terms in the numerator be less than the denominator.

minator. Thus let the fraction $\frac{by+c}{a}$ be abridged to $\frac{dy+f}{a}$. Then to find y .

The method consists in lessening the coefficient of y continually, till at last it becomes 1. And this is done by subtracting $\frac{dy+f}{a}$ or some multiple of it, from y , or any multiple of it, which comes very near it; that is, from $\frac{ay}{a}$, $\frac{2ay}{a}$, $\frac{3ay}{a}$, &c. or this from it. And the resulting fraction abridged, or its nearest multiple, is in like manner to be subtracted from the nearest foregoing fraction; or from any wb . which is nearer; or this from that. And these wb . may be $\frac{ay}{a}$, $\frac{2ay}{a}$, &c. or $\frac{ay+a}{a}$, $\frac{ay+2a}{a}$, $\frac{2ay+a}{a}$ &c. or any you can find, which has the nearest coefficient to y . By this means the coefficient of y is continually lessened, till at last we have $\frac{y+g}{a} = wb = p$; then will $y = ap - g$: where p may be any whole number taken at pleasure. And y being known, x will be found from the given equation.

You must observe in this whole process, to keep the same denominator a , throughout.

For whole numbers subtracted from one another, will always leave whole numbers. And whole numbers multiplied by whole numbers, will always produce whole numbers. And upon these principles the rule is founded.

Ex. 1.

Let $19x = 14y - 11$, to find x, y in whole numbers.

By reduction $x = \frac{14y - 11}{19} = wb.$ Also $\frac{19}{19}y$

$= wb.$ Then by subtraction, $\frac{19y}{19} - \frac{14y - 11}{19}$

$= \frac{5y + 11}{19} = wb.$ And multiplying by 4,

$\frac{20y + 44}{19} = \frac{20y + 6}{19} + 2 = wb.$ And $\frac{20y + 6}{19}$

$= wb.$ Subtract $\frac{19y}{19}$; and $\frac{y + 6}{19} = wb. = p.$

Whence $y = 19p - 6.$ Let $p = 1$, for the least affirmative value of y , and $y = 13.$ Whence $x = 9.$

Or thus,

$y = \frac{19x + 11}{14} = x + \frac{5x + 11}{14} = wb.$ Then

$\frac{5x + 11}{14} = wb.$ And multiplying by 3, $\frac{15x + 33}{14}$

$= wb.$ But $\frac{14x + 28}{14} = wb.$ And subtracting,

$\frac{x + 5}{14} = wb. = p.$ And $x = 14p - 5.$ Let $p = 1,$

to have x the least; and $x = 9,$ and $y = 13.$

Ex. 2.

Suppose $3x = 8y - 16$, query $x, y.$

Here $x = \frac{8y - 16}{3} = 2y - 5 + \frac{2y - 1}{3} = wb.$

And $\frac{2y - 1}{3} = wb.$ And multiplying by 2,

$\frac{4y - 2}{3} = wb.$ But $\frac{3y}{3} = wb.$ And their dif-

ference

ference $\frac{y-2}{3} = wb. = p$. Whence $y = 3p + 2$, and taking $p = 0$, $y = 2$. Whence $x = 0$.

Ex. 3.

$$\text{Let } 24x = 13y + 16.$$

Here $x = \frac{13y + 16}{24} = wb.$ multiply by 11, and $\frac{143y + 176}{24} = wb.$ But $\frac{6 \times 24y + 7 \times 24}{24}$ or $\frac{144y + 168}{24} = wb.$ From which subtract the former, and $\frac{y-8}{24} = wb. = p$. And then $y = 24p + 8$, and putting $p = 0$, $y = 8$, and $x = 5$.

Ex. 4.

$$\text{Let } 14x = 4y + 7.$$

Then $x = \frac{4y + 7}{14} = wb.$ And multiplying by 7, $\frac{28y + 49}{14}$ or $\frac{28y + 7}{14} + 3 = wb.$ And $\frac{28y + 7}{14} = wb.$ But $\frac{28y}{14} = wb.$ Therefore their difference $\frac{7}{14} = wb.$ which is absurd; for an even number cannot divide an odd number, nor a greater number a lesser. See Cor. 2. Prop. VIII.
B. II. Arithmetic.

Ex. 5.

$$\text{Let } 27x = 1600 - 16y.$$

Here $x = \frac{1600 - 16y}{27} = wb.$ abridged, $\frac{7 - 16y}{27} = wb.$ or $\frac{16y - 7}{27} = wb.$ Subtract it from $\frac{27y}{27}$, and

and $\frac{11y+7}{27} = wb.$ multiply by 2, and $\frac{22y+14}{27}$

$=wb.$ Subtract it from $\frac{27y+27}{27}$, and $\frac{5y+13}{27}$

$=wb.$ multiply by 2, and $\frac{10y+26}{27} = wb.$ sub-

tract it from $\frac{11y+7}{27}$, and $\frac{y-19}{27} = wb. = p,$ and

$y = p \times 27 + 19,$ and if $p = 0, y = 19,$ and $x = 48.$

Cor. 1. All the values of y are bad, by continual-
ly adding the coefficient of x ; as $y, y+a, y+2a,$
 $y+3a,$ &c. And all the values of x are bad, by
continually adding the coefficient of y ; as $x, x+b,$
 $x+2b,$ &c; or by subtracting them, for negative
numbers, and both are in arithmetical progression.

Cor. 2. When the process brings out an odd num-
ber divided by an even number, or a lesser number di-
vided by a greater, which should be a whole number;
the question is impossible.

Cor. 3. If it be required to find y a whole num-
ber, so that the fraction $\frac{by+c}{a}$ may also be a whole
number. You must proceed the very same way, by
abridging the fraction to $\frac{y+g}{a}$, and then find
 $y = aP + g,$ where P is any whole number, taken at
pleasure.

PROBLEM LXXVI.

To find such a whole number $x,$ that being divided
by the given numbers $a, b, c,$ &c. shall leave the
given remainders $f, g, h,$ &c.

R U L E.

Since the fractions $\frac{x-f}{a}, \frac{x-g}{b}, \frac{x-h}{c}$ &c. are
whole

whole numbers; put the first $\frac{x-f}{a} = P = wb$. Then $x = aP + f$. Put this value of x in the second fraction; then $\frac{aP+f-g}{b} = wb$. Then (Cor. 2. last Prob.) find $P = bQ + m$, where $Q = wb$. then will $x = abQ + am + f$. Put this value of x in the third fraction; then $\frac{abQ + am + f - h}{c} = wb$. Then, as before, find $Q = cR + n$; and put this instead of Q in the last value of x ; then this value of x must be put into the fourth fraction; and proceed the same way through all the fractions. This is the method of proceeding; but numbers must be used all along instead of the small letters. And the least wb . number R may be taken at pleasure.

Ex. 1.

To find a number which divided by 3, 5, 7, and 2; will leave the remainders 2, 4, 6, 0, respectively.

Let the number be x , then $\frac{x-2}{3}$, $\frac{x-4}{5}$, $\frac{x-6}{7}$ and $\frac{x-0}{2}$ are whole numbers. Let $\frac{x-2}{3} = P$, and $x = 3P + 2$; then $\frac{x-4}{5} = \frac{3P+2-4}{5} = \frac{3P-2}{5} = wb$. subtract it from $\frac{5P}{5}$, and $\frac{2P+2}{5} = wb$. Subtract this from $\frac{3P-2}{5}$; then $\frac{P-4}{5} = wb = Q$, and $P = 5Q + 4$, and $x = 15Q + 14$. Again $\frac{x-6}{7} = \frac{15Q+8}{7} = wb$. and $\frac{Q+1}{7} = wb = R$, and $Q = 7R - 1$, and $x = 105R - 1$.
Lastly,

Sect. VII. PROBLEMS. 221

Lastly $\frac{x-0}{2} = \frac{105R-1}{2} = wb.$ and $\frac{R-1}{2} = wb.$
 $=S$, and $R=2S+1$. Whence $x=210S+104$,
 the number fought; and putting $S=0$, the least
 value of x is 104.

Ex. 2.

To find a whole number, which being divided by
 16, 17, 18, 19, 20; will leave 6, 7, 8, 9, 10,
 remainders.

Let x = number. Then $\frac{x-6}{16}$, $\frac{x-7}{17}$, $\frac{x-8}{18}$,
 $\frac{x-9}{19}$, $\frac{x-10}{20}$ are whole numbers. Put $\frac{x-6}{16} = P$,
 then $x = 16P + 6$.

Then $\frac{x-7}{17} = \frac{16P-1}{17} = wb.$ And thence
 $\frac{P+1}{17} = wb. = Q$, and $P = 17Q - 1$, and
 $x = 272Q - 10$.

Also $\frac{x-8}{18} = \frac{272Q-18}{18} = wb.$ and $\frac{2Q}{18} = wb.$
 $=R$, and $Q = 9R$, whence $x = 2448R - 10$.

Again $\frac{x-9}{19} = \frac{2448R-19}{19} = wb.$ and
 $\frac{-3R}{19} = wb.$ or $\frac{3R}{19} = wb.$ and $\frac{18R}{19} = wb.$ whence
 $\frac{R}{19} = wb. = S$, and $R = 19S$. Then
 $x = 46512S - 10$.

Lastly $\frac{x-10}{20} = \frac{46512S-20}{20} = wb.$ and
 $\frac{12S}{20} = wb. = T$, and $S = 5T$. Whence
 $x = 232550T - 10$. And if $T = 1$, then the
 least value of $x = 232550$.

Ex.

Ex. 3.

To find a number (x), which being divided by 3, 7, 14, 20; there shall remain 1, 3, 7, 14.

Here $\frac{x-1}{3}$, $\frac{x-3}{7}$, $\frac{x-7}{14}$, $\frac{x-14}{20}$ are whole numbers. Let $\frac{x-1}{3} = P$, and $x = 3P + 1$.

Then $\frac{x-3}{7} = \frac{3P-2}{7} = \text{wb.}$ and $\frac{6P-4}{7} = \text{wb.}$

Whence $\frac{P+4}{7} = \text{wb.} = Q$, and $P = 7Q - 4$, and $x = 21Q - 11$.

Also $\frac{x-7}{14} = \frac{21Q-18}{14} = \text{wb.}$ and $\frac{7Q-4}{14} = \text{wb.}$ and $\frac{14Q-8}{14} = \text{wb.}$ Whence $\frac{8}{14} = \text{wb.}$ which is absurd.

Hence the question is impossible for the three first suppositions; but will hold good for two of them: in which case $x = 21Q - 11$, where the least value of x is 10.

2 R U L E.

When two divisors and their remainders are given; then find two fixed multipliers M , N ; such, that dividing them,

$\frac{M}{a}$ leaves 0, and $\frac{M}{b}$ leaves 1 remaining.

and $\frac{N}{a}$ leaves 1, and $\frac{N}{b}$ leaves 0 remaining.

Then divide $\frac{Mg + Nf}{ab}$, and the remainder is x , the number sought.

Likewise

Likewise for three divisors and remainders; find three fixed multipliers M , N , P ; such, that by dividing them,

$\frac{M}{a}$ leaves 1, and $\frac{M}{bc}$ leaves 0, remaining.

$\frac{N}{b}$ leaves 1, $\frac{N}{ac}$ leaves 0, remaining.

$\frac{P}{c}$ leaves 1, $\frac{P}{ab}$ leaves 0, remaining.

Then dividing $\frac{Mf + Ng + Pb}{abc}$, the remainder is x , required; and the like for more quantities.

To prove the truth of this. Since (Case 1)

$\frac{M}{a}$ as also $\frac{N}{b}$ leave 0, by division; therefore

$\frac{Mg}{a}$, and $\frac{Nf}{b}$ leave 0.

And since $\frac{M}{b}$, as also $\frac{N}{a}$ leave 1. Therefore

$\frac{M-1}{b}$ and $\frac{N-1}{a}$ leave 0. Therefore $\frac{Mg-g}{b}$

and $\frac{Nf-f}{a}$ leave 0; that is, $\frac{Mg}{b}$ leaves g , and

$\frac{Nf}{a}$ leaves f . Therefore $\frac{Mg+Nf}{a}$ leaves $0+f$,

and $\frac{Mg+Nf}{b}$ leaves $g+0$.

But since $Mg+Nf$ may exceed ab , and therefore is not the least number; therefore divide by ab , and the remainder is the least number required. And the same way, Case 2, or any other, is proved.

Ex. 4.

Having the cycle of the dominical letter f , and cycle of the moon g ; to find the year of the Dionysian period.

Let x be the year sought. Then $\frac{x-f}{28}$ and $\frac{x-g}{19}$ are whole numbers. Here $a=28$, and $\frac{M}{28} = wb. = P$, and $M=28P$. Also $\frac{M-1}{19} = wb. = \frac{28P-1}{19}$; and multiplying by 2, $\frac{56P-2}{19} = wb.$ Also $\frac{57P}{19} = wb.$ Therefore $\frac{P+2}{19} = wb. = Q$, and $P=19Q-2$. Whence $M=28 \times 19Q-2 = 532Q-56$, and if $Q=1$; then $M=476$.

Then $\frac{N}{b} = \frac{N}{19} = wb. = P$, and $N=19P$. Also $\frac{N-1}{28} = \frac{19P-1}{28} = wb.$ multiply by 3; then $\frac{57P-3}{28} = wb.$ and $\frac{56P}{28} = wb.$ therefore $\frac{P-3}{28} = wb. = Q$, and $P=28Q+3$. Whence $N=28 \times 19Q+57$, and if $Q=0$, $N=57$.

Therefore $x =$ remainder of $\frac{476g+57f}{532}$, which serves in general, for any numbers, f ; g .

Let $f=10$, $g=12$; then $x=430$.

Ex. 5.

Having the cycle of the Sunday letter f , the golden number g ; and indiction b ; to find the year of the Julian period.

Here $a=28$, $b=19$, $c=15$, $ab=532$, $ac=420$, $bc=285$, and $abc=7980$.

Then

Then $\frac{M}{285} = wb. = P$, and $M = 285P$. Also
 $\frac{M-1}{28} = \frac{285P-1}{28} = wb.$ This at last gives
 $P = 17$ the least; and then $M = 4845$.

Again, $\frac{N}{420} = wb. = P$, and $N = 420P$. Also
 $\frac{N-1}{19} = \frac{420P-1}{19} = wb.$ and $\frac{2P-1}{19} = wb.$ which
 will give $P = 10$, and $N = 4200$.

Lastly, $\frac{P}{532} = wb. = Q$, and $P = 532Q$. Also
 $\frac{P-1}{15} = \frac{532Q-1}{15} = wb.$ and $\frac{7Q-1}{15} = wb.$ at
 last $Q = 13$, and $P = 6916$. Whence the re-
 mainder of $\frac{4845f + 4200g + 6916b}{7980}$ is $= x$.

Let $f = 0$ or 28 , $g = 1$, $b = 2$; then $x = 2072$.

Cor. When the operation brings out, a lesser number divided by a greater, instead of a whole number; the problem is impossible.

PROBLEM LXXVII.

An equation being given, containing several unknown quantities; to find their limits.

When an equation contains several unknown quantities, the values of all of them, except one, may be taken at pleasure; and when their values are assigned, and numbers put for them in the equation, that single quantity may also be found, by reducing the equation. And such equations will admit of an infinite number of solutions, if we admit of fractional and negative numbers. But since these solutions are most useful where affirmative quantities are concerned; and more useful still, when only affirmative whole numbers are admitted;

Q

admitted; therefore I propose to consider only these two cases, and particularly the last: because in that case such an equation will have a determined number of solutions. And therefore it is necessary to know the limits of the unknown quantities; lest we go about to seek their values beyond these limits.

R U L E.

Transpose the negative quantities to the contrary side; that all the terms may be affirmative. Then to find the limits of any one, put all the rest $=0$, or suppose them to vanish; and from hence find the value of that quantity, which will be one limit thereof. And to know which limit it is, conceive the other quantities to increase and have some certain value; then if by this, the value (of the unknown quantity under consideration) increases; it is the least limit you found; if it decreases, it is the greatest limit. And in case you find no least limit, then 0 is its least limit. This process relates to fractional quantities.

But if you only desire whole numbers; put 1 for each of the other quantities, which is the least value they can have; then from the resulting equation, find your unknown quantity and its limit, as before directed.

Proceed the same way with all the unknown quantities.

Ex. 1.

Let $3a + 5e = 28$, to find the limits of a , e .

Let $e = 0$, then $3a = 28$, and $a = \frac{28}{3} = 9\frac{1}{3}$.

Now let e be some real quantity; it is plain the greater e is, the less a must be; therefore $9\frac{1}{3}$ is the greater limit. Whence $a = 9\frac{1}{3}$.

For

For e : let $a=0$, then $5e=28$, and $e=\frac{28}{5}=5\frac{3}{5}$. But if a increases, e decreases; therefore $5\frac{3}{5}$ is the greater limit, and $e \searrow 5\frac{3}{5}$, and the lesser limit of both a and e , is 0. All this including fractions.

For whole numbers.

Let $e=1$, then $3a=28-5=23$, and $a=\frac{23}{3}=7\frac{2}{3}$, the lesser limit, and $a \searrow 7\frac{2}{3}$.

Again, let $a=1$, then $5e=25$, and $e=\frac{25}{5}=5$, the greater limit, and $e \searrow$ or $\searrow 5$.

Ex. 2.

Let $3a-5e=28$, to find the limits of a , e in whole numbers.

Then $3a=28+5e$; let $e=1$, then $3a=33$, and $a=\frac{33}{3}=11$; but when e increases a increases, therefore 11 is the lesser limit, and $a \searrow$ or $\searrow 11$.

Let $a=1$, $28+5e=3$, and e will be negative, which we exclude. But whilst a increases e increases; therefore 0 is the least limit of e , or $e \searrow 0$: and it has no greatest limit.

Ex. 3.

Let $3x+5y+8z=10003$, to find the limits in whole numbers.

Suppose $y=1=z$. Then $3x=10003-13$, and $x=\frac{9990}{3}=3330$. And since x decreases, whilst y and z increase; therefore 3330 is the greater limit, and $x \searrow$ or $\searrow 3330$.

Again, let $x=z=1$; then $5y=9992$, and $y = \frac{9992}{5} = 1998\frac{2}{5}$; and y decreases whilst x, z increase: whence $y \supset 1998\frac{2}{5}$.

Lastly, for z ; let $x=y=1$, then $8z=10003-8=9995$, and $z = \frac{9995}{8} = 1249\frac{3}{8}$. But z decreases, whilst x, y increase; therefore $z \supset 1249\frac{3}{8}$.

Ex. 4.

Let $13x-5y+8z=10$, to find the limits of y .

Here $13x+8z=10+5y$; let $x=1=z$; then $5y+10=21$, and $5y=11$, and $y=2\frac{1}{5}$; and whilst x and z increase, y increases; therefore $2\frac{1}{5}$ is the least limit, and $y \supset 2\frac{1}{5}$.

Note, the limits of x and z cannot be found till the value of y be assigned.

P R O B L E M LXXVIII.

Two equations being given, containing three or more unknown quantities; to determine their limits.

R U L E.

Having pitched upon the quantity you would limit; expunge one of the other quantities, and you will have one limiting equation. Then expunge another of them, and this gives another limiting equation. By these two equations find the limits of the quantity pitched on separately, by the last problem.

But note, In any limiting equation, all the other unknown quantities therein, (being put on the same side of the equation, with the absolute number,) must have the same sign: otherwise, (if they have different signs) they cannot limit the quantity proposed, till the value of some of the rest be known.

If

If there be more equations, the process is the same with any of them.

Ex. 1.

Let $a + e + y = 56$.
and $32a + 20e + 16y = 1232$; to limit a .

Multiply the first equation by 20, produces $20a + 20e + 20y = 1120$. Subtract this from the second, and you have $12a - 4y = 112$: whence (Prob. lxxvii.) $a \sqsupset 9\frac{1}{2}$.

Multiply the first equation by 16, gives $16a + 16e + 16y = 896$. Subtract it from the second, and $16a + 4e = 336$; whence $a \sqsupset 20\frac{1}{2}$.

In like manner, to limit y , multiply the first equation by 32, and $32a + 32e + 32y = 1792$. Subtract the second from it, and $12e + 16y = 560$. This gives $y \sqsupset 34\frac{1}{2}$. And the equation $12a - 4y = 112$, gives $y \sqsupset 0$.

To limit e ; the equation $16a + 4e = 336$, gives $e \sqsupset 80$. And the equation $12e + 16y = 560$, gives $e \sqsupset 45\frac{1}{2}$. But here is no lesser limit for e ; therefore $e \sqsupset 0$, and $\sqsupset 45$.

Ex. 2.

Let $3x - y + 2u = 20$
and $12x + 6y + 5u = 150$

1st $\times 5$ is, $15x - 5y + 10u = 100$
2d $\times 2$ is, $24x + 12y + 10u = 300$
difference $9x + 17y = 200$

This equation gives $x \sqsupset 20\frac{1}{2}$, and $y \sqsupset 11\frac{1}{2}$.

1st $\times 6$ is, $18x - 6y + 12u = 120$; add this to the second: then $30x + 17u = 270$,

whence $x \sqsupset 8\frac{1}{3}$, and $u \sqsupset 14\frac{1}{3}$.

1st $\times 4$ is $12x - 4y + 8u = 80$. Subtract from the second,
 $10y - 3u = 70$.

And $y \sqsupset 7\frac{1}{2}$, and $u \sqsupset 0$.

There is no least limit for x ; therefore $x \sqsupset 0$, and $\sqsupset 8\frac{1}{3}$.

Ex. 3:

Let $a + e + y + u = 100$.and $16a + 10e + 8y + 6u = 1200$.

		1	$a + e + y + u = 100$	
		2	$16a + 10e + 8y + 6u = 1200$	
$1 \times (6)$		3	$6a + 6e + 6y + 6u = 600$	
$2 - 3$		4	$10a + 4e + 2y = 600$	
$4 \div$		5	$a = \frac{600 - 4e - 2y}{10} = 59\frac{4}{10}$ at most	
5		6	$a \geq 59\frac{4}{10}$	
$1 \times (10)$		7	$10a + 10e + 10y + 10u = 1000$	
$2 - 7$		8	$6a - 2y - 4u = 200$	
8 tr.		9	$6a = 200 + 2y + 4u$	
$9 \div (6)$		10	$a = \frac{200 + 2y + 4u}{6} = 34\frac{1}{3}$ at least, and $a \leq 34\frac{1}{3}$	
			So a is between $34\frac{1}{3}$ and $59\frac{4}{10}$. Then for the other quantities.	
$1 \times (8)$		11	$8a + 8e + 8y + 8u = 800$	
$1 \times (16)$		12	$16a + 16e + 16y + 16u = 1600$	
$2 - 11$		13	$8a + 2e - 2u = 400$. This equation will limit u but not e . Here $u \leq 0$.	
$12 - 2$		14	$6e + 8y + 10u = 400$	
14		15	$u = \frac{400 - 6e - 8y}{10} = 38\frac{6}{10}$, or $u \leq 38\frac{6}{10}$	
4		16	$y = \frac{600 - 4e - 10a}{2} = 293$, or $y \leq 293$; but, since the limits of a , are known; y may be determined more exactly; thus	
			$y = \frac{596 - 10 \times 35}{2} = 123$, or $y \leq 123$.	

14 | 17 Again $y = \frac{400-16}{8} = \frac{384}{8} = 48$, or
 $y =$ or $\neg 48$. But these are all
 greater limits of y , and there wants
 the lesser limit; therefore $y \sqsubset 0$, and
 $\neg 48$.

14 | 18 $e = \frac{400-18}{6} = \frac{382}{6} = 63\frac{2}{3}$, or $e = \neg$
 $63\frac{2}{3}$. But the least limit of e can-
 not be found; therefore take $e \sqsubset 0$.

SCHOLIUM.

When three numbers are sought by two equations; all the values of each of them, in whole numbers, make three series of arithmetical progression, taken within the limits of these numbers. And if four or more numbers are sought, the value of each is to be found in several arithmetical progressions. But yet the values of any three will be in arithmetic progression, when the values of all the rest are assigned, as before for three numbers.

For in the case of three numbers, and two equations; any one of the three may be expunged; and then you will have but one equation, and two unknown quantities; which brings it under Prob. lxxv. But by Cor. 1. of that problem, these two remaining quantities are contained in two series of arithmetical progression. And as any of the three may be expunged; therefore any two of them will constitute two series of arithmetical progression.

PROBLEM LXXIX.

The prices of several ingredients being given, to find the quantities thereof; so that the mixture may be sold at a given price.

Suppose four simples A, B, C, D, are to be mixed; and their prices to be as follows:

Q 4

Mean

$$\begin{aligned}
 \text{Mean price} &= m \\
 \text{Price of A} &= m+a \\
 \text{of B} &= m+b \\
 \text{of C} &= m-c \\
 \text{of D} &= m-d,
 \end{aligned}$$

And let the quantities to be taken of A, B, C, D, be x, y, z, v , respectively. Place them in order, thus :

$$\begin{array}{r}
 \text{prices} \quad \text{quantities} \\
 m \left\{ \begin{array}{l|l}
 m+a & x \\
 m+b & y \\
 m-c & z \\
 m-d & v
 \end{array} \right.
 \end{array}$$

Then by the nature of the question ; if each quantity be multiplied by its price, the sum of the products will be equal to the sum of all the quantities multiplied by the mean price ; that is,

$$\left. \begin{array}{l}
 \overline{m+a} \times x + \overline{m-d} \times v \\
 + \overline{m+b} \times y + \overline{m-c} \times z
 \end{array} \right\} = \overline{x+v+y+z} \times m.$$

$$\text{Let } \overline{m+a} \times x + \overline{m-d} \times v = \overline{x+v} \times m.$$

$$\text{And } \overline{m+b} \times y + \overline{m-c} \times z = \overline{y+z} \times m.$$

That is,

$$mx + ax + mv - dv = mx + mv$$

$$ny + by + mz - cz = my + mz.$$

By the former, $ax - dv = 0$, or $ax = dv$.

By the latter, $by - cz = 0$, or $by = cz$.

Now since x and y may be taken at pleasure. Therefore put $x = d$, and $y = c$. Then will $v = a$, and $z = b$. Whence the quantities will be ranged thus :

$$m \left\{ \begin{array}{l} m+a \\ m+b \\ m-c \\ m-d \end{array} \right. \left| \begin{array}{l} d \\ c \\ b \\ a \end{array} \right. \text{ which gives this}$$

R U L E.

Couple every greater rate with one lesser than the mean price ($m+a$ and $m-d$; also $m+b$ and $m-c$); then take the difference between each rate and the mean rate, and place it *alternately*, that is, against the quantity it is coupled with; do the same with all the rates, (thus place a against $m-d$, b against $m-c$, c against $m+b$, d against $m+a$); then if none of the quantities of A, B, C, D, be given. Then d , c , b , a will be the quantities of each to be taken for the mixture. But if any one quantity be given; then all the quantities d , c , b , a must be increased or decreased in proportion. Or if the sum of the quantities be given, then other quantities must be taken in proportion, so that $d+c+b+a$ may be to the sum given, as any of the differences d , c , &c. to the respective quantity required. And this is the common rule of *Alligation Alternate*.

Again,

Since $ax=dx$, and $by=cx$. Take $x=md$, and $y=nc$; then $v=ma$, and $z=nb$. Then putting md , nc , nb , ma , for x , y , z , v respectively; and the case will stand thus:

$$m \left\{ \begin{array}{l} m+a \\ m+d \\ m-c \\ m-d \end{array} \right. \left| \begin{array}{l} md \\ nc \\ nb \\ ma \end{array} \right. \text{ which gives this}$$

R U L E.

Having coupled the rates as before directed, and taken the differences. Then instead of any couple

couple of the differences, you may take any equimultiples thereof; and place them alternately. And these (or other quantities proportional to them), will be the quantities required. And this is the *Rule of Alligation improved*.

PROBLEM LXXX.

If the numbers *A* and *B* be produced from *a* and *b*, by any similar operation; to find the number from which *N* is produced, by the like operation. Supposing the differences of the numbers *A*, *B*, *N*, to be as the differences of *a*, *b*, and the unknown number.

Let *z* be the number sought, $\begin{matrix} a & b & z \\ A & B & N \end{matrix}$ and put the differences $N - A = r$, $N - B = s$. Then by the question, $r(N - A) : s(N - B) :: z - a : z - b$. Then $rz - rb = sz - sa$. And by transposition, $rz - sz = rb - sa$, and $z = \frac{rb - sa}{r - s}$; or if *s* be negative (or *B* greater than *N*), then $z = \frac{rb + sa}{r + s}$, the number sought.

Cor. i. Hence is derived the practice of the double Rule of False. For if both *A* and *B* be lesser than *N*, or both greater; then $z = \frac{rb - sa}{r - s}$. But if only one as *B* be greater than *N*, then *s* is negative, and $z = \frac{rb + sa}{r + s}$.

That is, if each supposed number be multiplied by the error of the other, and the difference of the products be divided by the difference of the errors, when the errors are like; or the sum of the products divided by the sum of the errors, when the errors are unlike; the quotient gives the number sought.

Cor.

Cor. 2. Hence also is derived another method of working the Rule of Approximation, or Rule of False, which is this.

Multiply the difference of the supposed numbers, by the least error, and divide the product, by the difference of the errors, if like; or by the sum if unlike. The quotient is the correction of the number belonging to the least error.

Then this correction is to be added or subtracted, according as that number was too little or too great.

For let s be the least error, being the error of b , and q = the correction; then if A, B be less than N , $b+q=z$, and $q=z-b=\frac{rb-sa}{r-s} - b =$

$$\frac{rb-sa-rb+sb}{r-s} = \frac{b-a}{r-s} s.$$

But if B is greater than N , then $b-q=z$, and

$$q=b-z=b-\frac{rb+sa}{r+s} = \frac{rb+sb-rb-sa}{r+s} =$$

$$\frac{b-a}{r+s} s.$$

SCHOLIUM.

Since it has been shewn, that the number sought will come out exactly, by this rule, when the errors are exactly proportional to the differences of the supposed numbers from the true one. Therefore it follows, that when the errors are nearly proportional to these differences, that the answer will come out nearly true. And these proportions will be the nearer to an equality, the nearer these supposed numbers are taken to the true number. And therefore in all questions where this rule is applied, every operation will bring us nearer the true answer, if we always take the nearest numbers, (where the errors are least) for new suppositions. And thus repeating the operation, one may

may continually approximate to the true number, within any degree of exactness required; let the particular question be of what nature it will.

Upon this rule also is founded the rule of finding proportional parts.

P R O B L E M LXXXI.

Suppose A, B, C, D, &c. to be several sorts of goods; and m, n, p, q, &c. given numbers; and the values of these goods are

$$mA = nB$$

$$pB = qC$$

$$rC = sD$$

$$tD = vE$$

To find what quantity of the last sort is equal to a given quantity of the first: and the reverse.

Let z times the last be $=y$ times the first, that is, let $zE = yA$.

Multiply all these equations together; the first side by the first, and the second by the second. Then we have

$mA \times pB \times rC \times tD \times zE = nB \times qC \times sD \times vE \times yA$. Then $mprtz = nqsvy$. Then if the quantity of the last

sort be required, $z = \frac{nqsvy}{mprt}$. But if the quantity

of the first sort be sought; $y = \frac{mprtz}{nqs}$. Whence this

R U L E.

Place the terms in two columns, so that there may not be two terms of a sort in either column. Then multiply the numbers in the lesser column for a divisor; and the numbers in the greater column (with the odd term) for a dividend. The quotient

quotient is the quantity of that sort which stands single in the two columns. And this is the *Rule of Exchange* in arithmetic.

PROBLEM LXXXII.

To investigate numbers for rational squares, cubes, &c.

Problems of this sort are often capable of an infinite number of answers; and yet none of the quantities can be assumed at pleasure, but must be investigated as follows.

R U L E.

Put one or more letters to denote the root of the square, cube, &c. Which letters must be so assumed, that when the equation is involved, either the given number, or the highest power of the unknown quantity, may be on both sides of the equation, and consequently vanishes out of it. And then if the unknown quantity be but of one dimension, the problem is solved, by reducing the equation. But if the unknown quantity is still a square or higher power; you must farther assume other new letters, to denote the root, and proceed as before; till you get the unknown quantity of one dimension; and from this unknown quantity all the rest are to be determined. For the whole art is, so to denote the root of the given power, that the unknown quantity may be reduced to one dimension.

But no general rule of proceeding can be given to suit all cases; and therefore the solution will often be left to the sagacity of the analyst, in contriving such a designation of letters as is proper for the purpose.

Ex. 1.

To find two such numbers, so that the sum of their squares is a square.

Let x, y, z be the roots of the squares, so that $xx+yy=zz$. Assume $z=y+r$, then $xx+yy=zz=yy+2ry+rr$, and $xx=2ry+rr$, and $2ry=xx-rr$, where y the unknown quantity is of one dimension, which reduced gives $y = \frac{xx-rr}{2r}$; and

$y+r = \frac{xx-rr}{2r} + r = \frac{xx+rr}{2r} = z$. Therefore the

numbers are $x, \frac{xx-rr}{2r}$ and $\frac{xx+rr}{2r}$, where x and r denote any numbers taken at pleasure.

But if the answer is required in whole numbers, then $2rx, xx-rr, xx+rr$ will denote the roots of the squares, where the sum of the two first is equal to the last square.

Cor. The three sides of a right-angled triangle will only be commensurable, when $xx+rr$ denotes the hypotenuse, and $xx-rr$, and $2rx$ the two sides; x, r being any numbers taken at pleasure, so as x is greater than r .

Ex. 2.

To find two numbers, the sum of whose squares is equal to the sum of two given squares.

Let x, y be the roots; aa, bb the given squares. Assume $x=a-v, y=vz-b$. Then $xx+yy=aa+bb=aa-2av+vv+vvzz-2bvz+bb$; and $vv+vvzz=2av+2bvz$, and $v+vvzz=2a+2bz$; and $v = \frac{2a+2bz}{zz+1}$. Where z is any number taken at pleasure.

Then $x = \frac{azz-2bz-a}{zz+1}$, and

$y = \frac{2ax+bzx-b}{zz+1}$.

Or

Or thus,

Let $x = a - v$, then $aa - 2av + vv + yy = aa + bb$; and
 $yy - 2av + vv = bb$. Put $y = vz - b$; then $vvzz -$
 $2bzv + bb - 2av + vv = bb$, and $vvzz + vv = 2bzv +$
 $2av$, or $vzz + v = 2bz + 2a$, and $v = \frac{2bz + 2a}{zz + 1}$,
 as before.

Ex. 3.

To find two numbers, such that when either of them
 is added to the square of the other, the sum will be a
 square number.

Let the numbers be x, y ; then $xx + y = \square$,
 and $yy + x = \square$. Let $xx + y = r - x = rr - 2rx$
 $+ xx$; then $y = rr - 2rx$, and $2rx = rr - y$, whence
 $x = \frac{rr - y}{2r}$.

Again, assume $yy + x$ or $yy + \frac{rr - y}{2r} = y + v^2 =$
 $yy + 2yv + vv$. Then $\frac{rr - y}{2r} = 2yv + vv$, whence
 $rr - y = 4ryv + 2rvv$, and $4ryv + y = rr - 2rvv$;
 whence $y = \frac{rr - 2rvv}{4rv + 1}$. And $x = \frac{2rvv + vv}{4rv + 1}$,
 where r, v may be taken at pleasure, provided
 r be greater than $2vv$.

Otherwise,

Since $x = \frac{rr - y}{2r} = \frac{1}{2}r - \frac{y}{2r}$, and $yy + x$ or $yy -$
 $\frac{y}{2r} + \frac{1}{2}r = \square$, put $yy - \frac{y}{2r} + \frac{1}{2}r = yy + \frac{y}{2r} + \frac{1}{2}rr =$
 $\left(\frac{y - \frac{1}{4}r}{2r}\right)^2$. Then $\frac{1}{2}r = \frac{1}{16rr}$, and $r^3 = \frac{1}{16}$, which is
 a cube

a cube number. And therefore will answer the question; and we have $r = \frac{1}{2}$; whence $x = \frac{1}{4} - y$, and y may be any thing less than $\frac{1}{4}$.

Ex. 4.

To find two numbers in a given ratio, so that either of them added to the square of the sum, may make a square.

Let the ratio of the two numbers be as b to c , and put $b+c=d$, and let the numbers be bx and cx . Then the square of the sum is $(bx+cx)^2 = ddx$. Therefore $ddxx+bx = \square$, and $ddxx+cx = \square$.

Put $ddxx+bx = dx-v^2 = ddx-2dxv+vv$; then $bx = vv-2dxv$, or $bx+2dxv = vv$, and $x = \frac{vv}{b+2dv}$.

Then $ddxx+cx$ or $\frac{ddv+bc+2cdv}{b+2dv} \times x = \frac{ddv+bc+2cdv}{b+2dv} \times \frac{vv}{b+2dv} = \square$, but $\frac{vv}{b+2dv} = \square$; therefore

$ddv+bc+2cdv = \square$ (See Cor. 27. II. Arithm.); assume $ddv+bc+2cdv = dv-z^2 = ddv-2dvz+zz$; then $2cdv+2dzv = zz-bc$; and $v = \frac{zz-bc}{2cd+2dz}$. Where zz must be greater than

bc , and expunging v , $x = \frac{zz-bc}{4ddz \times b + z \times c + z}$.

SCHOLIUM.

It appears from these operations, that when a quantity, which is to be a square by the problem, is not an algebraic square; we must make it so, by assuming some new quantities to compleat it. Then these squares being compared, an equation is had for determining the unknown quantity. And

in working, one may multiply or divide by any quantity which is a square, and what is left will be a square, in a more simple form. The like for other powers.

PROBLEM LXXXIII.

To determine the maximum or minimum of a quantity proposed.

When a quantity is required to be the greatest or least possible, it is called a *maximum* or *minimum*. And at the time it becomes such, it is at a stand, and at that moment neither increases nor decreases. Therefore to compute it.

R U L E.

Calculate the value of the maximum or minimum two different ways, which is done by increasing the unknown quantity therein, by an exceeding small part; then these values are to be put equal to one another. The same must be done, if there be several variable quantities. But go no farther than the first power of the small added part. Or,

If the maximum or minimum consists of two parts; compute the exceeding small increment of one, and the decrement of the other; and put them equal to one another.

Ex. 1.

What fraction is that whose square exceeds its cube the greatest possible.

Let x be the fraction, then $x^2 - x^3 = \text{max}$. Take e an exceeding small part to be added to x , then you will also have $\overbrace{x+e}^2 - \overbrace{x+e}^3 = \text{max}$. that is, $2x + 2xe - x^3 - 3x^2e = \text{max}$. Whence $x^2 - x^3 = 2x^2 + 2xe - x^3 - 3x^2e$, and $2xe - 3x^2e = 0$, or $3x^2e = 2xe$, and $3x = 2$, or $x = \frac{2}{3}$.

R

Or

Or thus,

Since $x^2 - x^3 = \max.$ let e be the small increase of x , then $2xe$ is the increment of xx , and $3x^2e$ is the decrement of x^3 ; therefore $2xe = 3x^2e$, and $x = \frac{3}{2}$ as before.

Ex. 2.

To divide a given quantity into two parts, that one of the parts multiplied by the cube of the other part; the product may be a maximum.

Let a be the quantity, and x one part, and $a-x$ the other part, and e a small additional part to x . Then $x^3 \times a-x$ or $ax^3 - x^4 = \max. = ax^3 + 3ax^2e - x^4 - 4x^3e$. Then $3ax^2e = 4x^3e$, and $x = \frac{3}{4}a$, for one part, and $a-x = \frac{1}{4}a$, the other part.

Ex. 3.

To find $a^3 - a^2x + x^3$ a minimum, x being unknown.

Put $x+e$ for x . Then $a^3 - a^2x + x^3 = \min. = a^3 - a^2x - a^2e + x^3 + 3x^2e$, and $-a^2e + 3x^2e = 0$, and $3xx = aa$; whence $x = a\sqrt{\frac{1}{3}}$. Then $a^3 - a^2x + x^3 = a^3 - a^2\sqrt{\frac{1}{3}} + \frac{1}{3}a^3\sqrt{\frac{1}{3}} = a^3 \times 1 - \frac{2}{3}\sqrt{\frac{1}{3}}$, the minimum.

Ex. 4.

Let $\frac{baax + aaxx - bx^3 - x^4}{baa + x^3} = a + x$ be a maximum.

This reduced to a common denominator is $\frac{2bbaax + aaxx - bx^3 - ba^3 - ax^3}{baa + x^3} = \max.$ Put $x+e$ for x ,

Then

$$\text{Then } \frac{2baax + aaxx - bx^3 - ba^3 - ax^3}{baa + x^3} =$$

$$\frac{2baax + 2baae + aaxx + 2aaxe - bx^3 - 3bx^2e - ba^3 -}{baa + x^3 + 3xxe} =$$

$$\frac{ax^3 - 2ax^2e}{};$$

Then multiplying alternately,

$$2baax + aaxx - bx^3 - ba^3 - ax^3 \times baa + x^3 + 3xxe =$$

$$: 2baax + 2baae + aaxx + 2aaxe - bx^3 - 3bx^2e - ba^3 -$$

$$ax^3 - 3ax^2e : \times baa + x^3. \text{ And throwing out}$$

what is common on both sides,

$$2baax + aaxx - bx^3 - ba^3 - ax^3 \times 3xxe = \overline{baa + x^3}$$

$$\times 2baae + 2aaxe - 3bxxe - 3axxe. \text{ That is (dividing}$$

by e), $6baax^3 + 3aax^4 - 3bx^5 - 3ba^3xx - 3ax^5 =$
 $2bba^4 + 2ba^4x - 3bbaaxx - 3ba^3xx + 2baax^3 + 2aax^4$
 $- 3bx^5 - 3ax^5. \text{ Reduced, } 4baax^3 + aax^4 = 2bba^4 +$
 $2ba^4x - 3bbaaxx; \text{ or dividing by } a, \text{ and transposing,}$
 $x^4 + 4bx^3 + 3bbxx - 2baax - 2bbaa = 0.$

Ex. 5.

Suppose $y^3 - 3yyx + 3yxx = nyx - nxx.$
 and $x - y = mn.$

Suppose the maximum $= m.$ Then $x = m + y.$
 This substituted in the first equation, and reduced,
 gives $y^3 + 3m^2y = 2yy - mn.$ And $y^3 + 12mmy -$
 $nyy + 4mnn = 0.$ Where m is a fixt quantity. Put
 $y + e$ for y ; then $y^3 + 12m^2y - nyy + 4m^2n = 0 =$
 $y^3 + 3y^2e + 12m^2y + 12m^2e - ny^2 - 2nye + 4m^2n = 0,$
 and $3y^2e + 12m^2e - 2nye = 0,$ whence $3yy + 12m^2 -$
 $2ny = 0,$ or $2ny - 3yy = 12mm.$ From this equa-
 tion, and $y^3 + 12m^2y - ny^2 + 4m^2n = 0,$ the quan-
 tities y and m will easily be determined.

Ex. 6.

Through a given point P within the angle BAC , Fig. 1.
 to draw a right line BPC , making the area of the
 triangle BAC , the least possible.

Draw AP , and bPc extremely near BPC ; then
 the area $ABP + ACP = \text{minimum.}$ In the very
 small

Fig. small triangles BPb , and CPc , the vertical angles
 1. at P are equal, and $BP = bP$, as also $CP = cP$,
 extreme near. Therefore the areas BPb , and CPc ,
 are to one another as BP^2 to CP^2 (Geom. 19. II).
 But CPc is the increment of the area APC ; and
 BPb is the decrement of the area APB . There-
 fore $BPb = CPc$, or $BP^2 = CP^2$; therefore
 $BP = CP$. Whence if PD be drawn parallel to
 CA , then $DB = DA$.

Ex. 7.

2. To find the greatest triangle inscribed in a circle
 $ACBD$.

Draw the diameter AB , and CD perpendicu-
 lar thereto; also draw AC , AD . Let $AB = d$,
 $AE = x$, $EC = y$: then triangle $ACD = xy = \text{max.}$
 or $xyy = \text{max.}$ but $yy = dx - xx$; therefore
 $dx^2 - x^2 = \text{max.} = dx^2 + 3dx^2e - x^2 - 4x^2e$ (putting
 $x + e$ for x), and $3dx^2e = 4x^2e$, or $4x = 3d$, whence
 $x = \frac{3}{4}d$.

SCHOLIUM.

When any quantity is a maximum or minimum,
 its root, or its square, or its cube, &c. will like-
 wise be a maximum or minimum. Also when any
 quantity is a maximum or minimum, any given
 quantity may be added to it, or subtracted from
 it, and it will still be a maximum, or minimum.
 Likewise it may be multiplied or divided by any
 given quantity, and still remain a maximum or
 minimum.

PROBLEM LXXXIV.

A number or quantity being given; to find its loga-
 rithms by a series, or to turn numbers into logarithms.

Let $\frac{x}{y}$ be the quantity given; $M = 1$, for
 Neper's logarithms, or $M = .434294482$, for the
 common

common logarithms. And let $x-y=v$, $x+y=z$.

Then the logarithm of $\frac{x}{y}$, will be denoted these several ways following, deduced from the nature of logarithms.

$$1. \text{ Log: } \frac{x}{y} = M \times \frac{v}{y} - \frac{v^2}{2y^2} + \frac{v^3}{3y^3} - \frac{v^4}{4y^4} + \frac{v^5}{5y^5} - \&c.$$

Or,

$$2. \text{ Log: } \frac{x}{y} = M \times \frac{v}{x} + \frac{v^2}{2x^2} + \frac{v^3}{3x^3} + \frac{v^4}{4x^4} + \frac{v^5}{5x^5} + \&c.$$

Or,

$$3. \text{ Log: } \frac{x}{y} = 2M \times \frac{v}{z} + \frac{v^3}{3z^3} + \frac{v^5}{5z^5} + \frac{v^7}{7z^7} + \&c.$$

Cor. 1. If v be far less than 1. Then

$$\text{Log: } \frac{1+v}{1-v} = M \times v - \frac{vv}{2} + \frac{v^3}{3} - \frac{v^4}{4} + \frac{v^5}{5} \&c.$$

This is plain by putting $y=1$. For then

$$x=1+v, \text{ and } \frac{x}{y} = 1+v.$$

$$\text{Cor. 2. Log: } y+v = \text{log: } y, + M \times \frac{v}{y} - \frac{v^2}{2y^2}$$

$$+ \frac{v^3}{3y^3} - \frac{v^4}{4y^4} \&c.$$

$$\text{or } \text{log: } y+v = \text{log: } y, + M \times \frac{v}{x} + \frac{v^2}{2x^2} + \frac{v^3}{3x^3} + \frac{v^4}{4x^4} \&c.$$

$$\text{or } \text{log: } y+v = \text{log: } y, + 2M \times \frac{v}{z} + \frac{v^3}{3z^3} + \frac{v^5}{5z^5} + \frac{v^7}{7z^7} \&c.$$

$$\text{For } \text{log: } x \text{ or } y+v = \text{log: } y \times \frac{x}{y} = \text{log: } y + \text{log: } \frac{x}{y}.$$

Cor. 3. If $l = \text{logarithm of } n$,
and $l+s = \text{logarithm of } n+v$. Then the additional part of the logarithm, that is,

$$s = M \times : \frac{v}{n} - \frac{v^2}{2n^2} + \frac{v^3}{3n^3} - \frac{v^4}{4n^4} \&c.$$

$$\text{or } s = M \times : \frac{v}{n+v} + \frac{v^2}{2 \cdot n+v} + \frac{v^3}{3 \cdot n+v} \&c.$$

$$\text{or } s = 2M \times : \frac{v}{2n+v} + \frac{v^3}{3 \cdot 3n+v} + \frac{v^5}{5 \cdot 2n+v} \&c.$$

For since $l+s = \text{log: } \overline{n+v}$, and $l = \text{log: } n$; therefore $l+s-l = \text{log: } \overline{n+v} - \text{log: } n = \text{log: } \frac{n+v}{n}$,

that is, $s = \text{log: } \frac{n+v}{n}$. And by this prop. (writing n

for y , $n+v$ for x , and $2n+v$ for z); s or $\text{log: } \frac{n+v}{n}$ will come out as above.

Cor. 4. If x be far less than a , then

$$\text{Log: } a+bx+cx^2+dx^3 \&c. = \text{log: } a + M \times : \frac{bx+cx^2+dx^3 \&c.}{a} - \frac{bx+cx^2 \&c.}{2aa}$$

$$+ \frac{bx \&c.}{3a^3} - \&c.:$$

$$\text{and log: } \frac{a}{a-bx-cx^2-dx^3 \&c.} = M \times :$$

$$\frac{bx+cx^2+dx^3 \&c.}{a} + \frac{bx+cx^2 \&c.}{2aa}$$

$$+ \frac{bx \&c.}{3a^3} \&c.:$$

and

$$\text{and log: } \frac{a+bx+cx^2+dx^3 \ \&c.}{a-bx-cxx-dx^2 \ \&c.} = 2M \times :$$

$$\frac{bx+cx^2+dx^3 \ \&c.}{a} + \frac{bx+cx^2 \ \&c.}{3a^3}$$

$$+ \frac{bx+\&c.}{5a^5} \ \&c.:$$

The first case appears from Case 1, Cor. 2. writing $a+bx+cx^2, \ \&c.$ for x , a for y , and $bx+cx^2 \ \&c.$ for v .

The second appears from Case 2. of this prop. writing a for x , $a-bx-cxx \ \&c.$ for y , and $bx+cx^2 \ \&c.$ for v .

The third appears from Case 3. of the prop. writing $a+bx+cx^2 \ \&c.$ for x , $a-bx-cxx \ \&c.$ for y , $2bx+2cx^2 \ \&c.$ for v , and $2a$ for z .

SCHOLIUM.

The $\log : y+v = \log : y : + \frac{2Mv}{2y+v}$ very near, when v is very small; which is only the first term of the series, Case 3. Cor. 2.

PROBLEM LXXXV.

A logarithm being given; to find the quantity belonging to it, or its number, by a series. Or to turn logarithms into numbers.

Let $l+s$ be the logarithm given, $n+v$ its number, and let l be the logarithm of the number n .

Put $m=2.302585093 = \frac{1}{M}$, for the common logarithms, or $m=1$, for *Neper's* logarithms. Then

by Cor. 3. last Prob. $s=M \times : \frac{v}{n} - \frac{v^2}{2n^2} + \frac{v^3}{3n^3} \ \&c.$

and $\frac{s}{M}$ or $ms = \frac{v}{n} - \frac{v^2}{2n^2} + \frac{v^3}{3n^3} \&c.$ Then by re-

version of series (Prob. lxii.), $\frac{v}{n} = ms + \frac{m^2 s^2}{2}$

$$+ \frac{m^3 s^3}{2 \cdot 3} + \frac{m^4 s^4}{2 \cdot 3 \cdot 4} \&c. \text{ Then}$$

$$1. v = n \times : ms + \frac{m^2 s^2}{2} + \frac{m^3 s^3}{2 \cdot 3} \&c. \text{ Whence}$$

$$2. n + v = n \times : 1 + ms + \frac{m^2 s^2}{2} + \frac{m^3 s^3}{2 \cdot 3} + \frac{m^4 s^4}{2 \cdot 3 \cdot 4} \&c. \text{ and}$$

$$3. \frac{n+v}{n} = 1 + ms + \frac{m^2 s^2}{2} + \frac{m^3 s^3}{2 \cdot 3} \&c.$$

That is,

$$\text{Number of } l+s = \text{number of } l \times : 1 + ms + \frac{m^2 s^2}{2} + \frac{m^3 s^3}{2 \cdot 3} + \frac{m^4 s^4}{2 \cdot 3 \cdot 4} \&c.$$

Cor. 1. If $n=1$, and $l=0$; then

$$1+v \text{ or number of } s = 1 + ms + \frac{m^2 s^2}{2} + \frac{m^3 s^3}{2 \cdot 3} + \frac{m^4 s^4}{2 \cdot 3 \cdot 4} \&c.$$

Cor. 2. If $l = \log: n$, and $l+s = \log: n+v$; then the additional part of the number, that is,

$$v = n \times : ms + \frac{m^2 s^2}{2} + \frac{m^3 s^3}{2 \cdot 3} + \frac{m^4 s^4}{2 \cdot 3 \cdot 4} \&c.$$

Cor. 3. If L be the log: of the number N , then

$$N^x = 1 + mxL + \frac{m^2 x^2 L^2}{2} + \frac{m^3 x^3 L^3}{2 \cdot 3} + \frac{m^4 x^4 L^4}{2 \cdot 3 \cdot 4} \&c.$$

For

Sect. VII: PROBLEMS: 249

For, by Cor. 1. $1+v$ (numb. of s log.) =
 $1+ms + \frac{m^2s}{2} + \frac{m^3s}{2.3}$ &c. Where $1+v$ may
 represent any number, and s its logarithm. There-
 fore let $1+v=N$, and $s=L$; then

$$N \text{ (numb. of } L \text{ log.)} = 1+mL + \frac{m^2L^2}{2} + \frac{m^3L^3}{2.3}$$

&c. therefore by the nature of logarithms,

$$N^x \text{ (numb. of } xL \text{ log.)} = 1+mxL + \frac{m^2x^2L^2}{2}$$

$$+ \frac{m^3x^3L^3}{2.3} \text{ \&c.}$$

Cor. 4. If $y=n+v$, $x=r+e$, $l=log: n$. Then

$$y^x \text{ or } \overline{n+v}^{r+e} = n^r \times \left(1 + mel + \frac{mel^2}{2} + \frac{mel^3}{2.3} \right)$$

$$\text{\&c.} \times \left(1 + \frac{r+e}{1} \times \frac{v}{n} + \frac{r+e}{1} \times \frac{r+e-1}{2} \times \frac{vv}{nn} \right. \\ \left. + \frac{r+e}{1} \times \frac{r+e-1}{2} \times \frac{r+e-2}{3} \times \frac{v^3}{n^3} \right) \text{\&c.}$$

$$\text{For } \overline{n+v}^{r+e} = n^{r+e} \times \overline{1+\frac{v}{n}}^{r+e} = n^r \times n^e \times$$

$$\overline{1+\frac{v}{n}}^{r+e}, \text{ but by Cor. 3. } n^e = 1 + mel +$$

$$\frac{mel^2}{2} + \frac{mel^3}{2.3} \text{ \&c. and } \overline{1+\frac{v}{n}}^{r+e} = 1 + \frac{r+e}{1} \times \frac{v}{n}$$

$$+ \frac{r+e}{1} \times \frac{r+e-1}{2} \times \frac{vv}{nn} \text{ \&c.}$$

Cor. 5. If $l=log: n$; then

$$N^{r+e} = n^r \times \left(1 + mel + \frac{mel^2}{2} + \frac{mel^3}{2.3} \right) \text{ \&c.}$$

For here $v=0$.

Cor.

Cor. 6. If v, e be exceeding small, then

$\frac{r}{n+v}^{r+e} = n^r \times \left(1 + me + \frac{r^2 v}{n} \right)$; nearly, being only by the first power of e and v .

Cor. 7. If n = number of the logarithm a ; then the number of the logarithm $a + bx + cxx + dx^3$ &c.

$= n \times$ into $1 + m \times \frac{bx + cx^2 + dx^3}{a}$ &c. $+ \frac{m^2}{2} \times \frac{bx + cxx}{a^2}$ &c. $+ \frac{m^3}{2 \cdot 3} \times \frac{bx + cxx}{a^3}$ &c. $+ \frac{m^4}{2 \cdot 3 \cdot 4} \times \frac{bx + \&c.}{a^4}$ $+ \&c.$

This follows from this problem, putting $l = a$, and $s = bx + cxx$ &c.

PROBLEM LXXXVI.

A problem being resolved analytically, to demonstrate it synthetically.

R U L E.

When a problem has been solved algebraically, the demonstration of it is to be deduced from the steps of the algebraic process; by going backward from the end of it to the beginning; observing how each step is formed from the foregoing, and forming your process accordingly.

S E C T. VIII.

*The Resolution of Equations ; and the extraction
of their roots in numbers.*

P R O B L E M LXXXVII.

To find the limits of the roots of an equation.

WHEN an equation is proposed to have its root extracted, it is proper to find the limits of the roots ; lest we lose our time in seeking the roots beyond these limits.

R U L E.

Reduce the equation, that the highest term may have 1 for its coefficient ; then square the coefficient of the second term, from which subtract twice the coefficient of the third term, then the square root thereof is greater than the greatest root of the equation. But the equation should be clear of impossible roots.

For that quantity is the sum of the squares of the roots, by Prob. xl. Art. 9. and that sum, is greater than the square of any one root.

Or thus,

Substitute several numbers successively for the unknown quantity ; till at last you find two numbers which give, one a positive, and the other a negative result. Then the root is between these numbers.

There are other rules among the writers of Algebra, which come nearer ; but then they are more laborious.

Ex.

Ex. 1.

$$\text{Let } x^3 + 3x^2 - 5x - 20 = 0.$$

$$\text{Then } 3 \times 3 - 2 \times -5 = 9 + 10 = 19.$$

and $\sqrt{19} = 4.3$, &c. Therefore 4.3 is greater than any of the roots.

Ex. 2.

$$\text{Suppose } xx - x - 5 = 0.$$

$$\text{If } x = 2, \text{ then the result is } 2 - 5 = -3.$$

$$\text{If } x = 3, \text{ the result is } 6 - 5 = +1.$$

Therefore the root is between 2 and -3.

P R O B L E M LXXXVIII.

To resolve a quadratic equation, and extract its root in numbers.

I comprehend all equations under the name of quadratics, in which are two terms involving the unknown quantity; and where the index of one is double to that of the other. As in these,

$$aa + ba = d$$

$$a^4 + ba^2 = d$$

$$a^6 + ba^3 = d, \text{ \&c.}$$

where b, d , may represent any numbers, affirmative or negative.

Every quadratic equation has two roots, though perhaps only one of them will answer the question proposed. And to find these roots the equation proposed must be first reduced, by dividing all, by the coefficient of the highest term; and then transposing the known quantity to the contrary side. Which done, the equation will appear thus, $aa + ba = d$. Now add to both sides $\frac{1}{4}bb$ the square of half the coefficient of a , and we have $aa + ba + \frac{1}{4}bb = \frac{1}{4}bb + d$, where the first side is a complet

complete square; therefore extract the square root, and $a + \frac{1}{2}b = \pm \sqrt{\frac{1}{4}bb + d}$, transpose $\frac{1}{2}b$, then $a = -\frac{1}{2}b + \sqrt{\frac{1}{4}bb + d}$. So a becomes known, being either equal to $-\frac{1}{2}b + \sqrt{\frac{1}{4}bb + d}$, or to $-\frac{1}{2}b - \sqrt{\frac{1}{4}bb + d}$. Whence this

R U L E.

The equation being cleared, complete the square by adding to both sides the square of half the coefficient of the second term. Then extract the root of both sides, which may be either $+$ or $-$; then transpose the known quantity.

Note. If the absolute number is negative, and greater than $\frac{1}{4}$ the square of the coefficient; the equation is impossible.

If $aa + ba = d$,

Then $a = \pm \sqrt{\frac{1}{4}bb + d} - \frac{1}{2}b$.

And the root extracted in numbers gives a ; but if $\frac{1}{4}bb$ is lesser than d , and d negative; it is impossible.

Ex. 1.

If $aa + 5a = 68$.

Then $a = \pm \sqrt{68 + 6\frac{1}{4}} - \frac{1}{2} = \pm \sqrt{74.25} - 2.5$;

74.25 (8.6168 &c.
64

166) 1025
+ 6) 996

+ 8.6168
- 2.5

1721) 2900
+ 1) 1721

+ 6.1168 = a

- 11.1168 = a

17226) 117900
+ 6) 103356

17232) 1454400

Ex.

Ex. 2.

Let $aa - 6a = 27$.

Then $a = 3 \pm \sqrt{9 + 27} = 3 \pm \sqrt{36}$.
 that is, $a = 3 + 6 = 9$.
 or $a = 3 - 6 = -3$.

Ex. 3.

Suppose $aa - 236a = -1155$.

Then $a = 118 \pm \sqrt{118^2 - 1155}$;
 that is, $a = 118 + 113 = 231$
 or $a = 118 - 113 = 5$.

2 R U L E.

When you have large numbers to deal with; it is better to proceed thus. Clear the equation,

And if $aa + ba = d$,then $a = \frac{d}{b+a}$, the form.

To find the first quotient figure, take $\frac{d}{b}$, when b is far greater than a ; or take \sqrt{d} , when a is far greater than b ; or take $\frac{d}{2b}$ when a and b are nearly equal; thus it will easily be found by a few trials. Or in general, take the first figure such, that when it is multiplied by the sum of itself and b , it will produce the first figure or figures of d , or the next less: this is all the difficulty. Then multiply and subtract as usual, the remainder is the resolvend.

Then to continue the division; you must find a new divisor for each quotient figure, thus. Add the last quotient figure to the last divisor (duly observing their places), for a new divisor; see how

how oft this is contained in the resolvend, set the answer in the quotient, and also add it to the divisor; then multiply the whole divisor by that quotient figure; and subtract the product, for a new resolvend. But when any of the signs are negative, the proper quantities are to be subtracted, instead of being added. This work is always to be repeated for each quotient figure.

When any quotient figure is so great that the product exceeds the resolvend, place a less figure in the quotient.

When you have got more than half your intended number of figures in the quotient, you may continue the division without adding the new quotient figures to the divisor.

Observe, each quotient figure is to be added twice to the divisor; once before multiplication, and once after; just as in extracting the square root, and for the same reason. For this method extracts the square root, when $b=0$.

When one root is had, the other is found, by adding this to the coefficient b ; for the sum, changing its sign, is the other root.

This rule is the foundation of the method for extracting the roots of affected equations.

Ex. 4.

$$\text{Let } aa + 32a = 4644.$$

$$\text{then } \quad \quad \quad 4644$$

$$a = \frac{\quad}{32+a}$$

Suppose $\frac{4600}{32} = 100$ too great for a .

$\sqrt{4644} = 60$, which is also too great for a . Take

$a = \frac{4600}{64} = 7$, too great. Take $a = 50$.

$$\begin{array}{r}
 32 \\
 + 50 \\
 \hline
 82) 4644 (50 \\
 + 54 \quad 410 \\
 \hline
 136 \quad 544 (4 \\
 \quad 544 \quad \hline
 \quad \hline
 \quad \quad | 54 = a.
 \end{array}$$

Ex. 5.

$$\begin{array}{r}
 \text{Let } aa + 35a = 28349994 \\
 \quad \quad \quad 28349994 \\
 a = \frac{\quad \quad \quad}{35 + a}
 \end{array}$$

Here $a = \sqrt{28}$ &c. = 5000 nearly.

$$\begin{array}{r}
 +35 \\
 5000 \\
 \hline
 5035) 28349994 (5307 = a \\
 5300) 25175 \dots \\
 \hline
 10335) 31749 \\
 307) 31005 \\
 \hline
 10642) 74494 \\
 \quad 74494 \\
 \quad \hline
 \quad \quad \quad
 \end{array}$$

Ex. 6.

$$\text{Suppose } aa - 5307a = -184520.$$

$$\text{then } a = \frac{-184520}{-5307 + a} = \frac{184520}{5307 - a}.$$

$$\text{Here } a = \frac{184}{5} = 3 \text{ nearly.}$$

Sec. VIII. EQUATIONS.

$$\begin{array}{r} 5307) 184520 \quad (35 = a. \\ -30 \quad 15831 \end{array}$$

$$\begin{array}{r} 5277) 26210 \\ -35 \quad 26210 \end{array}$$

$$\begin{array}{r} 5242) \dots \end{array}$$

Ex. 7.

Let $aa + 463a = 26698$

$$a = \frac{26698}{463 + a}$$

$$\begin{array}{r} 463) 26698 \quad (51.855342 = a. \\ +50 \quad 2565 \end{array}$$

$$\begin{array}{r} 513) 1048 \\ +51 \quad 564 \end{array}$$

$$\begin{array}{r} 564) 4800000 \\ +1.8 \quad 45264 \end{array}$$

$$\begin{array}{r} 565.8) 313000 \\ +.85 \quad 283325 \end{array}$$

$$\begin{array}{r} 566.65) 30275 \\ +5 \quad 28335 \end{array}$$

$$\begin{array}{r} 566.70 \quad 1940 \\ \quad \quad 1700 \end{array}$$

$$\begin{array}{r} \quad \quad \quad 240 \\ \quad \quad \quad 226 \\ \quad \quad \quad 14 \\ \quad \quad \quad 11 \\ \quad \quad \quad 3 \end{array}$$

$$\begin{array}{r} 566.70 \quad 1940 \\ \quad \quad 1700 \end{array}$$

$$\begin{array}{r} \quad \quad \quad 240 \\ \quad \quad \quad 226 \\ \quad \quad \quad 14 \\ \quad \quad \quad 11 \\ \quad \quad \quad 3 \end{array}$$

$$\begin{array}{r} 566.70 \quad 1940 \\ \quad \quad 1700 \end{array}$$

$$\begin{array}{r} \quad \quad \quad 240 \\ \quad \quad \quad 226 \\ \quad \quad \quad 14 \\ \quad \quad \quad 11 \\ \quad \quad \quad 3 \end{array}$$

Scholium. If $x^2 + bx^2 = d$. Put $a = xx$, then $aa + ba = d$; and find a as above. Then $x = \sqrt{a}$,
S by

258 RESOLUTION of B. I
 by extracting the root. And the same for higher equations.

To prove the truth of this rule. Let $x+y+z$ &c. be the true value of a ; x the first figure, y the second, and z the third, &c. Then since $aa+ba=d$, the value of d will be

$$b \times \overbrace{x+y+z}^2 + \overbrace{x+y+z}^2, \text{ whence } x+y+z \text{ \&c. or}$$

$$a = \frac{b+a}{d} = \frac{bx+by+bz+xx+x+y+z}{b+x+y+z}$$

The operation.

b)	$bx+by+bz+xx+2xy(x+y+z \text{ \&c.}$	
$+x$)	$+yy+2xz$	
		$+zz+2yz$	
1	divisor $b+x$)	$bx+xx$	
		$+x+y$	
2	divif. $b+2x+y$)	$by+bz+yy+2xy$ resolvend	
		$+zz+2xz$	
		$+2yz$	
		$by+2xy+yy$	
3	divif. $b+2x+2y+z$)	$bz+zz+2xz$ resolvend	
		$+2yz$	
		$bz+2xz+2yz+zz$	
		0	

Here $bx+by$ &c. being divided by b gives x in the quotient, and x added to b , gives $b+x$ for the divisor, and $bx+xx$ for the product, which subtracted, leaves the resolvend $by+bz+2xy+yy$ &c.

Then in order to get the second figure y , the resolvend $by+2xy+yy$ &c. is to be divided by $b+2x+y$. Therefore $x+y$ is to be added to the last divisor $b+x$, to get the new divisor $b+2x+y$. This divisor multiplied by y , gives $by+2xy+yy$, which

which subtracted, leaves $bz + 2xz + 2yz + zz$ for the resolvend.

Then to get the third figure z , it is plain, the resolvend $bz + 2xz + 2yz + zz$ must be divided by the divisor $b + 2x + 2y + z$, but this new divisor is $b + 2x + y + x + z$, that is, it is the old divisor with $x + z$ added. Then this divisor multiplied by z , and subtracted, 0 remains. Therefore the root is rightly extracted, and the rule true.

As I am upon this subject, I shall also shew the truth of the rule for extracting the square root in Arithmetic, which is the case here, when $b = 0$.

Let $x + y + z$ be the square, that is,

1 div. x) $xx + 2xy + yy + 2xz + 2yz + zz$
 $+ x$) xx

$2x$) $2xy + yy + 2xz + 2yz + zz$
 $+ y$ $2xy + yy$

2 div. $2x + y$) $+ 2xz + 2yz + zz$
 $+ y + z$ $+ 2xz + 2yz + zz$

3 div. $2x + 2y + z$) 0

Here x being the root of the first term, its square subtracted, leaves the resolvend $2xy + yy$ &c. Then to find y , the resolvend must be divided by $2x + y$. That is, to the old divisor x , add $x + y$ for a new divisor $2x + y$; this multiplied by y , and subtracted, leaves the resolvend $2xz + 2yz + zz$. Again to find z , the resolvend is to be divided by $2x + 2y + z$ that 0 remain; that is, to the old divisor $2x + y$ add $y + z$, the sum is the new divisor $2x + 2y + z$, which multiplied by z , is equal to the resolvend, so that 0 remains; and the root is $x + y + z$.

PROBLEM LXXXIX.

To extract the root of a cubic equation.

R U L E.

Take away the second term of the equation, (by Prob. li.) which then will be in this form,

$$a^3 + ba = d.$$

Then substitute numbers in either of the following forms, and extract the roots, by which means a will be found.

$$a = \sqrt[3]{\frac{1}{2}d + \sqrt{\frac{1}{4}dd + \frac{1}{27}b^3}} - \frac{\sqrt[3]{b}}{\sqrt[3]{\frac{1}{2}d + \sqrt{\frac{1}{4}dd + \frac{1}{27}b^3}}}.$$

$$\text{or } a = \sqrt[3]{\frac{1}{2}d + \sqrt{\frac{1}{4}dd + \frac{1}{27}b^3}} + \sqrt[3]{\frac{1}{2}d - \sqrt{\frac{1}{4}dd + \frac{1}{27}b^3}}.$$

Note, When b is negative, and $\frac{1}{27}b^3$ greater than $\frac{1}{4}dd$, the equation is impossible.

Ex. I.

$$\text{Let } x^3 - 6x = -9.$$

$$\text{Here } b = -6, d = -9, \text{ and } \sqrt[3]{\frac{1}{4}dd + \frac{1}{27}b^3} = \sqrt[3]{12\frac{1}{4}} = 3\frac{1}{2}.$$

$$\text{and } \sqrt[3]{-\frac{1}{2}d - \sqrt{\frac{1}{4}dd + \frac{1}{27}b^3}} = \sqrt[3]{-1} = -1. \text{ Therefore}$$

$$a = -1 - \frac{-2}{-1} = -1 - 2 = -3.$$

$$\text{or } \sqrt[3]{-\frac{1}{2}d + \sqrt{\frac{1}{4}dd + \frac{1}{27}b^3}} = \sqrt[3]{-8} = -2, \text{ whence}$$

$$a = -1 - 2 = -3, \text{ as before.}$$

Ex.

Ex. 2.

Let $a^3 + 6a = 20$.

Here $b=6$, $d=20$, and $\sqrt[3]{dd + \frac{1}{27}b^3} = \sqrt[3]{108}$.

And $\sqrt[3]{10 + \sqrt{108}} = 1 + \sqrt{3}$ (Prob. lxxii.) and

$\sqrt[3]{10 - \sqrt{108}} = 1 - \sqrt{3}$. Whence

$a = 1 + \sqrt{3} + 1 - \sqrt{3} = 2$.

Ex. 3.

Let $a^3 - 15a = 4$.

Here $b=-15$, $d=4$, and $\sqrt[3]{dd + \frac{1}{27}b^3} = \sqrt[3]{-121} = 11\sqrt{-1}$.

And $\sqrt[3]{2 + 11\sqrt{-1}} = 2 + \sqrt{-1}$. And

$\sqrt[3]{2 - 11\sqrt{-1}} = 2 - \sqrt{-1}$.

Whence $a = 2 + \sqrt{-1} + 2 - \sqrt{-1} = 4$.

Ex. 4.

Suppose $a^3 + 24a = 587914$.

Here $b=24$, $d=587914$. $\sqrt[3]{dd + \frac{1}{27}b^3} = 293957.000878$.

And $\sqrt[3]{\frac{1}{27}d + 29} \&c. = 83.7731$. And

$\frac{8}{83.77} = .0958$; therefore

$a = 83.7731 - .0958 = 83.6773$.

SCHOLIUM.

It sometimes happens that the root may be found, though the negative quantity $\frac{1}{27}b^3$ be greater than $\frac{1}{27}dd$; and that is when the surd cubic root can be extracted. For then the irrational parts, in different parts of the equation, will destroy one another, and vanish; as in Ex. 3.

To prove the truth of this rule. Put

$$r = \sqrt{\frac{1}{2}dd + \frac{1}{27}b^3}, \quad s = \sqrt[3]{\frac{1}{2}d+r}. \quad \text{Then } a = s - \frac{b}{3s}, \text{ and } a^3 = s^3 - bs + \frac{bb}{3s} - \frac{b^3}{27s^3}, \text{ and } ba = bs - \frac{bb}{3s},$$

$$\text{therefore } a^3 + ba = s^3 - \frac{b^3}{27s^3} = \frac{1}{2}d + r - \frac{\frac{1}{27}b^3}{\frac{1}{2}d+r}$$

$$= \frac{\frac{1}{2}dd + dr + rr - \frac{1}{27}b^3}{\frac{1}{2}d+r} = (\text{restoring } rr)$$

$$\frac{\frac{1}{2}dd + dr + \frac{1}{2}d\frac{1}{2}d + \frac{1}{27}b^3 - \frac{1}{27}b^3}{\frac{1}{2}d+r} = \frac{\frac{1}{2}dd + dr}{\frac{1}{2}d+r} = d; \text{ that}$$

is, $a^3 + ba = d$, according to the first part of the rule.

And the second part is proved, by shewing that

$$-\frac{b}{3s} = \sqrt[3]{\frac{1}{2}d-r}. \quad \text{It is plain } \frac{1}{2}d+r \times \frac{1}{2}d-r$$

$$= \frac{1}{2}dd - rr = -\frac{1}{27}b^3, \text{ therefore } \frac{1}{2}d-r = \frac{-\frac{1}{27}b^3}{\frac{1}{2}d+r} =$$

$$-\frac{b^3}{27s^3}, \text{ and } \sqrt[3]{\frac{1}{2}d-r} = -\frac{b}{3s}. \quad \text{Which was}$$

to be proved.

Some of the cases of cubic equations may also be resolved trigonometrically by the table of sines. As suppose the equation $x^3 - px = +q$, to be given. By Prop. 24, 25. Trigonometry, if $r = \text{radius}$,

$y = \text{fine of an arch}$, then $3y - \frac{4y^3}{rr} = S. 3ce$

the arch. And by Prop. 26. if $x = \text{cosine of an arch}$, then $\frac{4x^3}{rr} - \frac{3}{4}x = \text{cosine of } 3ce \text{ that arch.}$

These equations reduced give $y^3 - \frac{3}{4}rry = -\frac{rr}{4}$
 $\times \text{fine of } 3ce \text{ the arch. And } x^3 - \frac{3}{4}rrx = +\frac{rr}{4}$
 $\times \text{cosine of thrice the arch. Or putting } y \text{ for either the sine or cosine of the arch, } C \text{ for the sine or cosine of thrice the arch; then}$

$$y^3 -$$

$y^3 - \frac{1}{3} r r y = \pm \frac{r r}{4} C$, the sign + being for cosines, and - for sines.

Then, if the given equation $x^3 - p x = \pm q$ is to be resolved; it must be compared with the foregoing, and all the parts made similar in both. Therefore let the equation $x^3 - p x = \pm q$, be denoted thus, $x^3 - \frac{1}{3} R R x = \pm \frac{1}{3} R R S$, S being the sine or cosine of thrice the arch. Therefore $\frac{1}{3} R R = p$, and $R = \sqrt{\frac{1}{3} p}$. Also $q = \frac{1}{3} R R S = \frac{1}{3} p S$, and $S = \frac{3q}{p}$. Whence by proportion $R (\sqrt{\frac{1}{3} p}) : S (\frac{3q}{p})$

$:: r : C = \frac{3r q}{p \sqrt{\frac{1}{3} p}}$, the cosine or sine of an arch.

Of which, y is the cosine or sine of the third part.

Then y being found, it will be $r : y :: R (\sqrt{\frac{1}{3} p})$

$: x = \frac{y \sqrt{\frac{1}{3} p}}{r}$, as required. Hence this

2 R U L E.

Take away the second term (by Prob. li.) if it have any; and the equation will be reduced to this form,

$$x^3 - p x = \pm q.$$

Then take $\frac{3r q}{p \sqrt{\frac{1}{3} p}} =$ the cosine of an arch (if it be + q), or the sine (if - q). Find $y =$ cosine or sine of $\frac{1}{3}$ that arch; then $\frac{y \sqrt{\frac{1}{3} p}}{r} = x$ required.

And this last arch may be either that we found, or, that +120°, or the same +240°. By which means you will have three roots or values of y .

But note, when $\frac{3r q}{p \sqrt{\frac{1}{3} p}}$ is greater than 1, the question is impossible by this rule.

Therefore this rule supplies the defect of the first rule, which only solves equations that have but one root real, and two impossible ones: whilst this rule solves such as have three roots real.

Ex. 5.

Let $x^3 - 91x = -330$.

Here $x = a$, $p = 91$, $q = 330$, and $\sqrt[3]{p} = 11.015$, and if $r = 1$, $\frac{3rq}{p\sqrt[3]{p}} = .987655 =$ sine of 81° very near; and the third part is 27, or 147, or 267; whose sines are, $y = .45399$, or $.54467$, or $-.99863$; these multiplied by 11.015 produce 5.0004, and 5.9991, and -10.9998 ; therefore the three roots are 5, 6, and -11 .

Ex. 6.

Suppose $x^3 - 19x = 30$.

Here $p = 19$, $q = 30$, and $\sqrt[3]{p} = 5.03323$ and $\frac{3rq}{p\sqrt[3]{p}} = .94112 =$ cosine of $19^\circ : 45'$, and the third part is $6^\circ 35'$, or $126^\circ 35'$, or $246^\circ 35'$, whose cosine is $y = .99340$, or $-.59599$, or $-.39741$; which multiplied by 5.03323, produce 4.99998, and -2.99974 , and -2.00024 . So the three roots are these, 5, -3 , and -2 .

PROBLEM XC.

To resolve a biquadratic equation, by dissolving it into two quadratics.

Take away the second term (by Prob. li), and let the resulting equation be $x^4 + qx^2 + rx + s = 0$. Suppose it to be generated by the two quadratics, $xx + ex + f = 0$, and $xx - ex + g = 0$. These being multiplied

multiplied together produce $x^4 + fx^3 + egx^2 + fg = 0$.

$$\begin{array}{r} +g \\ -ef \\ -ee \end{array}$$

Comparing the terms of this with the first equation, we have $f+g-ee=q$, $eg-ef=r$, and $fg=s$; whence $g+f=q+ee$, and $g-f=\frac{r}{e}$; and conse-

quently $g = \frac{q+ee+\frac{r}{e}}{2}$, and $f = \frac{q+ee-\frac{r}{e}}{2}$. And

$(fg =) \frac{qq+2qee+e^4-\frac{rr}{ee}}{4} = s$. And by reduction,
 $e^6 + 2qe^4 + qqee - rr = 0$. Put $y = ee$, and then

$y^3 + 2qy^2 + qqy - rr = 0$. A cubic equation; whence the following

R U L E.

To resolve the biquadratic equation $x^4 + qx^2 + rx + s = 0$. Take the cubic equation $y^3 + 2qy^2 + qqy - rr = 0$; out of which take away the second term (by Prob. li.); and find the root by the last problem, or otherwise; and from thence find y . Then

take $e = \sqrt{y}$, and $f = \frac{q+ee-\frac{r}{e}}{2}$, and

$$g = \frac{q+ee+\frac{r}{e}}{2}$$

Lastly, find the roots of these two quadratic equations, $xx+ex+f=0$, and $xx-ex+g=0$. And these will be the four roots, of the biquadratic $x^4+qx^2+rx+s=0$.

Example.

Example.

$$\text{Let } x^4 - 25x^2 + 60x - 36 = 0.$$

From this you have the cubic equation $y^3 - 50y^2 + 769y - 3600 = 0$. Take away the second term, by writing $v + \frac{50}{3}$ for y . And we have

$$v^3 - \frac{193}{3}v = \frac{1150}{27}. \text{ And by Rule 2. Prob.}$$

last, $v = 8.3333 \text{ \&c.} = 8\frac{1}{3}$, whence $y = 8\frac{1}{3}$

$+ \frac{50}{3} = 25$, and $e = 5$; therefore $f =$

$$\frac{-25 + 25 - \frac{60}{5}}{2} = -6, \text{ and } g = \frac{-25 + 25 + \frac{60}{5}}{2} =$$

$+6$. Whence $xx + 5x - 6 = 0$, and $xx - 5x + 6 = 0$; and the roots of the former equation are 1 and -6 ; and of the latter, 3 and 2. Therefore the four roots of the biquadratic, $x^4 - 25x^2 + 60x - 36 = 0$, are 1, 2, 3, and -6 .

And the same roots will be found, by making use of the other values of v , which are $-\frac{2}{3}$, and $-\frac{23}{3}$.

Schol. But this and such like rules are of little value; for there is far more labour here in getting the roots than by the method of converging series, which is to follow.

P R O B L E M XCI.

To extract the root of any pure power in numbers.

Let G be the number given to be extracted; m the root required, r the nearest root, and e the

the remaining part of it; then $\overline{r+e^m} = G$, that is (Cor. 1. Prob. v.) $r^m + mr^{m-1}e + m \cdot \frac{m-1}{2}$

$r^{m-2}ee + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} r^{m-3}e^3 \&c. = G$, and rejecting e^3 and the higher powers, as very small;

we have $mr^{m-1}e + m \cdot \frac{m-1}{2} r^{m-2}ee = G - r^m$,

and $\frac{mr^{m-1}}{m \cdot \frac{m-1}{2} r^{m-2}} e + ee = \frac{G - r^m}{m \cdot \frac{m-1}{2} r^{m-2}}$. Hence

this

R U L E.

Let $G =$ absolute number.

$r =$ the nearest root you can find.

$r+e =$ the true root.

$m =$ the index of the root.

$$b = \frac{2r}{m-1}$$

$$D = \frac{G - r^m}{m \cdot \frac{m-1}{2} r^{m-2}}$$

Then $be + ee = D$, or $e = \frac{D}{b+e}$, nearly.

Which equation is to be resolved by Prob. lxxxviii. When e is had, then $r+e$ is to be taken for a new value of r , and the operation repeated, perhaps oftener than once. This rule generally triples the number of figures.

But if the third power of e be taken in, then

$$mr^{m-1}e + m \cdot \frac{m-1}{2} r^{m-2}ee + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} r^{m-3}e^3 = G - r^m$$

and since $be + ee = D$, therefore

fore $ce = D - be$, and $e^3 = De - bee$, and $m \cdot \frac{m-1}{2}$.

$$\frac{m-2}{3} r^{m-3} e^3 = m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} r^{m-3} \times \overline{De - bee},$$

whence $mr^{m-1} e + m \cdot \frac{m-1}{2} r^{m-2} ce + m \cdot \frac{m-1}{2}$.

$$\frac{m-2}{3} r^{m-3} \times \overline{De - bee} = G - r^m, \text{ and } rre + \frac{m-1}{2}$$

$$\frac{m-2}{3} De + \frac{m-1}{2} rce - \frac{m-1}{2} \cdot \frac{m-2}{3} bee = \frac{G - r^m}{mr^{m-3}} = rF,$$

by substitution, (putting $F = \frac{G - r^m}{mr^{m-2}}$); expunge b and

$$D, \text{ then } rre + \frac{m-1}{2} \cdot \frac{m-2}{3} \times \frac{G - r^m}{m \cdot \frac{m-1}{2} r^{m-2}} e +$$

$$\frac{m-1}{2} rce - \frac{m-1}{2} \cdot \frac{m-2}{3} \times \frac{2r}{m-1} ce = rF; \text{ that is,}$$

$$rre + \frac{m-2}{3} \times Fe + \frac{m-1}{2} rce - \frac{m-2}{3} rce = rF,$$

$$\text{that is, } rr + \frac{m-2}{3} F \times e + \frac{m+1}{6} rce = rF,$$

whence this

2 R U L E.

Let $G =$ absolute number.

$r =$ nearest root you can find.

$r + e =$ true root.

$m =$ index of the root.

$$F = \frac{G - r^m}{mr^{m-2}}$$

$$\text{Then } \frac{6r + \frac{2m-4}{r} \cdot F}{m+1} e + ee = \frac{6F}{m+1}, \text{ nearly;}$$

Which

Which is to be solved as Prob. lxxxviii, and repeated with new r , if there be occasion: This rule commonly quintuples the number of figures in the root, true; at each operation.

The root of any number may also be extracted by Prob. lviii. after this manner.

3 R U L E.

Let $P + Pq$, be the number given to be extracted.

P , the greatest power contained in it.

Pq , the remainder; and

q , the quotient, arising by dividing the remainder by the greatest power.

n , the index of the root. Then

$$\sqrt[n]{r + rq} = P^{\frac{1}{n}} + \frac{1}{n} Aq - \frac{n-1}{2n} Bq - \frac{2n-1}{3n} Cq - \frac{3n-1}{4n} Dq \text{ \&c.}$$

Where $A, B, C, \text{ \&c.}$ are the preceding terms. In this rule, when two or three figures are got, put them equal to

$P^{\frac{1}{n}}$, and begin the operation anew; and the series will then converge exceeding fast; and so much faster as q is less.

Cor. Hence it follows, that

$$\sqrt{P + Pq} = \sqrt{P} + \frac{1}{2} Aq - \frac{1}{4} Bq - \frac{3}{6} Cq - \frac{5}{8} Dq - \frac{7}{10} Eq \text{ \&c. for the square root.}$$

$$\sqrt[3]{P + Pq} = \sqrt[3]{P} + \frac{1}{3} Aq - \frac{2}{6} Bq - \frac{5}{9} Cq - \frac{8}{12} Dq - \frac{11}{15} Eq \text{ \&c. for the cube root.}$$

$$\sqrt[4]{P + Pq} = \sqrt[4]{P} + \frac{1}{4} Aq - \frac{3}{8} Bq - \frac{7}{12} Cq - \frac{11}{16} Dq \text{ \&c. for the biquadrate root.}$$

$$\sqrt[5]{P+Pq} = \sqrt[5]{P} + \frac{1}{5} Aq - \frac{4}{10} Bq - \frac{9}{15} Cq - \frac{14}{20} Dq \text{ \&c. for the fifth root.}$$

$$\sqrt[3]{P+Pq} = \sqrt[3]{P} + \frac{2}{3} Aq - \frac{1}{6} Bq - \frac{4}{9} Cq - \frac{7}{12} Dq - \text{\&c. for the cube root of the square; and so on.}$$

Ex. 1.

What is the cube root of 2.

Here $G=2$, $r=1$, $m=3$, by Rule 1, $b=1$.

$$D = \frac{1}{3} = .3333; \text{ and } e = \frac{D}{b+e}$$

$$\begin{array}{r} 1. \\ +2 \end{array}) .3333 \quad (.26 = e$$

$$\begin{array}{r} 1.2 \\ +26 \end{array}) .0933 \quad \text{and } r+e = 1.26$$

$$\begin{array}{r} 1.46 \\ \quad 57 \end{array}$$

Again, for a second operation;

Let new $r=1.26$; then $G-r^3 = .000376$,

$$\text{and } m \cdot \frac{m-1}{2} r = 3r = 3.78, \text{ and } D = \frac{.000376}{3.78} =$$

$.000099471$, and because e is negative here,

$$e = \frac{.000099471}{1.26 - e}$$

1.2600000

$$\begin{array}{r}
 1.2600000).0000994710(-.000078950106 \\
 \underline{-7} \qquad \qquad \qquad 881951 \\
 1.2599300 \qquad \qquad \qquad 11275900 \\
 \underline{-78} \qquad \qquad \qquad 10078816 \\
 1.2598520 \qquad \qquad \qquad 11970840 \\
 \underline{-89} \qquad \qquad \qquad 11338587 \\
 1.259843 \qquad \qquad \qquad 632253 \\
 \dots \qquad \qquad \qquad 629921 \\
 \qquad \qquad \qquad \underline{2332} \\
 \qquad \qquad \qquad 1259 \\
 \qquad \qquad \qquad \underline{73}
 \end{array}$$

$$\begin{aligned}
 r &= 1.26000 \\
 e &= -.000078950106 \\
 \sqrt[3]{2} &= 1.259921049894
 \end{aligned}$$

Ex. 2.

Extract the 5th root of 2327834559873.

First point every fifth figure thus

$$2327834559873.$$

Then for brevity's sake, take only the first period, as an integer, that is 232. Then proceeding by Rule 2, we shall find 2, the root of the greatest power contained therein; and thence, $r=2$, and

$$232 = G$$

$$32 = r^m$$

$$100 = G - r^m$$

and

$$40 = mr^{m-2}$$

Whence

$$\frac{100}{40} = 5 = F.$$

Therefore

Therefore $4\frac{1}{2}e + ee = 5$, or $e = \frac{5}{4.5 + e}$.

$$\begin{array}{r} 4.5) 5.00 (-92 = e. \\ +9 \ 4 \ 86 \end{array}$$

$$\begin{array}{r} 5.4) 1400 \\ +92 \ 1264 \end{array}$$

$$\begin{array}{r} 6.32) 136 \end{array}$$

Suppose again new $r = 290$.

Then $r^5 = 24389000$.

$$G = 2327834559873$$

$$r^5 = 2051114900000$$

$$G - r^5 = 276719659873$$

$$F = 2269.217 \quad \text{whence}$$

$$297.825e + ee = 2269.217$$

$$\text{or } e = \frac{2269.217}{297.825 + e}$$

$$\begin{array}{r} 297.825) 2269.217 (7.43375 \\ +7 \end{array}$$

$$\begin{array}{r} 304.825) 1354420 \\ +7.4 \end{array}$$

$$\begin{array}{r} 312.225) 105520 \\ +43 \end{array}$$

$$\begin{array}{r} 312.655) 11724 \\ +33 \end{array}$$

$$\begin{array}{r} 312.68) 2344 \\ \dots \end{array}$$

$$\begin{array}{r} 312.68) 2188 \\ \dots \end{array}$$

$$\begin{array}{r} 312.68) 156 \end{array}$$

$$\begin{array}{r} 312.68) 156 \end{array}$$

$$\begin{array}{r} 312.68) 156 \end{array}$$

$$\begin{array}{r} 312.68) 156 \end{array}$$

$$\begin{array}{r} 312.68) 156 \end{array}$$

$$\begin{array}{r} 312.68) 156 \end{array}$$

Whence $r + e = 297.4337$, which may be taken for new r , and the operation repeated, if there be occasion.

Ex. 3.

What is the 7th root of 100000.

The nearest root of 100000 is 5, whence by Rule 3d,

$$\begin{aligned} P + Pq &= 100000 \\ P &= 78125 \end{aligned}$$

$$\begin{aligned} Pq &= 21875. \\ \text{whence } q &= .28 \text{ \&c.} \end{aligned}$$

$$\begin{aligned} \text{And } \sqrt[7]{P + Pq} &= \sqrt[7]{P} + \frac{1}{7} Aq - \frac{6}{14} Bq - \\ &\frac{13}{21} Cq - \frac{20}{28} Dq - \frac{27}{35} Eq \text{ \&c.} \end{aligned}$$

That is,

$$\begin{aligned} \sqrt{P} &= +5.000 \\ +\frac{1}{7}Aq &= .200 \\ -\frac{3}{14}Bq &= -.024 \\ -\frac{13}{21}Cq &= +.004 \\ -\frac{20}{28}Dq &= -.001 \\ \text{\&c.} & \end{aligned}$$

$$5.204 - .025 = 5.179$$

But because this converges slow, take 5.179 for the root, and involve it to the 7th power, and we have

$$\begin{aligned} P + Pq &= 100000. \\ P &= 99935.8652873094 \\ Pq &= 64.1347126906 \\ q &= .0006417587 \end{aligned}$$

$$\begin{aligned} \text{Then } \sqrt[7]{P} &= 5.1790000000 = A \\ +\frac{1}{7}Aq &= +.0004748097 = B \\ -\frac{3}{14}Bq &= - \quad \quad \quad 1306 = C \\ -\frac{13}{21}Cq &= + \quad \quad \quad \quad \quad 1 = D \end{aligned}$$

$$\begin{aligned} 5.1794748098 \\ -1306 \end{aligned}$$

$$5.1794746792 = \sqrt[7]{100000.}$$

T

Schol.

Schol. 1. If the root is required for only a few places of figures; the easiest way by far, is to extract it by the help of logarithms.

Schol. 2. From the foregoing process, the rule for extracting the cube root in arithmetic, may be demonstrated.

Let $a+e$ be the root, a the first figure, e the second. Then the cube is $a^3+3a^2e+3ae^2+e^3$; then a^3 the greatest cube contained in it, being subtracted; there remains $3a^2e+3ae^2$, setting aside e^3 as being very small. Divide this remainder by 3, and we have a^2e+ae^2 , from which to find e , this remainder or resolvend must be divided by $aa+ae$. That is, the resolvend must be divided by aa , the square of the root, and then to the divisor, there must be added ae , the product of the root by the quotient figure; and the whole will be the true divisor for finding e . But as e^3 was left out of the account; the root got this way will deviate from the true root; and therefore you must, after a few figures are had, begin the operation again, with the new root which you have already got.

PROBLEM XCII.

To extract the root of any affected equation, in numbers.

Preparation.

Suppose $Ax+Bx^2+Cx^3+Dx^4+Ex^5$ &c. = N . Put $r+e=x$, r being the first figure of the root; and to find r , put 1, 10, 100 successively for x ; and the nearest value of these being found, try the intermediate numbers 5, 50, &c. then expunging x , we have

$Ar+Ac$

$$\left. \begin{array}{l} Ar + Ae \\ Br^2 + 2Bre + Bee \\ Cr^3 + 3Cr^2e + 3Cree + Ce^3 \\ Dr^4 + 4Dr^3e + 6Dr^2e^2 + 4Dre^3 + De^4 \\ \&c. \end{array} \right\} = \begin{array}{l} Ax \\ Bxx \\ Cx^3 \\ Dx^4 \\ \&c. \end{array} = N.$$

Sum $P + ae + bee + ce^3 + de^4 \&c. = N.$

And $ae + bee + ce^3 + de^4 \&c. = N - P = f.$

Then since $e = \frac{f}{a}$ nearly, we shall have

$ae + be \times \frac{f}{a} = f.$ Or $ae + \frac{bf}{a}e = f.$ From whence we shall have this

1 RULE.

$$e = \frac{f}{a + b \times \frac{f}{a}} \text{ nearly.}$$

Or, if more exactness be required, we may bring in ee ; then $ae + bee = f$, whence this

2 RULE.

$$\frac{a}{b}e + ee = \frac{f}{b}, \text{ or } e = \frac{\frac{f}{b}}{\frac{a}{b} + e} = \frac{ce^3 + de^4 \&c.}{a + ee}$$

nearly, to be wrought by lxxxviii. Rule 2.

Or if e^3 be taken in for more exactness; proceed thus, $bee = f - ae$, and $ce^3 = \frac{cfe - caee}{b}$, whence

$$ae + bee + ce^3 = ae + bee + \frac{cf}{b}e - \frac{ca}{b}ee = f, \text{ whence}$$

3 R U L E.

$$\frac{a + \frac{cf}{b}}{b - \frac{ca}{b}} e + ee = \frac{f}{b - \frac{ca}{b}}, \text{ very near; to be wrought}$$

as Prob. lxxxviii. Rule 2.

In any of these rules the operation must be repeated after a few figures are had, by taking a new value of r , and proceeding as before.

Ex. 1.

$$\text{Let } 120x^3 + 3657x^2 - 38059x = 8007115.$$

By a few trials, you will find x to be greater than 30, and less than 40. Therefore suppose $r=30$, and $30+e=x$ the root sought, which being involved, and taking the least powers first, as in the rule, we have

$$\begin{array}{r} -1141770 - 38059e \\ + 3291300 + 219420e + 3657ee \\ + 3240000 + 324000e + 108000ee + 120e^3 \end{array} \} = 8007115.$$

Which being added,

$$\begin{array}{r} 5389530 + 505361e + 14457ee + 120e^3 = 8007115 \\ \text{and} \quad 505361e + 14457ee + 120e^3 = 2617585. \\ \text{or} \quad ae + bee + ce^3 = f. \end{array}$$

Then to shorten the work, divide by 1000, and then $505e + 14ee$ &c. $= 2617$, and by Rule 1,

$$e = \frac{2617}{505} = 5.18; \text{ or rather } e = \frac{2617}{505 + 14 \times 5.18} =$$

$\frac{2617}{577} = 4.53$. Whence $r+e$ or $x=34.5$ for a new operation. Which being involved, beginning at the highest power first, we have

$$\begin{array}{r} 4927635 + 42849e + 12420ee \\ + 4352744\frac{1}{2} + 252333e + 3657ee \\ - 1313035\frac{1}{2} - 38059e \end{array} \} = 8007115.$$

That

Sect. VIII. EQUATIONS.

277

That is,

$$7967343\frac{1}{2} + 257123e + 16077ee = 8007115.$$

or $257123e + 16077ee = 39771.25$

whence $15.9932e + ee = 2.473799$

and by rule 2d, $e = \frac{2.473799}{15.9932 + e}$.

$$\begin{array}{r} 15.9932) 2.473799 \quad (.1532 = e \\ + .1 \quad 160932 \end{array}$$

$$\begin{array}{r} 16.0932) 864479 \quad r = 34.5 \\ + 15 \quad 812160 \quad e = .1532 \end{array}$$

$$\begin{array}{r} 16.2432) 52319 \\ + .53 \quad 48888 \quad r + e = 34.6532 = x. \end{array}$$

$$\begin{array}{r} 10.296\frac{1}{2}) 3431 \\ + 3 \quad 3260 \end{array}$$

$$\begin{array}{r} 16.30 \quad 171 \end{array}$$

Ex. 2.

Let $z^4 - 3z^2 - 75z = 10000$.

Here by a few trials z will be found very near 10. Therefore let $r = 10$, and $r + e + z$. Then

$$\left. \begin{array}{l} z^4 = 10000 + 4000e + 600ee \text{ \&c.} \\ -3z^2 = -300 - 60e - 3ee \\ +75z = +750 + 75e \end{array} \right\} = 10000.$$

Being added,

$$10450 + 4015e + 597ee = 10000.$$

or $4015e + 597ee = -450$

or $6.725e + ee = -0.753769$

therefore e is negative, and by Rule 2.

$$e = \frac{-0.753769}{6.725 + e}$$

$$\begin{array}{r}
 6.725) \quad -0.753769 \quad (-0.114=e) \\
 \underline{-1} \quad \quad \quad \underline{-6625} \quad \cdot \\
 \hline
 6 \ 625) \quad -9126 \quad \quad \quad r=10.000 \\
 \underline{-11} \quad \quad \underline{-6515} \quad \quad \quad e=-.114 \\
 \hline
 6.515) \quad -26119 \quad \quad \quad r+e=9.886=z. \\
 \underline{-14} \quad \quad \underline{-26004} \\
 \hline
 6.501 \quad \quad \underline{-114}
 \end{array}$$

Again, put $r=9.886$, and $r+e=z$; then

$$\begin{array}{r}
 9551.738507135 + 3864.753593824e \\
 -293.198988 \quad \quad -59.316e \\
 +741.450 \quad \quad \quad +75e \\
 +586.397976ee = 10000; \text{ that is by addi-} \\
 \quad \quad \quad \underline{-3ee} \\
 \text{tion, } 9999.989519135 + 3880.437593824e + 583. \\
 397976ee = 10000; \text{ and transposing,} \\
 3880.437593824e + 583.397976ee = .010480865.
 \end{array}$$

$$\text{And } e = \frac{.010480865}{3880.437} = .00000270095, \text{ nearly.}$$

$$\text{Then } r+e = 9.88600270095 = z.$$

Ex. 3.

$$\text{Suppose } 7y^5 + 2100y^3 - 8000y^2 = 3850000000.$$

By a few trials y will be found between 50 and 60; therefore put $r=50$, and $r+e=y$; then expunge y . Or rather thus: Since the numbers are large, transform the equation (by Prob. xlii.), by putting $x = \frac{1}{10}y$, or $y = 10x$, which done we have $700000x^5 + 2100000x^3 - 800000x^2 = 3850000000$, or $7x^5 + 21x^3 - 8x^2 = 38500$. Then to extract the root of this, put $r=5$, and $r+e$ or $5+e=x$; and x being expunged, we have

$$\left. \begin{array}{r} 21875 + 21875e + 8750ee + 1750e^3 \\ + 2625 + 1575e + 315ee + 21e^3 \\ - 200 - 80e - 8ee \end{array} \right\} = 38500.$$

That is,

$$24300 + 23370e + 9057ee + 1729e^3 = 38500.$$

$$\text{and } 23370e + 9057ee + 1729e^3 = 14200.$$

Then by Rule 3, $\frac{f}{b} = 2711$, $\frac{ca}{b} = 4461$, whence

$$5.675e + ee = 3.0900.$$

$$\begin{array}{r} 5.675) 3.0900 (.5004 = e \\ + 5 \quad 30875 \\ \hline 6.175 \quad 25 \\ + 5 \\ \hline 6.675 \end{array}$$

Again, put $r = 5.5$, and repeating the operation,

$$\left. \begin{array}{r} 35229.90625 + 32027.187e + 11646.25ee \\ + 3493.875 + 1905.75e + 346.5ee \\ - 242. - 88.e - 8.ee \end{array} \right\} = 38500.$$

That is when added,

$$38481.78125 + 33844.937e + 11984.75ee = 38500.$$

And

$$33844.937e + 11984.75ee = 18.21875$$

$$\text{Then } \frac{18.21875}{33845} = 0005383 = \frac{f}{a} = e \text{ nearly,}$$

$$\text{Then } b \times \frac{f}{a} = 6451, \text{ and (Rule 1.)}$$

$$\frac{18.21875}{33844.937 + 6451} = .000538198 = e, \text{ more exactly.}$$

Then

$$r + e = 5.500538198 = x, \text{ and } y = 10x = 55.00538198.$$

The root may also be extracted as follows. Having got $ae + be^2 + ce^3 + de^4 = f$, as before directed; let v be the first figure of the value of e , s the second. Then putting $v + s$ for e ; $a \times v + s + b \times v + s^2 + c \times v + s^3 \&c. = f$; that is, $av + as + bv^2 + 2bvs \&c. + cv^3 + 3cv^2s \&c. = f$. And $as + 2bvs + 3cv^2s \&c. = f - av - bv^2 - cv^3 \&c.$ Whence

$$s = \frac{f - av - bv^2 - cv^3 \&c.}{a + 2bv + 3cv^2 \&c.} \quad \text{Whence this}$$

4 R U L E.

Having any equation given, proceed as in the other rules, till you get $ae + be^2 + ce^3 + de^4 \&c. = f$. Then find by repeated trials, the first figure v , of the value of e , so that $v \times a + bv + cv^2 + dv^3 \&c.$ may be nearly $= f$; and take that product from f , to find the remainder.

Then to find the next figure or figures; divide this remainder, by $a + 2bv + 3cv^2 + 4dv^3 \&c.$ the quotient is the said figure, which must be added to v , for a new value of v . Then repeat the operation with new v , viz. take $v \times a + bv + cv^2 + dv^3 \&c.$ from f , and divide the remainder by $a + 2bv + 3cv^2 \&c.$ and add the quotient to last v ; and so on.

And note, after the divisor once takes place, each new quotient may be continued to near as many figures, as all the preceding ones. Also in the divisor, you need not continue the parts of the divisor $2bv, 3bv^2 \&c.$ any farther in decimals, than to answer the number of figures, you would have true in the root.

General form.

$$v = \frac{f}{a}; \text{ or } v = \frac{f}{a+bv} \text{ nearer; or } v = \frac{f}{a+bv+cv^2} \text{ nearer still; \&c. then, next figure} \\ = \frac{f-av-bv^2-cv^3 \&c.}{a+2bv+3cv^2 \&c.}$$

Ex. 4.

Let $x^3 - 17x^2 + 54x = 350$.

Here x is greater than 10, and less than 20. Let $r=10$, $r+e=x$; then

$$\left. \begin{array}{l} 1000 + 300e + 30e^2 + e^3 \\ -1700 - 340e - 17e^2 \\ + 540 + 54e \end{array} \right\} = 350.$$

or $160 + 14e + 13e^2 + e^3 = 350$.

that is $14e + 13e^2 + e^3 = 510$.

To find e , try 1, 2, 3, &c. and you will find e very near 5, but something less. Therefore take $v=5$, and $v \times a + bv + cv^2 = 5 \times 14 + 65 + 25 = 520$, and $510 - 520 = -10$, then $a + 2bv + 3cv^2 = 219$, and

$\frac{-10}{219} = -.045$	$v = 5.000$
	$- .045$
	$e = 4.955$

Let new $v=4.95$; then $a + bv + cv^2 \times v = 509.119875$, and $510 - 509.119875 = 0.880125$.
 Also $a + 2bv + 3cv^2 = 216.2075$. Whence
 $s = \frac{0.880125}{216.2075} = .00407$, and $e = 4.95407$;
 whence

whence $z = 14.95407$. Or, if you please, put $v = 4.95407$ for a new operation.

Ex. 5.

$$\text{Let } 2x^4 - 16x^3 + 40x^2 - 30x = -1.$$

By a few trials, it appears that x is between 1 and 2. Therefore put $r = 1$, $r + e = x$. Then expunging x ,

$$\left. \begin{array}{r} 2 + 8e + 12ee + 8e^2 + 2e^3 \\ -16 - 48e - 48ee - 16e^2 \\ +40 + 80e + 40ee \\ -30 - 30e \end{array} \right\} = -1.$$

The sum is

$$\begin{array}{r} -4 + 10e + 4ee - 8e^2 + 2e^3 = -1 \\ \text{or } 10e + 4ee - 8e^2 + 2e^3 = 3 \\ \quad a \quad b \quad c \quad d \quad f \end{array}$$

Here we have $e = \frac{3}{10} = .3$, or more exactly

$$e = \frac{3}{10 + 4e} = \frac{3}{11.2} = .26 = v.$$

Then for the next figures of the root, $v \times a + bv + cv^2 + dv^3 = 2.73893$, and $3 - 2.73893 = .26106$. Also $a + 2bv + 3cv^2 + 4dv^3 = 10.598$, and

$$\frac{26106}{10.598} = .246$$

$$v = .26$$

$$\text{then } e = .2846$$

Take new $v = .2846$, then $v \times a + bv + cv^2 + dv^3 = 2.99869539$, and $3 - 2.99869539 = .00130461$. Also $a + 2bv + 3cv^2 + 4dv^3 = 10.51728$. Then

$$.00130461$$

$$\frac{.00130461}{10.5173} = .00012404$$

$$.2846 = v$$

$$.28472404 = e,$$

whence $r + e$ or $x = 1.28472404$.

The roots of equations may also be extracted by help of the Rule of Falso Position in Arithmetic, as follows.

5 R U L E.

In such equations as contain surds, exponential quantities, &c. make two suppositions in numbers, for the root, as near as you can get them. Then each of these being put in the equation instead of the root, you must mark the errors (that is, the excess or defect) arising from each of them.

Then multiply the difference of the supposed numbers by the least error, and divide the product, by the difference of the errors, if they are like, (that is, both excesses or both defects); or by the sum, if unlike. Then

The quotient is the correction of the number belonging to the least error; and is to be added if that number was too little; or subtracted, if too great. This gives the root nearer than before.

In like manner try this root, and the nearest of the former, or else take a new supposed number; then find their errors, and proceed as before, and you will get a root still nearer. And thus by repeating the operation, you may continually approximate, as near as you will, to the true root.

Ex.

Ex. 6.

Suppose $x^x = 100$, to find x .

By the nature of logarithms $x \times \log: x = \log: 100 = 2$.

Here x , by a few trials, will be found greater than 3, and less than 4. Suppose $x = 3\frac{1}{2}$; then $l:x = .5440680$, and $xl:x = 1.9042380$, which should be equal to

$$\frac{2}{- .0957620} = 1 \text{ Er. too little.}$$

Again, suppose $x = 3.6$, then $l:x = .5563025$, and $xl:x = 2.0026890$

we have $\frac{2}{+.0026890} = 2 \text{ Er. too great.}$ Hence

1 num. 3.5	1 er. —.095762
2 num. 3.6	2 er. +.002689
diff. 0.1	sum .098451

Then $\frac{0.1 \times .002689}{.09845} = .00273 = \text{cor.}$

2 num. 3.60000	
correct. —.00273	

$$3.59727 = x.$$

Again, suppose $x = 3.597$, then $l:x = .5559404$, and $xl:x = 1.9997176$, which subtracted from 2, gives —.0002824 the error, too little. Whence

2 num. 3.600,	2 er. = + .0026890
3 num. 3.597,	3 er. = — .0002824
diff. .003,	sum .0029714

Then

Sect. VIII. EQUATIONS.

Then $\frac{.003 \times .0002824}{.0029714} = .000285$ the cor.

$$\begin{array}{r} 3 \text{ num. } 3.597 \\ \text{cor. } + .000285 \\ \hline \end{array}$$

$x = 3.597285$ as required.

Ex. 7.

If s be the sine of an arch z , $\text{rad.} = 1$, and $4sz = 5$, to find s and z .

By division $sz = 1.25$. The length of 1 degree is $= .01745329$ &c. By a few trials, we may find z between 70 and 80 degrees. Suppose $z = 70$ deg. then $.01745 \times 70 = 1.2215$; also $S.70 = .939 = s$, and $sz = 1.1469$, and $1.25 - 1.1469 = .1031$ the first error, too little.

Again, suppose $z = 75$ deg. then $.01745 \times 75 = 1.3087$, and $s = .966$, and $sz = 1.2642$; and $1.2642 - 1.25 = .0142$ the second error, too much. Hence

1 num. 70	1 er. —.1031
z num. 75	2 er. +.0142
5.	sum .1173

Then $\frac{5 \times .0142}{.1173} = \frac{.0710}{.117} = .60$ the cor.

$$\begin{array}{r} 2 \text{ num. } 75.0 \\ \text{— cor. } .6 \\ \hline \end{array}$$

$z = 74.4$

Again, let $z = 74.4 = 74^\circ : 24'$, $s = .9631626$; then $.01745329 \times 74.4 = 1.298524$, and $1.2985245 = 1.2506895$, from which subtract 1.25, then $.0006895 = 3d$ error, too much.

2 num. 75 3 num. 74.4 <hr style="width: 50%; margin-left: auto; margin-right: auto;"/> diff. .6	2 er. +.0143937 3 er. +.0006895 <hr style="width: 50%; margin-left: auto; margin-right: auto;"/> diff. .0133105
---	---

Then $\frac{.6 \times .0006895}{.01331} = .0310.$

$$\begin{array}{r} 3 \text{ num. } 74.400 \\ \text{--- cor. } .031 \\ \hline \end{array}$$

and $z = 74.369 = 74^\circ : 22' : 8''$
 and $s = .9630372.$

SCHOLIUM.

There are also other ways of extracting the roots of equations, though not much different from some of the foregoing ones, particularly a method of Sir *I. Newton's*, which is like the process used in the second method foregoing; the principal difference being, that he every where takes a new letter, where we find a new value of e .

Also surd or transcendental equations, may be resolved by reducing some of the quantities to infinite series; proceeding by the rules of Sect. VI.

In equations, where the terms involve a great many factors, which makes it tedious to multiply them together; it will be a shorter way to add the logarithms of the several factors together; and then find the number belonging, which will be the numeral coefficient of that term. And thus all the coefficients of the particular terms may be found.

We may note, that though the third rule converges faster than the rest; yet, as there is so much trouble in finding the coefficients, and divisors, it will be found not so expeditious as the second, or even the first.

In

In making use of the second rule, after half the number of places are found for the value of e ; it will be needless to form new divisors; for the rest of the figures will be as truly found by plain division. For what is added to the divisor, in places so far back, does not at all affect the quotient.

The root may also be extracted as in the following problem, and the coefficients $a, b, c, \&c.$ found as there directed; which is a compendious method, when the equation consists of many terms.

P R O B L E M X C I I I .

To extract the root of the infinite series $Az + Bz^2 + Cz^3 + Dz^4 + Ez^5 \&c. = N$, in numbers; supposing this series to converge fast enough.

Preparation.

Take r as near the root z , as you can find it; and let $r + e = z$, and z being expunged, we have

$$\left. \begin{array}{l} Ar + Ae \\ + Br^2 + 2Bre + Be^2 \\ + Cr^3 + 3Cr^2e + 3Cre^2 + Ce^3 \\ + Dr^4 + 4Dr^3e + 6Dr^2e^2 + 4Dre^3 + De^4 \\ + Er^5 + 5Er^4e + 10Er^3e^2 + 10Er^2e^3 + 5Ere^4 + Ee^5 \\ \&c. \end{array} \right\} = N$$

the sum

$$P + ae + be^2 + ce^3 + de^4 + ge^5 = N$$

$$\text{and } ae + be^2 + ce^3 + de^4 + ge^5 = N - P = f.$$

Whence this

R U L E .

Take r very near z , and let $r + e = z$, then substitute the powers of $r + e$ for those of z , till you get $P + ae + be^2 + ce^3 \&c. = N$, and $ae + be^2 \&c. = N - P = f$, which equation is to be resolved by Prob. lxxxviii; or else the equation $ae + be^2 + ce^3 + de^4$

$+de^4$ &c. $=f$, is to be resolved by some of the rules in the last problem, and the operation repeated if there be occasion.

And here the coefficients $a, b, c, d, \&c.$ are most easily had from the terms, which compose the value of P ; for we have $P = Ar + Br^2 + Cr^3 + Dr^4$ &c. Whence

$$a = \frac{Ar + 2Br^2 + 3Cr^3 + 4Dr^4 \&c.}{r}$$

$$b = \frac{Br^2 + 3Cr^3 + 6Dr^4 + 10Er^5}{rr}$$

$$c = \frac{Cr^3 + 4Dr^4 + 10Er^5 \&c.}{r^3}, \text{ and so on; where}$$

the numbers in a , are 1, 2, 3, 4, &c; in b , 1, 3, 2×3 , 2×5 , 3×5 , 3×7 , 4×7 , 4×9 , &c. in $c = 1, 4, \frac{5}{2}p, \frac{6}{3}q, \frac{7}{4}r, \frac{8}{5}s, \frac{9}{6}t$ &c. where p, q, r, s, t , &c. are the foregoing terms. And in finding a, b, c , &c. you must go through all the terms, till they grow very small, and at last vanish. But you need not find above two or three of these coefficients a, b, c , &c. and each succeeding one may consist of fewer places of figures.

Example.

$$\text{Let } x - \frac{1}{2}x^2 + \frac{1}{2.3}x^3 - \frac{1}{2.3.4}x^4 + \frac{1}{2.3.4.5}x^5$$

$$\&c. = \frac{2}{7}.$$

Here by several trials x is found nearly $=\frac{1}{3}$; therefore put $r = \frac{1}{3}$, and $r + e = x$. Then $P = A + Br^2 + Cr^3$ &c. that is,

$$\begin{aligned} r &= .333333 \\ + \frac{1}{2}r^3 &= 6172 \\ + \frac{1}{12}r^5 &= 34 \end{aligned}$$

$$\begin{aligned} -\frac{1}{2}r^2 &= -.055555 \\ -\frac{1}{24}r^4 &= 514 \\ -\frac{1}{720}r^6 &= 2 \end{aligned}$$

$$\begin{aligned} &-.339539 \\ &-.056071 \end{aligned}$$

$$P = .283468.$$

$$\begin{array}{r} .333333 \text{ --- } .111110 \\ + 18516 \text{ --- } 2056 \quad .352019 \\ \text{then } a = + 170 \quad - 12 \quad \text{--- } .113178 \\ \hline \phantom{\text{---}} \times 3 = \\ \phantom{\text{---}} = \end{array}$$

.716523. Also

$$\begin{array}{r} -.055555 + .018516 \\ - 3084 + 340 \quad \text{--- } .058669 \\ b = - 30 \quad \text{--- } + .018856 \\ \hline \phantom{\text{---}} \times 9 \\ \phantom{\text{---}} = \end{array}$$

= -.3583. Hence

$$.283468 + .71652e - .3583ee = .285714$$

$$\text{and } .71652e - .3583ee = .002246$$

$$\text{and } 1.9998e - ee = .00627$$

$$e = \frac{.00627}{1.999 - e}$$

$$\begin{array}{r} 1.999 \cdot .006270 \text{ (}.00314 = e \\ - 3 \quad 5988 \end{array}$$

$$\begin{array}{r} 1.996 \quad 282 \\ \quad 199 \\ \quad 83 \\ \quad 79 \\ \hline \quad 4 \end{array}$$

U

Then

$$\text{Then } r = .33333$$

$$e = .00314$$

$$z = .33647$$

or put $r = .33647$ for another operation.

SCHOLIUM.

If the series breaks off, then it is no matter whether it converges or not. And in that case it coincides with the last problem, and may be solved by any of the rules therein.

And if e be very small, the equation $ae + bee + ce^3 \&c. = f$, may be expeditiously solved by Prob. lxii. Rule 1, in which you need only use the three first terms; which will be shorter than taking new r . But that rule cannot so conveniently be applied to the given series, because it does not converge so fast as this.

PROBLEM XCIV.

To extract the root in numbers of the infinite series $Az + Bz^3 + Cz^5 + Dz^7 \&c. = N$; supposing it to converge fast.

Preparation.

Take r as near the root as it can be found by trials, and put $r + e = z$, and expunging z , we shall have,

$$\left. \begin{aligned} &Ar + Ae \\ &+ Br^3 + 3Br^2e + 3Bree + Be^3 \\ &+ Cr^5 + 5Cr^4e + 10Cr^3e^2 + 10Cr^2e^3 + 5Cre^4 \&c. \\ &+ Dr^7 + 7Dr^6e + 21Dr^5e^2 + 35Dr^4e^3 + 35Dr^3e^4 \\ &+ Er^9 + 9Er^8e + 36Er^7e^2 + 84Er^6e^3 + 126Er^5e^4 \end{aligned} \right\} = N.$$

the sum

$$P + ae + bee + ce^3 + de^4 \&c. = N.$$

and

and $ae + bee + ce^2 + de^3 \&c. = N - P = f.$

Whence this

R U L E.

Assume r by trials very near z , and $r + e = z$, then substitute the powers of $r + e$ for z , in the given series, till you get $P + ae + be^2 + ce^3 \&c. = N$,

And $ae + be^2 + ce^3 \&c. = N - P = f.$

Where $P = Ar + Br^2 + Cr^3 + Dr^4 + Er^5 \&c.$

$$a = \frac{Ar + 3Br^2 + 5Cr^3 + 7Dr^4 \&c.}{r}$$

$$b = \frac{3Br^2 + 10Cr^3 + 21Dr^4 \&c.}{rr}$$

$$c = \frac{Br^2 + 10Cr^3 + 35Dr^4 \&c.}{r^2}$$

&c.

Where the numbers of a , are 1, 3, 5, 7, &c. of b ; 3, 2×5 , 3×7 , 4×9 , 5×11 , 6×13 , &c. And each series is to be continued till the terms become very small and vanish; which will happen in a little time, because the given series converges. The terms of a , b , c , are easily had from the terms of P , as above, without much labour; then having got $ae + bee + ce^3 \&c. = f$, in numbers; find the root e , by Prob. lxxxviii. or by some of the rules in Prob. xcii.

Example.

$$\text{Let } y + \frac{1}{6} y^3 + \frac{3}{4 \cdot 5} y^5 + \frac{3 \cdot 5}{7 \cdot 2 \cdot 4 \cdot 6} y^7 + \frac{3 \cdot 5 \cdot 7}{9 \cdot 2 \cdot 4 \cdot 6 \cdot 8} y^9 \&c. = .698132, \text{ to find } y.$$

The series abridged will be $y + \frac{1}{3} y^3 + \frac{1}{5} y^5$

U 2

+ $\frac{5Ryy}{7}$ &c. = 698132; Q, R, S, being the numerators. By a few trials, y is nearly = .6, put $r = .6$; then

$r = 600000$	$r = .600000 = Ar$
$Q = 108000$	3) $Q = 36000 = Br^3$
$R = 29160$	5) $R = 5832 = Cr^5$
$S = 8748$	7) $S = 1249 = Dr^7$
$T = 2756$	9) $T = 306$ &c.
$V = 893$	11) $V = 81$
$W = 295$	13) $W = 23$
$X = 98$	15) $X = 6$
$Y = 33$	17) $Y = 2$
$Z = 11$	19) $Z = 1$

$$\underline{\underline{.643500 = P.}}$$

Then $a = 1.2500$, the sum of the first column divided by r . $b = .585$; whence

$$.643500 + 1.250e + .585e = .698132$$

$$\text{and } 1.250e + .585e = .054632$$

$$\text{and } e = \frac{.0546}{1.25} = .043 \text{ nearly.}$$

$$\text{and } e = \frac{.054632}{1.2500 + .585 \times .043} = \frac{.054632}{1.2500 + .0251}$$

$$= \frac{.054632}{1.275} = .04284 \text{ more exactly.}$$

$$\text{add } r = .60000$$

$$\underline{\underline{.64284 = y.}}$$

or take new $r = .6428$ for another operation.

PROBLEM XCV.

To extract the roots of two given equations, containing two unknown quantities x, y ; though never so compounded.

R U L E.

By several trials find two near values of x and y , viz. r and s , and put $r+e=x$, and $s+v=y$. And instead of the powers of x and y , put in those of $r+e$, and $s+v$. Then involve all surds by the binomial theorem (Prob. lviii.), also reduce logarithmic quantities to series (Prob. lxxxiv, lxxxv.), and the like for all compound quantities; so that at last the equations may consist only of simple terms. And in doing this, reject all powers of e and v above the first, and also their products.

Then you will have two simple equations of e and v , which being resolved, will give their values; and from hence x and y will be known. Then put new r and s for these values of x and y , and repeat the operation, which may be done as often as you please, till you get the roots as near as you have a mind. And the same form may stand and serve for all these operations.

Ex. 1.

$$\text{Suppose } \sqrt{yy-xx} + \frac{2xy}{\sqrt{yy+2x}} = 20 = b$$

$$\text{and } \frac{\log x + \sqrt{xx+yy}}{y} = 0.096 = c.$$

Let $a=2$. And by some trials we find x near 4, and y near 13; then put $r=4$, $s=13$, and by involution, and putting $r+e$ for x , and $s+v$ for y , we have

$$\frac{ss-rr+2sv-2re^{\frac{1}{2}}}{ss+2sv+ar+ae^{-\frac{1}{2}}}=b \quad \text{and} \quad \frac{2rs+2rv+2se}{ss+2sv+ar+ae^{-\frac{1}{2}}}=c$$

$$\frac{ss-rr+2sv-2re^{\frac{1}{2}}}{ss+2sv+ar+ae^{-\frac{1}{2}}}=b \quad \text{and}$$

$$\frac{\log: r+e+rr+2re+ss+2sv^{\frac{1}{2}}}{s+v}=c.$$

$$\text{but } \frac{ss-rr+2sv-2re^{\frac{1}{2}}}{ss+2sv+ar+ae^{-\frac{1}{2}}} = \sqrt{\frac{ss-rr}{ss+ar}} + \frac{sv-re}{\sqrt{ss-rr}}$$

$$\text{and } \frac{2rs+2rv+2se}{ss+2sv+ar+ae^{-\frac{1}{2}}} = \frac{1}{\sqrt{ss+ar}} - \frac{2sv+ae}{2 \times \sqrt{ss+ar}^{\frac{3}{2}}}$$

$$\text{also } \frac{rr+ss+2re+2sv^{\frac{1}{2}}}{ss+2sv+ar+ae^{-\frac{1}{2}}} = \sqrt{\frac{rr+ss}{ss+ar}} + \frac{re+sv}{\sqrt{rr+ss}}$$

Put $dd=ss-rr$, $ff=ss+ar$, $gg=ss+rr$ Then we

$$\text{have } d + \frac{sv-re}{d} + 2rs+2rv+2se \times \frac{1}{f} - \frac{2sv+ae}{2f^{\frac{3}{2}}}$$

$$= b, \text{ that is, (1) } \frac{2s}{f} - \frac{r}{d} - \frac{ars}{f^{\frac{3}{2}}} \times e + \frac{s}{d} + \frac{2r}{f} - \frac{2rs}{f^{\frac{3}{2}}}$$

$$\times v = b-d - \frac{2rs}{f}.$$

$$\text{Again, } \frac{\log: r+e+rr+ss+2re+2sv^{\frac{1}{2}}}{s+v} = c =$$

$$\frac{\log: r+e+g+\frac{re+sv}{g}}{s+v}, \text{ and } \log: r+e+g+\frac{re+sv}{g}$$

$$= s+v \times c. \text{ Put } t=r+g, l=\log:t.$$

$$m = .4342945, \text{ then (Prob. lxxxiv.) } l + \frac{mte+msv}{tg} = cs+cv, \text{ which reduced, is (2)}$$

$$mte+msv = tg \times cs - l. \text{ Then numbers being}$$

substituted in these two equations, give

$$(1) 1.588e + 1.075v = -0.190$$

$$(2) 7.643e - 17.3326v = +0.59536;$$

And

Sect. VIII. EQUATIONS. 295

And these equations being resolved, give $e = -.0743$, and $v = -.0671$; whence $r + e$ or $x = 3.926$; and $s + v$ or $y = 12.933$.

Or for another operation, put $r = 3.926$ and $s = 12.933$, and finding new values for d, f, g, t , and l , you will have two equations, which will give e and v more exactly.

Ex. 2.

Let $x + \log y = b = 8.7679114$.

and $y + \log x = c = 3.4760046$.

By a few trials, we find x nearly $= 8$, and $y = 2\frac{1}{2}$. Put $r = 8$, and $r + e = x$; also $s = 2\frac{1}{2}$, and $s + v = y$. Also $M = .4342945$; then we have

$$r + e + l \overline{s + v} = b = r + e + l \cdot s + \frac{Mv}{s} \text{ (Pr. 84.)}$$

$$\text{and } s + v + l \overline{r + e} = c = s + v + l \cdot r + \frac{Me}{r}.$$

These equations reduced become

$$e + \frac{Mv}{s} = b - r - l \cdot s.$$

$$v + \frac{Me}{r} = c - s - l \cdot r.$$

And put into numbers are

$$e + .1737v = .3700.$$

$$\text{and } v + .0543e = .0730.$$

Which equations being resolved give $e = .3608$, and $v = .0535$; whence

$$r = 8.0000$$

$$+ e = .3608$$

$$x = 8.3608$$

$$s = 2.5$$

$$+ v = .0535$$

$$y = 2.5535$$

U 4

Again,

Again, put $r=8.3608$, and $s=2.5535$, for another operation; whence will be found $\frac{M}{s} =$

$.170078$, $\frac{M}{r}=.051944$; and $b-r-l: s = -$
 $.0000245$, $c-s-l:r=.0002568$, and from thence will arise these two equations,

$$e + .170078v = -.0000245,$$

$$\text{and } v + .051944e = .0002568.$$

Which being resolved, give $e=-.0000688$, and $v=.0002604$; therefore

$r = 8.3608000$	$s = 2.5535000$
$+e = -.0000688$	$+v = .0002604$
<hr style="width: 100%;"/>	<hr style="width: 100%;"/>
$x = 8.3607312$	$y = 2.5537604.$

S E C T. IX.

The geometrical Construction of Equations.

THE *construction* of equations, is the drawing right lines or curves, after such a manner, as by their interfections, to give the roots of the equation proposed. This method is used for avoiding the tediousness of computation; and is exact enough for finding two or three of the first figures of the root, but not more. For where great exactness is required, we are not to trust to a construction by lines; but make a computation in numbers, to find the root.

In geometrical constructions, the simplest is always to be made use of, or that by which we can come the shortest way, to the roots of the equation proposed.

But since the extraction of roots by converging series, is now brought to so great perfection; geometrical constructions are almost laid aside. Therefore I intend to trouble the reader only with the shortest methods of constructing equations as far as the fourth power. When we come to higher powers, there is so much trouble and difficulty in drawing the lines proper for them, that their interfections cannot be depended on; and one may sooner extract the root in numbers.

P R O B L E M X C V I.

To construct a simple equation.

R U L E.

1. When there are several simple quantities, connected by the signs + and —. From a certain point,

point, draw a right line, from which point set all affirmative quantities one way, one adjoining to another; from the last point, set all the negative quantities the contrary way, adjoining to each other as before. Where the last ends, the distance from thence, to the first point, gives the sum (or difference) of all; which is affirmative or negative, according as it lies on the affirmative or negative side of the first point assumed.

2. When you have the square root of two quantities, find a mean proportional between them, by Prob. 16. B. VIII. Geometry.

3. To reduce two compound quantities to the same designation. By Prob. 15. B. VIII. Geometry, find one or more proportionals thus; say, as the first letter of the first quantity, to the first in the second, so the second in the second to that fourth proportional. Again, as the second letter in the first quantity to the third letter in the second; so the fourth proportional last found, to another fourth proportional. Proceed thus till all the letters in one quantity be exhausted.

Note, when any term is of too low a dimension, make 1 to be one of the factors, as oft as it is wanted. And when you have several simple quantities, add them into one, by Art. 1.

4. For many compound quantities, reduce them all to the same designation by Art. 3.

Ex. 1.

Suppose $a + b - c = x$.

Fig. Draw the line DAB, and from the first point
 3 A, set off $AB = a$, and make $BC = b$, both forward; lastly, make $CE = c$, backward. Then $+AE = x$. *Ex.*

Ex. 2.

Let $ax = bc$, to find x .Make $AB (a) : AC (b) :: AD (c) : AE$ \leftarrow
 $= x$, (by Prob. 15. Geom.)

Ex. 3.

Suppose $2abcx = 5defg$.Make as $2a : 5d :: e : m$ (Pr. 15. Geom.)and $b : f :: m : n$ and $c : g :: n : p$ then $2abc : 5dfg :: e : p$ and $2abcp = 5defg = 2abcx$ or $p = x$.

Ex. 4.

Let $abx - f\sqrt{bc} \times x = dd\sqrt{ac - bd}$.

By Prob. 15. Geom. make $a : b :: d : m$;
then $bd = am$; and $\sqrt{ac - bd} = \sqrt{ac - am}$.
make $c - m = n$, then $\sqrt{ac - bd} = \sqrt{an}$.
Find p a mean proportional between a and
 n , and q a mean between b and c , (Prob. 16.
Geom.) then the given equation becomes
 $abx - fqx = ddp$.

Reduce these three terms to the same designation, thus $a : f :: q : r$, whence $fq = ar$, in like manner $dd = as$; then the equation is $abx - arx = asp$, or $bx - rx = sp$. Put $b - r = t$, then $tx = sp$, and $t : s :: p : x$ required.

Ex. 5.

Let $2abcd - efgb + 3klmn = 4qrstx - 5noplx$,
to find x .

Reduce all the quantities to the same designation, then

 $4qrstx$

Fig.

$$4qrsx = 5nopl$$

$$4qrstv = 2abcd$$

$$4qrstw = eefgb$$

$$4qrsty = 3kllmn.$$

then the equation becomes

$$4qrstv - 4qrstw + 4qrsty = 4qrstz - 4qrsxz.$$

that is, $tv - tw + ty = tz - xz$

Put $v - w + y = A$, $t - x = B$, then $At = Bz$,
or $B : A :: t : z$.

P R O B. XCVII.

To construct a quadratic equation.

I R U L E.

If it is a pure quadratic; reduce the quantities concerned therein to the same designation (Prob. xcvi. Art. 3.) by which means surds will be denoted by simple quantities, and at last you will get all the known quantities equal to a known square, whose side is the root.

Ex. 1.

$$\text{Suppose } yy = ab - \frac{cdd}{b} + d\sqrt{aa - bc}.$$

Make $b : c :: d : m$, (Prob. 15. Geom.)

then $cd = bm$, and $\frac{cdd}{b} = \frac{bmd}{b} = md$. Also make $a : b :: c : n$, then $bc = an$; whence $yy = ab - md + d\sqrt{aa - an}$.

Let p be a mean between a and $a - n$, (Prob. 16. Geom.) then $\sqrt{aa - an} = p$. whence $yy = ab - md + dp$.

Let $d : a :: b : q$, then $ab = dq$.

then $yy = dq - dm + dp$.

Lastly, put $q - m + p = r$, and find s a mean between d and r ; then $yy = dr = ss$, and $y = s$.

2 R U L E.

2 RULE.

In adfected quadratics, reduced to this form $aa+ba=nn$. Draw a right line AD, then take any point C; and make $CB=\frac{1}{2}b$, towards the right hand if $+b$, or towards the left, if $-b$. Erect the perpendicular $BF=n$. From the center C through F, describe the circle AFD, to cut AD. Then (BD, BA) the distances of B, from the intersections A, D, are the two roots, the affirmative to the right hand, the negative to the left of B. 5.

Ex. 2.

Let $aa+3a=10$.

Draw the line AD, make $CB=1\frac{1}{2}$ on the right; find a mean proportional between 1 and 10, set it in the line BF, perpendicular to AB, with the radius CF describe the circle AFD; then $a=BD=+2$, and $a=BA=-5$, the two roots required. 5.

Ex. 3.

Suppose $aa-3a=10$.

Draw the line AD, make CB (on the left of C) $=1\frac{1}{2}$, find a mean proportional between 1 and 10; at B erect the perpendicular BF, and make BF = the mean; with the radius CF describe the circle AFD; to cut AD in A and D; then $a=BD=+5$, and $a=BA=-2$, the roots required. 6.

3 RULE.

In such quadratic equations as may be reduced to this form, $aa+ba=-nn$. From any point C as a center, in the right line BD, with radius $\frac{1}{2}b$, describe the circle BFD, erect a perpendicular at D on the right, if it be $+b$, or on 7.

Fig. on the left at B, if it is $-b$; whose length is $BA = n$. Through A draw AFG parallel to BD, to intersect the circle in F and G; then AF and AG are the two roots of the equation; which are affirmative, if they lie towards the right hand from A; or negative, if on the left.

Note, if the parallel does not cut the circle, or touch it, the equation is impossible.

Ex. 4.

Suppose $aa + 7a = -10$.

8. With the radius $3\frac{1}{2}$, and center C, describe the circle BFD. At the end of the diameter D, on the right, raise the perpendicular DA, a mean between 1 and 10. Through A draw AFG parallel to the diameter BD, to cut the circle in F and G; and AF, AG, being on the left from A, are two negative roots; $a = AF = -2$, and $a = AG = -5$.

Ex. 5.

Let $aa - 7a = -10$.

7. With the radius $CB = 3\frac{1}{2}$, and center C, describe the circle BFD; at the end B, of the diameter BD on the left, raise the perpendicular BA, equal to the mean between 1 and 10. Through A, draw AFG parallel to the diameter BD, to cut the circle in F and G; then AF, AG, lying on the right hand from A, are the two affirmative roots; and $a = AF = 2$, and $a = AG = 5$.

4 R U L E.

When the unknown quantity is higher than the square, and the index in one term double to that in the other; it may be brought to some of the foregoing forms, whose highest term is a square.

Sect. IX. EQUATIONS.

303

a square. Assume an unknown quantity, whose rectangle with some given quantity, is equal to the square of the unknown quantity proposed; for this substitute that rectangle; and you will have an equation as required.

Ex. 6.

Let $xx - bzz = n$.

Assume $dx = zz$, then by substitution, $ddxx - bdx = n$, and $xx - \frac{b}{d}x = \frac{n}{dd}$. Let $d : b :: 1 : p$; then $b = dp$; also make $d : n :: 1 : q$, and $d : 1 :: q : r$, then $dd : n :: 1 : r$, and $ddr = n$. And the least equation becomes $xx - \frac{dp}{d}x = \frac{ddr}{dd}$, that is, $xx - px = r$, which is to be constructed by some of the former rules.

To demonstrate these rules. Let $aa + ba = nn$. Here we have $CB = \frac{1}{2}b$, $BF = n$, and if $BD = a$, then CD or $CF = a + \frac{1}{2}b$, and $CF^2 = CB^2 + BF^2$, that is, $a + \frac{1}{2}b = \frac{1}{2}bb + nn$. But if $BA = -a$, then CA or $CF = -a - \frac{1}{2}b$, and $-a - \frac{1}{2}b = \frac{1}{2}bb + nn$. In both cases $aa + ba = nn$. 5.

Again, if $aa - ba = nn$, we have as before $CB = \frac{1}{2}b$, $BF = n$, and if $BD = a$, then CD or $CF = a - \frac{1}{2}b$, and $a - \frac{1}{2}b = \frac{1}{2}bb + n$. 6.

But if $AB = -a$, then AC or $FC = -a + \frac{1}{2}b$, and $-a + \frac{1}{2}b = nn + \frac{1}{2}bb$, in both cases $aa - ba = nn$.

Again, if $aa - ba = -nn$; here $BC = \frac{1}{2}b$, $BA = n$, and $AG = BD - AF$, therefore if $AF = a$, then $AG = b - a$, and $AG \times AF = AB^2$, that is, $b - a \times a = nn$. 7.

But

Fig. But if $AG = a$, then $AF = BD - AG = b - a$, and $AF \times AG = AB^2$, or $b - a \times a = nn$. In both cases $aa - ba = -nn$.

8. Lastly, when $aa + ba = -nn$, then $BC = \frac{1}{2}b$, $BA = n$, as before; then if $AF = -a$, then $AG = BD - AF = b + a$, and $AF \times AG = AD^2$, or $-a \times b + a = nn$.

But if $AG = -a$, then $AF = BD - AG = b + a$, and $AF \times AG = AD^2$, or $b + a \times -a = nn$. In both cases $aa + ba = -nn$.

P R O B. XCVIII.

To construct cubic and biquadratic equations.

9. To construct a cubic equation, that has all its roots real, by a circle. Let the radius $OB = R$, sine $EF = s$, GH the sine of the arch GB or $3BE$. Then by trigonometry, $3s - \frac{4s^3}{RR} = GH$. Draw CD parallel to AB , and put $SF = c$, $ES = x$, $GH = b$, then $c + x = s$, whence $3 \times c + x - \frac{4}{RR} \times c + x^3 = b$, this reduced gives $x^3 + 3cx^2 + 3ccx + c^3 = 0$.
- $$- \frac{1}{2}RR + \frac{1}{2}bRR$$
- $$- \frac{1}{2}cRR$$

Suppose this cubic equation be given,

$x^3 + px^2 + qx + r = 0$. Comparing this with the former, and equating the coefficients, we have $p = 3c$, and $c = \frac{1}{3}p$. Also $q = 3cc - \frac{1}{2}RR = \frac{1}{2}p^2 - \frac{1}{2}RR$, whence $R = \frac{2}{3}\sqrt{pp - 3q}$, and $r = c^3 + \frac{1}{2}bRR - \frac{1}{2}cRR$; whence $b = \frac{gr - pq}{pp - 3q} + \frac{2}{3}p$.

Hence arises the following

I R U L E.

I RULE.

Having the equation $x^3 + px^2 + qx + r = 0$. 10.
given;

1. With the radius $\sqrt[3]{pp-3q}$, describe the circle BGAK.
2. Draw the diameter AB, and CD parallel to it, at the distance of $\frac{1}{3}p$; above it, if it be $+p$, but below it, if $-p$.
3. Draw also ZG parallel to AB, at the distance $\frac{qr-pq}{pp-3q} + \frac{2}{3}p$, above it, if it is affirmative; or below it, if negative. Let it cut the circle in G.
4. Take the arch $BP = \frac{1}{3}BG$; and make $PQ = QK = KP$.
5. From the points P, Q, K, let fall perpendiculars, upon the line CD, which will be the roots of the equation; the affirmative above the line, and the negative below it.

SCHOLIUM.

If $3q$ be greater than pp , the equation is impossible; for in this case the equation has two impossible roots.

Also if $p=0$, then the radius of the circle $OB = \sqrt[3]{-3q}$; and CD coincides with AB; and the distance of ZG from AB is $-\frac{3r}{q}$.

And if q is affirmative, the equation is impossible. These constructions are extremely easy.

Ex. 1.

$$\text{Let } x^3 + 9x^2 - 22x - 120 = 0.$$

Here the radius $OB = \sqrt[3]{pp-3q} = \sqrt[3]{81+66} = 8.0829$, and $\frac{1}{3}p = 3$, the distance of CD, above AB. 11.

X

And

Fig.

- And $\frac{9r-pq}{pp-3q} + \sqrt[3]{p} = \frac{-1080+198}{81+66} + \sqrt[3]{6}$
11. $= -6+6=0$, the distance of GZ from AB; therefore ZG coincides with AB; and the arch BG and also its third part is 0, and P falls on B; and making $PQ=QK=KP$, and letting fall perpendiculars on CD, we shall have $PS=-3$, $QT=+4$, and $KT=-10$, the three roots required.

Ex. 2.


Suppose $x^3-17x^2+82x-120=0$.

12. The radius $OB = \sqrt[3]{289-246} = 4.37$,
 $p = -5.66$, the distance of CD below AB,
 and $\frac{9r-pq}{pp-3q} + \sqrt[3]{p} = \frac{-1080+1394}{289-246} - \frac{34}{3} =$
 $7.302-11.333 = -4.031$, the distance of
 GZ below AB. Take BP the third part of
 BG, and making $PQ=QK=KP$; and mea-
 suring the perpendiculars upon CD, we have
 $PS=+4$, $QT=+10$, and $KV=+3$, the
 roots of the equation.

Ex. 3.

Let $y^3-13y+12=0$.

13. In this example $p=0$, therefore CD coin-
 cides with AB; and radius $OB = \sqrt[3]{-3q} =$
 $\sqrt[3]{39} = 4.18$; and $-\frac{3r}{q} = -\frac{36}{-13} = 2.77$ the
 distance of ZG above AB. Take arch
 $BP = \frac{1}{3}$ arch BG, and make $PQ=QK=KP$;
 and let fall perpendiculars on AB, then
 $PS=+1$, $QT=+3$, and $KV=-4$, the
 three roots required.
14. Cubic equations may also be construed by
 a cubic parabola and a right line. Let EVAC
 be a cubic parabola, whose latus rectum is 1.
 Draw





Draw VE the tangent at the vertex, perpendicular to the axis VS, and BI parallel to it, and SBC perpendicular to AB. Fig. 14.

Then put $VH=b$, $VD=c$, VI or $SB=n$, $BC=a$, $VS=x$, then $SC=n+a$, and by the property of the parabola $VS=SC^2$, or

$$x = n+a^2. \text{ By similar triangles, } VH (b) : VD (c) :: SH (x-b) : SC = \frac{cx-bc}{b}; \text{ and}$$

$$BC \text{ or } \frac{cx-bc}{b} = n, \text{ and } ba = cx - cb - nb,$$

$$\text{whence } cx - ba = cb + nb, \text{ or } x - \frac{b}{c} a = b + \frac{nb}{c},$$

$$\text{that is (expunging } x) n^2 + 3n^2 a + 3na^2 + a^2 - \frac{b}{c} a = b + \frac{nb}{c}, \text{ which reduced is}$$

$$a^2 + 3na^2 + 3na + n^3 = 0.$$

$$-\frac{b}{c} a - b$$

$$-\frac{nb}{c}$$

Let $a^3 + pa^2 + qa + r = 0$, be any cubic equation. By comparing them, and equating the like terms; we have $3n=p$, and $n=\frac{1}{3}p$. Also

$$3nn - \frac{b}{c} = q, \text{ and } \frac{b}{c} = 3nn - q = \frac{1}{3}pp - q.$$

Again, $n^2 - b - \frac{bn}{c} = r$, or $\frac{1}{9}p^2 - b - \frac{1}{3}p$

$$\times \frac{1}{3}pp - q = r, \text{ whence } b = \frac{1}{3}pq - \frac{2}{9}p^2 - r.$$

$$\text{And since } \frac{b}{c} = \frac{1}{3}pp - q, \text{ } c = \frac{b}{\frac{1}{3}pp - q} = \frac{pq - \frac{2}{9}p^2 - r}{\frac{1}{3}pp - q}.$$

Whence we have the following construction, by this

CONSTRUCTION of B. I.

2 R U L E.

Given the equation $a^3 + pa^2 + qa + r = 0$.

1. With the parameter 1, and the axis VS, describe the cubic parabola FVAC, draw the diameter RAB, distant $\frac{1}{3}p$ from the axis VS, to the right hand, if affirmative; and draw the tangent at the vertex IVD.
2. In the axis VS take $VH = \frac{1}{3}pq - \frac{2}{27}p^3 - r$, downwards, if affirmative.
3. In the tangent IVD, take $VD = \frac{VH}{\frac{1}{3}p - q}$
 $= \frac{pq - \frac{2}{27}p^3 - 3r}{pp - 3q}$, to the left, if affirmative.
4. Through the points D, H, draw the right line FDHC, to cut the parabola. From all the points of intersection, let fall perpendiculars on the diameter AB, which will be the roots of the equation; those on the right hand of AB affirmative; these on the left, negative.
5. When any of the aforesaid quantities are negative, they must be laid the contrary way to what is directed above (Art. 1, 2, 3).

SCHOLIUM.

If the second term be wanting, $p = 0$, and AB coincides with VS; and then $VH = -r$,

and $VD = \frac{r}{q}$.

If the numbers given in the equation, be too great for your parabola; the equation is easily changed into another with less numbers, by Prob. xlii.

Secl. IX. EQUATIONS.

309
Fig.

Ex. 4.

Let the equation be $a^3 + 1.8a^2 - .5125a - 1.05 = 0$.

$$\frac{1.05}{r} = 0.$$

The parabola being described, we have $VI = \frac{1}{2}p = .6$, the distance of IB from the axis VS, on the right; and $VH = \frac{1}{2}pq - \frac{1}{2}p^2 - r = .3105$, taken downwards from the vertex V.

And $VD = \frac{VH}{\frac{1}{2}pp - q} = .195$, on the left from V.

Through D, H draw the right line FDHC, cutting the parabola in F, G, C; from which points letting fall perpendiculars on AB, we have $FR = -1.75$, $GL = -.8$, and $CB = +.75$, the three roots of the equation.

Ex. 5.

Let $x^3 - 4x + \frac{2}{3} = 0$.

Here $p = 0$, and therefore AB coincides with VS; then make $VH = -r = -\frac{2}{3}$, upward; and $VD = \frac{r}{q} = -\frac{2}{3}$, to the right hand.

Then through H, D draw the line FDC, to cut the parabola, in F, G, C; from which letting fall perpendiculars on VS, we have $FR = -1\frac{1}{2}$, $GL = +\frac{1}{2}$, and $CB = +1$, the three roots required.

Cubic and biquadratic equations may also be constructed by the common parabola. Let FVAC be a parabola, VS its axis, AB a diameter parallel to it. EA, and SBC two ordinates perpendicular to VS. Draw also HD perpendicular, and HQ parallel to VS; draw HC.

210 CONSTRUCTION of B. I.
 Fig. 16. Put EA or SB = c , AD = d , DH = g ,
 HC = f , and BC = a . Then QC = $g + a$, la-
 tus rectum of the figure = 1.

Then by the property of the figure AB =
 FB × BC = $2ca + aa$, and DB or HQ = $2ca +$
 $aa - d$, and $HC^2 = HQ^2 + QC^2$, that is, $ff =$
 $a^2 + 4ca^3 + 4ccaa + dd - 4cad - 2daa + gg + 2ga$
 $+ aa$; which reduced is,

$$\begin{aligned} a^2 + 4ca^3 + 4ccaa + dd - 2d - 4dc + dd \\ + 1 - ff = 0. \end{aligned}$$

Let $a^2 + pa^3 + qa^2 + ra + s = 0$, be any
 biquadratic equation; compare this term by
 term with the other; to find the values of
 the quantities c, d, f, g . Then we have
 $4c = p$, and $c = \frac{1}{4}p$.

Again, $4cc - 2d + 1 = q$, and $2d = 4cc + 1 -$
 $q = \frac{1}{2}pp + 1 - q$, and $d = \frac{pp + 1 - q}{2}$.

$$\begin{aligned} \text{Again, } 2g - 4cd = r, \text{ and } g = \frac{4cd + r}{2} \\ = \frac{pd + r}{2}. \end{aligned}$$

Lastly, $gg + dd - ff = s$, and $ff = gg + dd - s$.
 From hence arises the following construction.

3 R U L E.

Having the equation,

$$\begin{aligned} a^2 + pa^3 + qa^2 + ra + s = 0. \\ \text{or } a^2 + pa^2 + qa + r = 0. \end{aligned}$$

1. Describe a parabola FVAC, whose pa-
 rameter is 1, and axis VS. Draw the diame-
 ter AB at the distance of $\frac{1}{2}p$ from the axis,
 on the right hand, if p is affirmative. Then
 for the central rule.

2. From







Fig. 13.




14



15



16



124

2. From A, the top of the diameter, take Fig.
16.
 $AD = \frac{\frac{1}{2}pp + 1 - q}{2}$, downwards, if affirmative.

3. From D in the perpendicular DH, take
 $DH = \frac{p \times AD + r}{2}$ towards the left, if affirmative.

4. But when any of these quantities are negative, set them the contrary way.

5. From the center H with the radius $\sqrt{AD^2 + DH^2 - s} = \sqrt{HA^2 - s}$, describe a circle which will cut the parabola in several points as C.

6. From the points of intersection, let fall perpendiculars upon the diameter AB, and these will be the roots of the equation; these (BC) on the right side of AB, are affirmative roots; and on the left side, negative. And there are always as many real roots, as there are points of intersection; and the rest are impossible.

SCHOLIUM.

If the second term be wanting; then $p = 0$, and the diameter AB coincides with VS. Then

also $AD = \frac{1 - q}{2}$, and $DH = \frac{1}{2}r$.

In cubics s is wanting, and then the radius HC becomes $= HA$.

If the numbers or coefficients be too large for your parabola, you must transform the equation, into another to suit your parabola, by Prob. xlii. and then construct it; and lastly, restore the true roots.

Fig.

Ex. 6.

Suppose $y^3 + 20y^2 - 500y - 6000 = 0$.

The numbers being too large, put $x = \frac{1}{10}y$, or $y = 10x$; then the equation becomes $1000x^3 + 2000x^2 - 5000x - 6000 = 0$, that is, $x^3 + 2x^2 - 5x - 6 = 0$, where the numbers are small.

17. The parabola FVC being described, make $EA = \frac{1}{2}p = \frac{1}{2}$, on the right, and draw AB parallel to the axis VS.

Make $AD = \frac{\frac{1}{2}pp + 1 - q}{2} = \frac{2 + 5}{2} = 3\frac{1}{2}$, downwards. Draw DH perpendicular to AD, and make $DH = \frac{p \times AD + r}{2} = \frac{2 \times 3\frac{1}{2} - 6}{2} = \frac{1}{2}$, to the left. From the center H, with radius HA, describe a circle, cutting the parabola in R, A, C, F; from which letting fall perpendiculars on AB; we have $RA = -1$, $BC = +2$, $FB = -3$, the roots of the equation $x^3 + 2x^2 - 5x - 6 = 0$, and multiplying by 10, we have -10 , $+20$, and -30 , for the roots of the given equation $y^3 + 20y^2 - 500y - 6000 = 0$.

Ex. 7.

Let $x^4 - 1.75x^3 - 4.625x^2 + 4.875x + 6.75 = 0$.

18. Describe the parabola FVC, and draw the axis VS, and make $EA = \frac{1}{2}p = -.44$, to the left, and draw AB parallel to the axis, make $AD = \frac{\frac{1}{2}pp + 1 - q}{2} = 3.19$ downwards. Draw DH perpendicular to AB, and make $DH = \frac{p \times AD + r}{2} = -.36$, to the right.

From

Sect. IX. EQUATIONS. 313

From the center H, with the radius $\sqrt{AD^2 + DH^2} = 0.75 = 2$, describe a circle cutting the parabola in F, R, G, C; from which drawing perpendiculars to AB, we have $RO = -1$, $FB = -1\frac{1}{2}$, $GI = 2$, $CP = 2.25$, the roots required. Fig. 18.

Ex. 8.

Given the equation $x^4 - 1.5x^3 + 5x^2 - 9x - 6 = 0$.

Describe the parabola RVC to the axis VS, and make $EA = p = -0.375$, to the left, and draw AB parallel to VS. Take $AD = \frac{pp + 1 - q}{2} = -1.72$ upwards. Then (per-

pendicular to AD) take $DH = \frac{p \times AD + r}{2} = -3.21$, to the right. With the center H, and radius $4.39 (= \sqrt{AD^2 + DH^2 + 6})$, describe a circle, to intersect the parabola; from the points of intersection, letting fall perpendiculars on AB, gives the roots, $RO = -5$, and $CB = +2$. The other two roots are impossible, which is known from this, that the circle intersects the parabola in no more points than these two.

Ex. 9.

Let $x^4 - 5.67x^2 + 3.864 = 0$.

Here $p = 0$, therefore describe the parabola FAC, whose axis is AS; and make $AD = \frac{1 - q}{2} = 3.335$, downwards; and $DH = r = 0.403$, perpendicular to AD, to the left. With radius $\sqrt{AD^2 + DH^2} = 3.864 = 2.72$, describe a circle cutting the parabola, in R, G, C, F, and

Fig. and the perpendiculars from these points upon AS, give $RE = -.8$, $GI = +1$, $CB = +2.1$, and $Fs = -2.3$, the roots of the equation.

S C H O L I U M.

Geometrical equations may be constructed by lines as well as by numbers. For proper lines may be found for the coefficients, by proceeding according to Prob. xcvi; and so the whole may be done geometrically.


Quadratic equations, whose general form is $a^2 + pa + q = 0$, may also be constructed by the last rule; and then r and s will be $= 0$; but the method of constructing by the circle, is easier.


4 R U L E.

Any cubic or biquadratic equation $x^4 + px^3 + qx^2 + rx + s = 0$, may be constructed mechanically thus:

21. 1. Upon a plain smooth wall, draw a horizontal line AB, and CD perpendicular to it, and take $CP = \frac{1}{2}p$, to the left hand if p is affirmative. Hang a thread and plummet EPE to any point E, in the perpendicular EP; make a knot in the thread at n , and tie the other end so to the fixt point E, that Pn may be $= \frac{1}{2}$. Then with a pin or the point of a compass, move the thread EF sideways toward CD, till the knot n falls in the point C; mark the point D in the line CD, where the pin is, when that happens.

2. From D take $DG = \frac{pp + 1 - q}{2}$ ($= d$), downwards, if affirmative. And in the perpendicular GH, take $GH = \frac{dp + r}{2}$, to the left, if





Sect. IX. EQUATIONS.

if it is affirmative. But if any of these quantities be negative, they must be taken the contrary way, to what is directed above.

3. Then with the radius or distance $\sqrt{HD^2 - s}$, and one foot of the compasses in H, move the other foot along with the thread, round in a circle, and the weight F will ascend and descend, as the thread EF moves laterally. Observe always, when the knot n falls in the line AB, and mark all these points, Q, N, O, R. Then the distances of these points from C, are the roots of the equation; the affirmative on the right, the negative on the left hand of C; thus RC is an affirmative root, and QC, NC, OC, negative ones.

It is plain, this rule is founded on the last. For the moving point of the compass is always in the curve of a parabola, when the point n is in the line AB. To prove which, suppose the parabola ADB, to be described, whose focus is E. Then by the property of the figure, $EL + LR = EP + \frac{1}{2}$ parameter = $EP + Pn$ or $En = ED + DC$. Therefore the circle cuts the parabola in L; and the distance of L from DC, that is RC is one root of the equation; and the like for the rest.

S E C T. X.

Rules and Directions for the investigation and solution of Problems.

P R O B L E M X C I X.

To find if a question be truly limited.

A Question is said to be truly limited, when it admits but of one solution; or at most, of as many as is the index of the highest power of the unknown quantity in the final equation. And whether a question be limited or not, may be known from the equations, by this

R U L E.

When the number of unknown quantities, is just as many as the number of given equations, not depending on one another; then the question is truly limited.

But when the number of unknown quantities exceeds the number of equations given; then the question is unlimited, and capable of innumerable answers.

And when the number of unknown quantities is less than the number of given equations; then the question is absurd and impossible, except these equations be dependent upon one another; in which case the dependent ones may be struck out.

Equations are said to be *dependent* on one another, when they may be formed or derived from one another, by any operations, with the help of the known axioms.

For

For by Cor. 1, 2. Prob. liii, one unknown quantity may be taken away by each equation; so that at last there will remain but one equation, and one unknown quantity in it; and therefore it is truly limited.

But if there were more unknown quantities than equations, there will remain more unknown quantities than one, in the last equation. And then the question is not limited; for all of them, but one, may be taken at pleasure: and this is the reason of questions being unlimited.

Lastly, if there be more equations than unknown quantities, then at last there will remain one unknown quantity for several equations; and then the question is more than limited; and will therefore be impossible. For the unknown quantity being exterminated, there will be an equation consisting of all known quantities; which must be contradictory to one another, except they were some way or other depending on one another, so as to make an equality.

S C H O L I U M.

As a problem is truly limited, when the number of independent equations, is equal to the number of unknown quantities: so likewise a problem is truly limited, though there be never so many equations, provided all, above that number, are depending upon these, and derived from them. This is plain from any algebraic process; for in the operation, all the succeeding equations, are derived from these, first given; and all equations so derived, make no alterations in the limitation of the problem.

A problem may be impossible and more than limited, though the number of equations be less than the number of unknown quantities; and that is when the equations are contradictory.

As

As if $a + e + 2y = b$,

And $2a + 2e + 4y = c$; a, e, y being unknown quantities, and b, c , known ones. Now if it happen that $c = 2b$, the problem is unlimited; but if c is not $= 2b$, then the problem is impossible.

And therefore in general, problems are absurd, when the equations given are derived from absurd equations, or may be reduced to such: even though the number of equations be equal to or less than the number of unknown quantities.

The equations given in a problem, ought to be independent, otherwise they will either be consequential, or contradictory to one another. In the first case, you will at last find some quantity equal to itself. And in the second case you will arrive at some absurdity, where a greater quantity is equal to a less. And it often happens, that at the end of an operation, the equations given, are found to be either dependent or inconsistent with one another; which at first, could not so easily be discovered.

P R O B L E M C.

To investigate an algebraic problem.

R U L E S.

I. When a question is proposed to be resolved algebraically; the first thing to be done, is to consider the nature and circumstances thereof, to find out what is given therein, and what required. And the nature and tenor thereof being clearly understood; reject every condition or circumstance, which has no necessary connection or relation with the thing enquired after. Then give names to all the quantities concerned in the calculation, whether given or sought; that is, for the several numbers or quantities, or at least for the principal of them, put so many different letters, as directed in the notation,

notation; taking care to make the same letter stand invariably for the same thing, throughout the whole operation.

And in general problems, it will be convenient to make choice of such letters or symbols, as may some way represent to the mind, the things designed by them; as r for radius, s for sine, l for latus rectum, v for velocity, t for time, &c.

And if there be never so many quantities of different sorts, we may represent them by any numbers we like; or even all of them by 1, which is the most simple notation. Thus we may call any degree of motion 1, any degree of velocity 1, and we may put 1 for any quantity of space, time, matter, &c. But then we must take care to represent other quantities of the same sort, by proportional numbers.

We can also measure any kind of quantity by any other kind of quantity, by taking parts or degrees of one sort, proportional to the parts or degrees of the other. Thus, quantities of force may be measured by right lines proportional to them; bodies or quantities of matter by their weights; velocities by the spaces described in equal times; and all sorts of quantities or things by numbers.

2. But as that solution of a question is reckoned the more artificial, the fewer unknown quantities are assumed at first. Therefore when the principal quantities are denoted by letters; some of the quantities, that may be easily derived from the rest, are left without a name. As when the whole is given and a part, the other part is easily had from thence; or the parts being given, you may find the whole. Also when two sides of a right-angled triangle are denoted in algebraic terms, the third side is had from these, by addition or subtraction of squares. Likewise three terms of a proportion being given, the fourth term is easily derived

derived from these three; and in all such like cases, where the values of some are easily derived from the rest. And by this means, there will be fewer terms to exterminate.

3. After the designation of the quantities, by letters, as aforesaid; we must next abstract it from words, and translate it out of the English into the Algebraic language: that is, we must denote all the conditions of it, by so many algebraic equations: and this is called *stating the question*. In order to this, we must suppose the thing done which was required; and then, without making any difference between the known and unknown quantities; assume any of them, known or unknown, to begin your computation from; taking such as you think will bring out the simplest equation, or give the easiest solution. And it is best to assume that quantity to begin with first, which is easiest found or brought to an equation. And therefore it is often more convenient, not to begin with that which is directly required, but with some other, from which the quantity required may be easily had.

From these first assumed quantities, you must proceed in a synthetic method to find other quantities wanted, and from these to find others, &c. according as the nature of the question directs, till you get what equations you want. To this purpose, you must attend strictly to the nature, design, and meaning of the question, and search into all the circumstances of it, and examine into the particular relations of the quantities to one another; so that from thence you may get a proper number of equations. But sometimes these equations cannot be had from the words of the question; but depend upon the hidden properties of the quantities concerned therein; and then the equations are to be deduced from them, by a proper chain of reasoning.

reasoning, according to the nature of the subject under consideration. Thus, in numerical questions, we must proceed by the properties of numbers: in geometrical problems, by the elements of geometry: in mechanical problems, by the principles of mechanics: in trigonometrical problems, by the rules of trigonometry: in philosophical problems, by the laws of motion; and so of other subjects. And here great care must be taken that your equations do not depend upon one another; and that there be as many as there are unknown quantities, otherwise the question will not be limited.

4. Having got a proper number of equations, our business is now, to exterminate them one by one, as fast as we can, till there only remains one unknown quantity, in one final equation: then the problem is said to be *brought to a solution*. And by these equations, you must exterminate these quantities first, that are most easily exterminated; that is, the simplest first, and so on; till the quantity that remains at last, may give the simplest equation possible; or more simple than if any other of the unknown quantities remained in the final equation. And in all your process, great care must be taken, to keep to a just equality; which will certainly be, if you observe all along, to work according to these just rules or axioms, at the beginning of this book.

5. As to the chusing fit terms or quantities to begin the calculation with; it sometimes happens that there is such a relation of two terms of the question, when compared with the rest, that in making use of either of them, they will bring out equations exactly alike; or that both, if they are made use of together, shall bring out the same final equation, as to form. Then it will be the best way to make use of neither of these terms; but instead
Y
thereof,

thereof, to chuse some third, which has a like relation to both. As suppose the half sum or half difference, or perhaps a mean proportional; or any other quantity related to both indifferently, and without a like.

6. The proper designation of the terms will often much abridge the operation. As if two numbers are sought, whose sum or difference (n) is given, it will be convenient to take $\frac{1}{2}n+a$, and $\frac{1}{2}n-a$, for the numbers.

Also when several numbers are sought in arithmetical progression, where the common difference (d) is given; we may properly put $a-d$, a , $a+d$, for the numbers, when there are three: or $a-\frac{1}{2}d$, $a-\frac{1}{2}d$, $a+\frac{1}{2}d$, $a+\frac{1}{2}d$, for the numbers, when four are required; and so on.

Again, if several numbers are sought in geometrical progression; put aa , ae , ee , for three numbers: and a^3 , a^2e , ae^2 , e^3 , for four numbers: and a^4 , a^3e , a^2e^2 , ae^3 , e^4 , for five numbers; and so on. Or denote them by such other series, as will give them all, with the fewest letters.

7. Sometimes problems will run up into very high equations, where the unknown quantities cannot be expunged without great difficulty. Therefore, in such a case, if you can substitute new letters for the sums, or products, or powers, &c. of some of the old quantities; and then expunge all these old ones, and get a proper number of equations; you may often find the value of these new quantities, by easy and low equations; from whence the old quantities may be more easily determined. And you must find these new quantities by trials, such, that when they are substituted, they may render the equations easier. See Prob. xxiv, xxv. B. II.

Likewise in any operation, when you have a multitude of unknown quantities, for the coefficient of any power of the unknown quantity; put a single letter

letter for them all, which will much abridge the operation.

8. In geometrical problems, there is often more labour and skill required, than in numerical ones. In these you must first draw a figure, according to what the question requires to be done. And then it is often requisite to produce right lines; or to draw lines parallel or perpendicular to other lines; and to certain points, or through certain points; or to make similar triangles, and such like; all preparatory for the solution of the problem: always endeavouring to resolve the scheme into similar triangles, or right-angled ones, or given ones. Then assume such a line, &c. for your unknown quantity, as you judge will bring out the simplest equation. For you may begin your computation with any quantity, known or unknown: which done, you must proceed synthetically to find the rest. In general, let these quantities be denoted by letters, that lie nearest the given parts of the figure, and by means of which other parts adjoining may be easily had, without surds. In triangles, draw a perpendicular from the end of a given side, and opposite to a given angle. Such preparations as these being made, just as you find necessary for the method of solution you intend to try; pursue your computation according to the nature and property of lines, and the conditions given in the question, proceeding from the quantities assumed, to other quantities, as the relation of the lines direct; till you get two values for one and the same quantity, or find one quantity denoted two different ways, by which you will get an equation. The general principles for carrying on the computation are such as these; the addition or subtraction of lines, to find the sum or difference. The proportionality of lines (arising from similar triangles), where three terms being given to find a fourth. The addition or subtraction of squares in right-angled triangles, where

two sides being given, the third may be found. Likewise the doctrine of proportion will be of frequent use. Besides we must make use of such propositions in geometry as are suitable to the purpose; such as B. I. prop. 1, 2, 4, 8, 10, 11. B. II. 2, 3, 10, 13, 14, 15, 18, 21, 24, 25, 26. B. III. 1, 6, 7, 17, 20. B. IV. 9, 12, 13, 14, 16, 17, 20, 27, 31; and some in the following books, as occasion requires. By help of these principles, and a chain of right reasoning, we shall obtain as many equations as unknown quantities, which being had, we must change our method, and exterminate the superfluous quantities, and find the root of the final equation.

9. If the method you go upon at first, for the solution of the problem, proceeds but badly, as running into high equations and surds. You must draw fresh schemes, and begin your computation anew, till you have hit on a method as elegant as possible. For the principal art, of resolving problems, is to frame the positions with such judgment, that the solution may end in as simple an equation as possible. For some methods will produce more intricate equations and solutions, than others. But the skill of finding out the most simple and easy ways of resolution is not to be prescribed by any rules, but is only attained by constant practice and experience.

10. If you have any doubt what quantity to take for the quantity sought, so as to bring out the simplest equation. Suppose you have got a final equation with x ; take some other quantity y , which you suspect may be as simple, seek an equation between x and y ; then if y be of as high a power as x ; the final equation, if y were used, would be as high as is the final equation with x .

Or, Having got an equation between x and y ; substitute for x its value in terms of y , in the said final equation with x ; and you will find what power

er y will arise to, without forming the process anew for y . But if the equation between x and y be not a simple equation; it will often be as well to begin the process anew for y .

Or, If there be several quantities, and you do not know which will bring out the simplest equation. Put letters for them all, and get as many equations. Then by expunging such as are most easily expunged; you will, for the most part, get the most simple equation.

11. Lastly, when the final equation is obtained, extract its root by Sect. VIII, and you have the answer in numbers.

Note, The numbers given in a question, cannot always be taken at pleasure, but must often be subject to one or more determinations or restrictions, which for the most part are discoverable by the theorem resulting from the resolution of the question.

12. When you have an equation containing the quantity sought; and the equation is also effected with a second unknown quantity, which you want to get rid of; the extermination of which runs you to a very high power. Now if it happens that this second unknown quantity, is but in a few of the terms, which are but small in respect of the rest. Then if you can nearly guess at its value, you may retain it in the equation, putting that value for it, which will make little difference in the equation, among so many quantities, if you miss its value a little. Then the root of the equation being extracted will give the other unknown quantity very near. And this being had, the second unknown quantity will then be found more exactly, and may be substituted for it again, and the operation repeated, &c.

And one may often guess nearly at the value of this second quantity, from the conditions of the problem; especially if it be a geometrical one, from the construction of the figure.

These sorts of equations may also be resolved by the Rule of False Position, as directed in Pr. xcii. Rule 5; as also by Prob. xcv.

13. When you want to compute a problem for some practical use in common life, but by pursuing it in its mathematical rigour, you fall upon some irresolvable equations or intricate furd or series. Then you may often resolve it on very simple principles, by neglecting such quantities or such conditions, as serve only to embarrass the problem, but make little or no difference for the purpose you want it. In such case, neglect such quantities or such conditions, as are of little moment; and instead of such quantities as make the calculation difficult, take others nearly equal to them, which will make the operations more simple, or as simple as possible. Or some of the least moment may be entirely left out. And thus one may come at an easy solution of the problem.

These are the general rules of working; all which will be made clear, by the examples in the following Book II.

B O O K II.

The Solution of Problems.

A *Problem* is a question proposed to be resolved; and the *Solution* of a problem, is the finding such numbers, lines, &c. as will fulfil the conditions of the question.

Of problems these are *determined*, that have a determinate number of answers; and *indetermined*, which have innumerable answers.

Problems are of several kinds, as numerical, geometrical, trigonometrical, philosophical, mechanical, &c.

A problem of one, two, three, &c. dimensions, is that which has one, two, three, &c. solutions or answers.

We have hitherto been laying down such rules, as are necessary for the investigation and solution of problems. The reader must take particular care, to make himself well acquainted with these rules, and keep them in mind, so that he may have them ready for use, upon all occasions; for without them no problem can be solved. But as precepts are but of little use without examples, and generally reach no farther than mere speculation; I shall therefore, in the next place, apply them to practice, and that in the solution of a great variety of problems, in the most material branches of the mathematics; which I shall now begin with directly.

S E C T. I.

Numerical Problems.

P R O B. I.

There are two numbers whose sum is 25, and the proportion of one to the other is as 2 to 3, what are the numbers?

Suppose	}	1	$a =$ greater.
		2	$e =$ lesser.
per quest.		3	$e : a :: 2 : 3$
3 ×		4	$2a = 3e$
4 ÷ (2)		5	$a = \frac{3}{2}e$
per quest.		6	$a + e = e + \frac{3}{2}e = 25$
6 × (2)		7	$2e + 3e = 5e = 50$
7 ÷ (5)		8	$e = \frac{50}{5} = 10$, the lesser.
5,		9	$a = \frac{3}{2}e = 15$, the greater.

Otherwise thus,

Suppose		1	$a =$ greater, $s =$ sum, $= 25$; then $s - a =$ lesser.
per quest.		2	$2 : 3 :: s - a : a$
2 ×		3	$2a = 3s - 3a$
3 transf.		4	$5a = 3s$
4 ÷ (5)		5	$a = \frac{3s}{5} = \frac{75}{5} = 15$, the greater number.
		6	$s - a = 10$, the lesser.

P R O B.

P R O B. II.

A man having a certain number of pence, gives to A $\frac{1}{2}$ of them, to B $\frac{1}{4}$, to C $\frac{1}{8}$, and to D $\frac{1}{11}$, and then had 3 remaining. How many had he at first?

Let	1	$a =$ number of pence he had.
per quest.	2	$\frac{1}{2}a + \frac{1}{4}a + \frac{1}{8}a + \frac{1}{11}a = a - 3$
2,	3	$\frac{23}{88}a = a - 3.$
3 transf.	4	$\frac{1}{88}a = 3$
4 \times	5	$a = 72.$

P R O B. III.

A man hired a labourer, on condition, that for every day he wrought, he should have 12 d. and for every day he idled, he should forfeit 8 d. After 390 days, neither of them was in debt. To find the number of work days and play days.

Let	1	$a =$ the working days, $b = 390$; then $b - a =$ the play days.
per quest.	2	$12a = 390 - a \times 8 = 3120 - 8a,$
2 transf.	3	$20a = 3120,$
3 $\div (20)$	4	$a = \frac{3120}{20} = 156,$ the work days.
1,	5	$ba = 234,$ the playing days.

PROB. IV.

Some young men and maids had a reckoning of 37 shillings; and every man was to pay 3 shillings, and every maid 2; now if there had been as many men as maids, and maids as men, the reckoning would have come to 4 shillings less. What is the number of each?

Suppose	1	$a = \text{men}, e = \text{maids}, b = 37, c = 4.$
per quest }	2	$3a + 2e = b$
	3	$2a + 3e = b - c$
	4	$9a + 6e = 3b$
$2 \times (3)$	5	$4a + 6e = 2b - 2c$
$3 \times (2)$	6	$5a = b + 2c$
$4 - 5$	7	$a = \frac{b + 2c}{5} = 9, \text{ the men.}$
$6 - (5)$	8	$2e = b - 3a$
$2 - 3a$	9	$e = \frac{b - 3a}{2} = 5, \text{ the women.}$
$8 \div (2)$		

PROB. V.

A man being asked what a clock it was? answered, that it was between 8 and 9; and that the hour and minute hands were both together.

Let	1	$x = \text{time required}, b = 8, c = 12,$ $d = 1.$
	2	since the two hands divide the hour, and the whole circumfe- rence in the same proportion, therefore $c : x :: d : x - b,$
$2 \times$	3	$cx - cb = dx$
3 transf.	4	$cx - dx = cb$
$4 \div$	5	$x = \frac{cb}{c - d} = \frac{96}{11} = 8 \frac{8}{11} \text{ h. } 43 \frac{38}{11} \text{ min.}$

PROB.

PROB. VI.

A man gives the first beggar he meets with, $\frac{1}{6}$ of the pence he had and 4 d. more: to the second $\frac{1}{6}$ the remaining pence and 8 d. more: to the third $\frac{1}{6}$ the remaining pence and 12 d. more, and so on, increasing 4 d. every time, till at last he had nothing left; and then all the beggars had equal shares. Query, the number of pence and beggars.

Suppose	1	$a =$ pence he had at first.
per quest.	2	$\frac{a}{6} + 4 =$ pence given to the first beggar.
1-2	3	$\frac{5}{6}a - 4 =$ remainder.
$3 \div (6)$	4	$\frac{5}{36}a - \frac{4}{6} = \frac{1}{6}$ the remainder
$4 + (8)$	5	$\frac{5}{36}a - \frac{4}{6} + 8 =$ pence given to the second beggar.
$2 = 5$	6	$\frac{a}{6} + 4 = \frac{5}{36}a - \frac{4}{6} + 8,$ per quest.
$6 \times (36)$	7	$6a + 144 = 5a - 24 + 288$
7 transf.	8	$a = 120.$
2,	9	$20 + 4 = 24 =$ share of each.
8, 9	10	$\frac{120}{24} = 5 =$ number of beggars.

P R O B. VII.

There are three numbers, the first with $\frac{1}{3}$ the other two makes 14, the second with $\frac{1}{4}$ the other two makes 8; the third with $\frac{1}{5}$ the other two makes 8. What are the numbers?

Let	1	a, e, y be the numbers.
per quest	2	$a + \frac{e+y}{3} = 14$
	3	$e + \frac{a+y}{4} = 8$
	4	$y + \frac{a+e}{5} = 8$
	$2 \times (3)$	5
$3 \times (4)$	6	$4e + a + y = 32$
$5 - 6$	7	$2a - 3e = 10$
$5 \times (5)$	8	$15a + 5e + 5y = 210$
$4 \times (5)$	9	$a + e + 5y = 40$
$8 - 9$	10	$14a + 4e = 170$
$7 \times (4)$	11	$8a - 12e = 40$
$10 \times (3)$	12	$42a + 12e = 510$
$11 + 12$	13	$50a = 550$
$13 \div (5)$	14	$a = 11$
7 transf.	15	$3e = 2a - 10$
$15 \div (3)$	16	$e = \frac{2a - 10}{3} = 4$
5 transp.	17	$y = 42 - 3a - e = 5.$

P R O B. VIII.

Having given the sum of two numbers 8, and the difference of their squares, 16; to find the numbers.

Let	1	$x =$ lesser number, $a = 8, b = 16.$
	2	$a - x =$ greater number.

1	2	3	$xx =$ square of the lesser.
2	2	4	$aa - 2ax + xx =$ square of the greater
4	3	5	$aa - 2ax = b$ per quest.
5	transf.	6	$2ax = aa - b$
6	$\div 2a$	7	$x = \frac{aa - b}{2a} = 3$, the lesser.
2,		8	$a - x = 5$, the greater.

PROB. IX.

There are three numbers, the sum of the first and second is 9, of the first and third 10, of the second and third 13. What are the numbers?

Let	1	x, y, z , be the numbers, $a=9$,
		$b=10, c=13$;
per quest	2	$x + y = a$
	3	$x + z = b$
	4	$y + z = c$
	5	$y = a - x$
2 - x	6	$z = b - x$
3 - x	7	$a - x + b - x = c$
4, 5, 6	8	$2x = a + b - c$
7 transf.	9	$x = \frac{a + b - c}{2} = 3$
8 $\div (2)$	10	$y = a - x = 6$
5,	11	$z = b - x = 7$
6,		

PROB. X.

Two travellers A and B, 360 miles distance, set out at the same time. A travels 10 miles an hour; B 8. How far does each travel before they meet?

Suppose	1	A travels x miles, then B travels $360 - x$.
by propor.	2	$x : 360 - x :: 10 : 8$
2 x	3	$8x = 3600 - 10x$

3 transf.

3 transf.	4	$18x = 3600$
$4 \div (18)$	5	$x = \frac{3600}{18} = 200$, A's journey.
1,	6	$360 - x = 160 =$ B's journey.

P R O B. XI.

If three agents A, B, C, can produce the effects a , b , c , in the times e , f , g , respectively. In what time will they all jointly produce the effect d ?

Let	1	$x =$ time sought.
by pro- portion	2	$e :: x :: a : \frac{ax}{e}$, A's effect in the time x .
		$f :: x :: b : \frac{bx}{f} =$ B's effect in time x .
		$g :: x :: c : \frac{cx}{g} =$ C's effect in time x .
2, 3, 4,	5	$\frac{ax}{e} + \frac{bx}{f} + \frac{cx}{g} = d$
5 reduced	6	$x = \frac{d}{\frac{a}{e} + \frac{b}{f} + \frac{c}{g}} =$ time required.

P R O B. XII.

A woman can buy apples at 10 a penny, and pears at 25 for 2 pence; if she lay out $9\frac{1}{2}$ pence for 100 apples and pears together. How many of each must she have?

Let	1	$a =$ apples, then $100 - a =$ pears.
by pro- portion	2	$10 : 1 d. :: a : \frac{a}{10}$, price of the apples.
		$25 : 2 d. :: 100 - a : \frac{200 - 2a}{25}$, price of the pears.

per

per quest.	}	4	$\frac{a}{10} + \frac{200-2a}{25} = 9\frac{1}{2}$
4 × (50)		5	$5a + 400 - 4a = 475$
5 transp.		6	$a = 75$, the apples:
		7	$100 - a = 25$ the pears.

PROB. XIII.

A vintner would mix wine at 10 d. the quart, with another sort at 6 d; to make a 100 quarts to be sold at 7 d. How much of each must be take?

Let	}	1	$a =$ quarts of 10 penny, $e =$ quarts of 6 penny, $b = 10$, $c = 6$, $m = 100$, $f = 7$.
by proportion		2	$1 : b :: a : ba$, value of a quarts.
		3	$1 : c :: e : ce$, value of e quarts.
per qu.		4	$ba + ce = mf$.
		5	$a + e = m$
5 — a		6	$e = m - a$
4, 6		7	$ba + cm - ca = mf$
7 tr.		8	$ba - ca = fm - cm$
8 ÷		9	$a = \frac{fm - cm}{b - c}$
6,		10	$e = \frac{bm - fm}{b - c}$

PROB. XIV.

A factor exchanged 6 French crowns and 2 dollars for 45 shillings; and 9 French crowns and 5 dollars for 76 shillings. What is the value of a French crown and a dollar?

Suppose	}	1	$x =$ French crowns, $y =$ dollars,
			$a = 6$, $b = 2$, $d = 9$, $e = 5$, $c = 45$, $f = 76$.

per

per qu.	}	2	$ax + by = c$	
		3	$dx + ey = f$	
		$2 \times e$	4	$eax + cby = ec$
		$3 \times b$	5	$bdx + eby = bf$
		$4 - 5$	6	$aex - bdx = ec - bf$
		$6 \div$	7	$x = \frac{ec - bf}{ae - bd} = 6\frac{1}{11}$
		2,	8	$\frac{ec - bf}{ae - bd} a + by = c$
		8 tr.	9	$by = c - \frac{ec - bf}{ae - bd} a = \frac{bfa - bdc}{ae - bd}$
			10	$y = \frac{af - cd}{ae - bd} = 4\frac{1}{11}$

PROB. XV.

To divide a number b , into four parts; so that the first being increased with d , the second diminished by d , the third multiplied by d , and the fourth divided by d ; may be all equal.

Let	}	1	a, e, u, y , be the four parts.	
		2	$a + e + u + y = b$,	
		3	$a + d = e - d$	
		4	$a + d = dy$	
		5	$a + d = \frac{u}{d}$	
		$3 + d$	6	$e = a + 2d$
		$4 \div d$	7	$y = \frac{a + d}{d}$
		$5 \times d$	8	$u = ad + dd$
		2, 6, 7, 8,	9	$a + a + 2d + \frac{a + d}{d} + ad + dd = b$
		9 reduced	10	$a = \frac{bd - d^2 - 2dd - d}{dd + 2d + 1}$

6	11	$e = \frac{bd+d^2+2dd+d}{dd+2d+1}$
7	12	$y = \frac{b}{dd+2d+1}$
8	13	$u = \frac{bdd}{dd+2d+1}$

PROB. XVI.

A merchant bought (a) bushels of wheat, (b) bushels of barley, and (c) bushels of oats for (m) pounds. Afterwards he bought (d) bushels of wheat, (e) bushels of barley, and (f) bushels of oats, for (n) pounds.

And after that, (g) bushels of wheat, (h) bushels of barley, and (k) bushels of oats for (p) pounds, each sort at one price. What was each per bushel?

Let	1	$x, y, z,$ be the prices of the wheat, barley, and oats.
per qu. }	2	$ax+by+cz=m$
	3	$dx+ey+fz=n$
	4	$gx+by+kz=p.$
	5	$afkx+bfky+cfkz=fkm$
2xsk	6	$ckdx+ckey+cfkz=ckn$
3xck	7	$cfgx+cfby+cfkz=cfp$
4xcf	8	$afkx-ckdx+bfky-ckey=fkm-ckn.$
5-6	9	$ckdx-cfgx+ckey-cfby=ckn-cfp.$
6-7	10	$Ax+By=C$
substit. 8,	11	$Fx+Gy=H.$
9,	12	$y = \frac{C-Ax}{B}$
10 reduc.	13	$y = \frac{H-Fx}{G}$
11 reduc.	14	$\frac{C-Ax}{B} = \frac{H-Fx}{G}$
12=3	15	$GC-GAx=BH-BFx$
14x		Z

	8	$ac = bx + cbz$
per qu.	9	$df = ex + fez$
	10	$yb = gx + bgz$
8 × ef	11	$acef = efbx + efcbz$
9 × cb	12	$cbdf = cbex + cbefz$
11—12	13	$acef - cbdf = efbx - cbex.$
13 ÷	14	$x = \frac{acef - cbdf}{efb - ecb}$
9 × g	15	$gdf = gex + gfez$
10 × e	16	$eyb = egx + ebgz$
16—15	17	$eyb - gdf = ebgz - gfez$
17 ÷	18	$z = \frac{eyb - gdf}{ebg - efg}$
8 ÷ b	19	$\frac{ac}{b} = x + cz$
19, 14, 18	20	$\frac{ac}{b} = \frac{acef - cbdf}{efb - ecb} + \frac{ceyb - cgdf}{ebg - efg}$
Put	21	$p = f - c, r = b - f, r + p = b - c.$
20, 21,	22	$ac = \frac{acef - cbdf}{pe} + \frac{ceyb - cgdf}{reg} b.$
22 × preg	23	$acpreg = acefrg - cbdfrg + pceybb - pcdfgb$
23 tr.	24	$pceybb = apreg - aefrg + bdfrg + pcdfgb.$
24,	25	$pebby = areg \times p - f + bdfg \times r + p.$
21, 25	26	$pebby = -aregc + bdfg \times r + p.$
26, 21,	27	$pebby = aceg \times f - b + bdfg \times b - c$
27 ÷	28	$y = \frac{aceg \times f - b + bdfg \times b - c}{beb \times f - c}$

P R O B. XVII.

To divide ten thousand into two such parts, that when each of them is divided by the other, the sum of the quotients, may be 5.

Let	1	a, e be the parts, $b=10000, c=5$.
per qu. {	2	$a+e=b$
	3	$\frac{a}{e} + \frac{e}{a} = c$
	4	$e=b-a$.
$2-a$	5	$ee=bb-2ba+aa$
$4 \text{ } \odot \text{ } 2$	6	$cae=aa+ee$
$3 \times ae$	7	$cab-caa=2aa-2ba+bb$
$4, 5, 6$	8	$2+c.a.a-2b+bc.a=-bb.$
$7 \pm$	9	$aa-ba = \frac{-bb}{2+c}$
$8 \div$	10	$a = \sqrt{\frac{bb}{4-c+z} + \frac{b}{2}} = 8273,268$
9 reduc.	11	$e=1726,732.$
4,		

P R O B. XVIII.

A general would range his army in a square battle, but finds he has 284 soldiers to spare; but if he increases the side of the square with one man, he wants 25 to fill up the square. How many soldiers had he?

Let	1	$a =$ side of the square.
per qu. {	2	$aa+284 =$ number of men
	3	$\frac{aa+1}{a+1} - 25 =$ number of men
	4	$aa+284=aa+2a+1-25$
$2=3$	5	$2a=308$
4 tran.	6	$a=154$
$5 \div 2$	7	$aa+284=24000$, the number of men.

P R O B.

P R O B. XIX.

Several persons dining at an inn, the reckoning came to 75 shillings; but two of them sinking away, the rest had 10 shillings a-piece more to pay. Query, the number of persons?

Put	}	1	$a =$ number of persons at first.
per qu.		2	$\frac{175}{a} =$ each man's shot.
		3	$\overline{a-2} \times \overline{\frac{175}{a}} + 10 = 175.$
3 ×		4	$175a + 10aa - 350 - 20a = 175a$
4 transf.		5	$10aa - 20a = 350$
5 ÷		6	$aa - 2a = 35$
6 resolved		7	$a = 7$
2,		8	$\frac{175}{a} = 25,$ the club.

P R O B. XX.

There is a number, consisting of three digits, whose product is 315, and the sum of the first and last is double the second; and that number with 396 added makes a number consisting of the same digits but inverted.

Suppose	}	1	a, e, y the digits.	$a \ e \ y$
		2	$a + y = 2e$	$y \ e \ a$
per qu.		3	$ae y = 315$	
		4	$100a + 10e + y + 396 = 100y + 10e + a,$	
4 ±		5	$99a + 396 = 99y$	
5 ÷ (99)		6	$a + 4 = y$	
2 - a		7	$2e - a = y$	
6 = 7		8	$a + 4 = 2e - a$	

8,	9	$a = e - 2$
6,	10	$y = e + a$
3, 9, 10	11	$acy = e \times e - 2 \times e + 2 = e^3 - 4e = 315$
11 resolv.	12	$e = 7$
9, 10	13	$a = 5, y = 9.$

P R O B. XXI.

To find the value of a , when $\sqrt{a^3} - \sqrt[3]{aa} = 4.962a$.

Put	1	$b = 4.962$
per quest.	2	$a^{\frac{3}{2}} - a^{\frac{2}{3}} = ba$
2,	3	$a^{\frac{9}{6}} - a^{\frac{4}{6}} = ba$
assume	4	$x^6 = a$, then
4,	5	$a^{\frac{9}{6}} = x^9$, and $a^{\frac{4}{6}} = x^4$
3, 4, 5	6	$x^9 + x^4 = bx^6$
$6 \div x$	7	$x^5 - 1 = bxx$
7 transf.	8	$x^5 - bxx = 1$
8 resolv.	9	$x = 1.74256$
4,	10	$a = 27.998$

P R O B. XXII.

Given $\left\{ \begin{array}{l} 1 \quad a^3e^3 + a^2e^3 = 234000 = b, \\ 2 \quad ae + ae^2 + ae^3 = 1860 = c, \\ \text{to find } a \text{ e.} \end{array} \right.$

$$2 \div \quad 3 \quad a = \frac{c}{e + ee + e^3}$$

$$1 \div e^3 \quad 4 \quad a^3 + a^2 = \frac{b}{e^3}$$

$$3, 4 \quad 5 \quad \frac{c^3}{(e + ee + e^3)^3} + \frac{cc}{e + ee + e^3} = \frac{b}{e^3}$$

$5 \times e^3$

5 × e ³	6	$\frac{c^3}{e+e+ee} + \frac{cce}{1+e+ee} = b$
6 × reduced	7	$b \times 1 + c + ee = cce \times 1 + e + ee + c^2$
	8	$ \begin{array}{r} be^6 + 3be^5 + 6be^4 + 7be^3 + 6be^2 \\ \quad \quad \quad - c^2e^2 - c^2e^2 \\ \quad \quad \quad + 3be + b \\ \quad \quad \quad - c^2e \quad \quad \quad \} = c^2 \end{array} $

PROB. XXIII.

There is a cask of rum, out of which was taken 42 gallons, and filled up with water, and the same repeated three times more. At last there was found by the proof, to remain 25.2935 gallons of rum in it. What was the content of the cask?

Let	1	$b = 42, c = 25.2935, a = \text{cask's content, then } a - b = \text{first remainder.}$
	2	<p>And since the quantity of liquor is as the space it possesses; therefore</p> $a : a - b \text{ (1 rem.)} :: a - b :$ $\frac{a - b^2}{a} \text{ (2 rem.)} :: \frac{a - b^2}{a} : \frac{a - b^3}{aa}$ $\text{(3 rem.)} :: \frac{a - b^3}{aa} : \frac{a - b^4}{a^3} \text{ (4 rem.)}$
per quest.	3	$\frac{a - b^4}{a^3} = c.$
3 ©	4	$a^4 - 4ba^3 + 6bba^2 - 4b^3a + b^4 = ca^3$
4 tr.	5	$a^4 - 4ba^3 + 6bba^2 - 4b^3a = -b^4.$
5 resol.	6	$a = 124.84 \text{ gallons.}$

PROB. XXIV.

Given	1	$x^3 + x^2y + y^2x + y^3 = dxy.$	
	2	$x^6 + x^4y^2 + y^4x^2 + y^6 = bbxxyy.$	
1,	3	$\frac{x^3 + y^3}{x^2 + y^2} \times \frac{x + y}{x + y} = dxy$	
2,	4	$\frac{x^4 + y^4}{x^2 + y^2} \times \frac{x^2 + y^2}{x^2 + y^2} = bbxxyy$	
Put	}	5	$xy = a$
		6	$x + y = e$
3, 5, 6	7	$\frac{ee - 2a}{e} \times e = da = \frac{x^3 + y^3}{x^2 + y^2} \times \frac{x + y}{x + y}$	
7 ÷ e	8	$ee - 2a = \frac{da}{e} = xx + yy$	
4, 8	9	$\frac{ddaa}{ee} - 2aa = x^4 + y^4$	
	10	$\frac{ddaa}{ee} - 2aa \times \frac{da}{e} = \frac{x^4 + y^4}{x^2 + y^2} \times$	
9 × $\frac{da}{e}$		$\frac{x^2 + y^2}{x^2 + y^2} = bbxxyy = bbaa.$	
10 ×	11	$\frac{dd - 2ee}{bbe^3} \times da = bbe^3$	
11 ÷	12	$a = \frac{d^2 - 2dee}{d^2 - 2dee}$	
8, 12	13	$e^3 - \frac{2bbe^4}{d^2 - 2dee} = \frac{db^2e^3}{d^2 - 2dee}$	
13 ×	14	$d^2e^3 - 2de^5 - 2bbe^4 = dbbe^3$	
14 ÷	15	$d^2 - 2dee - 2bbe = dbb$	
		$ee + \frac{bb}{d}e = \frac{dd - bb}{2},$ a quadratic	
		which gives e , and then a by	
		step 12, and x and y may be	
		found from step 5 and 6.	

PROB. XXV.

Given	}	1	$a^5e - bda^4 + 2ba^3ee + bbae^3 - 2dbba^2e$ $- b^3de^2 = d^5$
		2	$a^4ee - 2bda^3e + bbdda^2 + ba^2e^3 -$ $2dbbae^2 + bbd + b^3d^2e = b^4d^2.$
that is,	}	3	$\frac{aa+be}{ae-bd} \times \frac{ae-bd}{ae-bd} = d^5.$
1, 2		4	$\frac{ae-bd}{ae-bd} \times \frac{aa+be}{aa+be} = b^4d^2.$
Put	}	5	$aa+be = x$
		6	$ae-bd = y$
3, 5, 6		7	$x^2y = d^5$
4, 5, 6		8	$y^2x = b^4dd.$
7 × 8		9	$x^3y^3 = b^4d^8.$
9 lw 3		10	$xy = bdd \sqrt[3]{b^4dd}$
7 ÷ 10,		11	$x = \frac{d^5}{bdd \sqrt[3]{b^4dd}} = \frac{d^5}{bb} \sqrt[3]{bbd}$
8 ÷ 10		12	$y = \frac{b^4dd}{bdd \sqrt[3]{b^4dd}} = \frac{bb}{d} \sqrt[3]{bbd}.$
5 - be		13	$aa = x - be$
6, +		14	$a = \frac{y+bd}{e}$
13, 14		15	$aa = \frac{yy + 2bdy + bbdd}{ee} = x - be$
15 red.		16	$xec - be^3 = yy + 2bdy + bbdd, \text{ a cubic}$ equation which gives e , whence a is known by step 14.

PROB. XXVI.

To find four numbers x, y, z, v , having the product of every three given.

Suppose	}	1	$xyz = b$
		2	$yzv = c$
		3	$zvx = d$
		4	$vxy = f$

$1 \times 2 \times 3 \times 4$	5	$x^3 y^3 z^3 v^3 = bcd f$
5 lw 2	6	$xyzv = \sqrt[3]{bcd f}$
$6 \div 2$	7	$x = \frac{\sqrt[3]{bcd f}}{c}$
$6 \div 3$	8	$y = \frac{\sqrt[3]{bcd f}}{d}$
$\div 4$	9	$z = \frac{\sqrt[3]{bcd f}}{f}$
$6 \div 1$	10	$v = \frac{\sqrt[3]{bcd f}}{b}$

PROB. XXVII.

To find 3 numbers, x, y, z , having the product of each and the sum of the other two, given.

per qu. }	1	$x \times y + z = b$
	2	$y \times x + z = c$
	3	$z \times x + y = d$
$1 + 2 + 3$	4	$2xy + 2xz + 2yz = b + c + d$
$4 \div (2)$	5	$xy + xz + yz = \frac{b+c+d}{2} = s$, by subst.
$5 - 1$	6	$yz = s - b$
$5 - 2$	7	$xz = s - c$
$5 - 3$	8	$xy = s - d$
$6 \times 7 \times 8$	9	$xyyzxz = \overline{s-b} \times \overline{s-c} \times \overline{s-d}$
9 lw 2	10	$xyz = \sqrt{\overline{s-b} \times \overline{s-c} \times \overline{s-d}}$
$10 \div 6$	11	$x = \frac{\sqrt{\overline{s-c} \times \overline{s-d}}}{s-b}$
$10 \div 7$	12	$y = \frac{\sqrt{\overline{s-b} \times \overline{s-d}}}{s-c}$
$10 \div 8$	13	$z = \frac{\sqrt{\overline{s-b} \times \overline{s-c}}}{s-d}$

PROB. XXVIII.

To find any polygonal or figurate number.

A figurate or polygonal number is the sum of a series of numbers in arithmetical progression from 1. And these are so called, because they denote the number of points, which fill a regular polygon, placed at equal distances, on lines drawn parallel and equidistant, to the sides of the figure. The following table shews the arithmetic proportionals, and the polygonal numbers formed from them. The numbers of the arithmetical series shew what number of points are placed on the several parallel lines of the polygon; and the polygonal numbers, shew the whole number of the points contained in the figure. 21.

Rank.	Arithm. proportionals.	Polygonal numbers.	Names.
1	1, 1, 1, 1, 1, 1	1, 2, 3, 4, 5, 6	laterals.
2	1, 2, 3, 4, 5, 6	1, 3, 6, 10, 15, 21	triangul.
3	1, 3, 5, 7, 9, 11	1, 4, 9, 16, 25, 36	quadrang.
4	1, 4, 7, 10, 13, 16	1, 5, 12, 22, 35, 51	pentang.
5	1, 5, 9, 13, 17, 21	1, 6, 15, 28, 45, 66	hexang.
6	1, 6, 11, 16, 21, 26	1, 7, 18, 34, 55, 81	heptang.
7	1, 7, 13, 19, 25, 31	1, 8, 21, 40, 65, 96	octang.

Let

arith. prop Pr. 6.	1	$r =$ any rank, $x =$ polygonal number sought, $n =$ place of x ; then $r - 1 =$ common diff. of the arithm. series
ib. Pr. 7.	2	$1 + n - 1 \times r - 1 =$ the n^{th} term in the arithmetic progression.
1, 3,	3	$\frac{2 + n - 1 \times r - 1}{2} \times n = n^{\text{th}}$ term in the polygonal numbers.
	4	$x = \frac{2 + n - 1 \times r - 1}{2} \times n.$

S E C T. II.

Of Interest and Annuities.

P R O B. XXIX.

The principal, time, and rate of interest being given; to find the amount, or money due at the end of that time; at simple interest.

Let	1	$p =$ principal, $t =$ time, $r =$ rate of interest of 1 l. for a certain time, as a year, &c. $s =$ sum of all the arrears.
by proportion	}	2 $1 : r :: p : rp$, the interest of p for a year.
		3 $1 : rp :: t : prt$, the interest for the time t .
		4 $p + prt =$ whole arrear at the end of the time t .
1, 4	5	$p + prt = s$, the arrear sought.

Cor. 1. Hence $p = \frac{s}{rt+1}$, when s, r, t , are given.

Cor. 2. $t = \frac{s-p}{pr}$, when s, p, r , are given.

Cor. 3. $r = \frac{s-p}{pt}$, when s, p, t , are given.

P R O B.

PROB. XXX.

The annuity, time, and rate of interest being given; to find the arrear, at the end of that time, at simple interest.

Put	1	$a =$ annuity or yearly rent; $t =$ time of forbearance; $r =$ interest of 1 l. for a year, &c. $s =$ whole arrear.
by proportion	2	$0 =$ interest due at 1 year's end.
	3	$ra =$ interest at 2 year's end.
	4	$2ra =$ interest at 3 year's end.
	5	$3ra =$ interest for 4 years.
	6	$t-1 \cdot ra =$ interest for t years.
	7	$ta =$ rents due at the end of t years.
2,3,4,5,6,7	8	$0 + 1 + 2 + 3 \dots$ to $t-1 \times$ into $ra + ta = s.$
arith. prop. Prop. 7.	9	$0 + 1 + 2 + 3 \dots t-1 = \frac{t \times t - 1}{2}.$
8, 9,	10	$\frac{t \times t - 1}{2} ra + ta = s.$
10,	11	$\frac{t-1 \cdot r + 2}{2} ta = s.$

Cor. 1. $a = \frac{2s}{t-1 \cdot r + 2 \times t}$

Cor. 2. $t = \sqrt{\frac{2s}{ar} + \frac{2-r}{2r}} - \frac{2-r}{2r}$

Cor. 3. $r = \frac{2s - ta}{t-1 \times ta}$

PROB.

PROB. XXXI.

To find the present worth of an annuity, to continue a given time, at a given rate of simple interest.

Let	1	$p =$ present worth, $a =$ annuity, $t =$ time, $r =$ interest of 1 l.
Prob. 29.	2	$p + prt = s$
Prob. 30.	3	$\frac{t - 1.r + 2}{2} ta = s$
$2 = 3$	4	$p + prt = \frac{t - 1.r + 2}{2} ta$
$4 \div$	5	$p = \frac{t - 1.r + 2}{2rt + 2} ta = \frac{t - 1}{rt + 1} r + 1 ta.$

$$\text{Cor. 1. } a = \frac{rt + 1}{\frac{t - 1}{2} r + 1} \times \frac{p}{t}.$$

Cor. 2. $tt + \frac{2}{r} - \frac{2p}{a} - 1 \times t = \frac{2p}{ra}$, whence t may be found.

$$\text{Cor. 3. } r = \frac{2ta - 2p}{2p - t - 1.a \times t}$$

PROB. XXXII.

The principal, time, and rate of interest being given, to find the amount at the end of that time, at compound interest.

Let	1	$p =$ principal, $t =$ time, $r =$ interest of 1 l. $R = 1 + r$ the amount of 1 l. and its interest. $s =$ sum of money due at the end of that time.
-----	---	--

per quest. by proportion 1, 6	2	$1+r$ or $R =$ money due at 1 year's end.
	3	$1 : R :: R : RR =$ money due at 2 years end.
	4	$1 : R :: RR : R^3 =$ money due at 3 years end.
	5	$R^t =$ money due at t year's end.
	6	$1 : R^t :: p : pR^t =$ the amount of p for the time t .
	7	$pR^t = s$.

Cor. 1. $p = \frac{s}{R^t}$.

Cor. 2. $R^t = \frac{s}{p}$, or $t = \frac{\log:s - \log:p}{\log:R}$.

Cor. 3. $R = \sqrt[t]{\frac{s}{p}}$, or $\log:R = \frac{\log:s - \log:p}{t}$.

P R O B. XXXIII.

The annuity, time, and rate of interest, being given: to find the arrears due at the end of that time, at compound interest.

Let by Prob. 32.	1	$a =$ annuity, or yearly rent, $t =$ time of forbearance, $r =$ interest of 1 l. for a year, &c. $R = 1+r$, $s =$ sum of all the arrears.
	2	$a =$ money due at 1 year's end. $2a + ra = a + Ra =$ arrear at 2 years end.
	3	$a + aR + aRR =$ arrear in 3 years.
	4	$a + aR + aR^2 + aR^3 =$ arrear for 4 years.
	5	$a + aR + aR^2 + aR^3 \dots$ to
	6	$aR^{t-1} =$ the arrear for t years.

geom.

geom. propor. prop. 26	}	7		$1 + R + R^2 + R^3 \dots$ to R^{t-1}
				$= \frac{R \times R^{t-1} - 1}{R - 1} = \frac{R^t - 1}{r}$
6, 7		8		$\frac{R^t - 1}{r} a =$ money due at the end of t years.
1, 8,		9		$\frac{R^t - 1}{r} a = s.$

Cor. 1. $a = \frac{rs}{R^t - 1}$.

Cor. 2. $R^t = \frac{rs}{a} + 1$, or $t = \frac{\log \frac{rs}{a} + 1}{\log R}$.

Cor. 3. $\frac{s}{a} R - R^t = \frac{s-a}{a}$, whence R may be found, and then r .

P R O B. XXXIV.

To find the present worth of an annuity, to continue a given time, at a given rate of compound interest.

Let		1		$p =$ present worth, $a =$ the annuity, $t =$ the time, $r =$ interest of 1h. $R = 1 + r.$
Prob. 32.		2		$pR^t = s.$
Prob. 33.		3		$\frac{R^t - 1}{r} a = s.$
$2 = 3$		4		$pR^t = \frac{R^t - 1}{r} a$
$4 \div R^t$		5		$p = \frac{R^t - 1}{rR^t} a = \frac{1 - \frac{1}{R^t}}{r} a.$

Cor.

Cor. 1. $a = \frac{pr}{1 - \frac{1}{R^t}}$

Cor. 2. $R^t = \frac{a}{a - pr}$, or $t = \frac{\log:a - \log:a - pr}{\log:R}$

Cor. 3. $R^t + \frac{a}{p} R^t - R^{t+1} = \frac{a}{p}$, whence R and r will be found.

PROB. XXXV.

To find the value of an annuity to continue for ever, at a given rate of compound interest.

Let	1	$p =$ present worth, $a =$ annuity, $r =$ interest of 1 l. $R = 1 + r$.
Pr. 34. { step 5. {	2	$p = \frac{R^t - 1}{rR^t} a$.
Prob. 73. cor. 7.	3	but since t is infinite, R^t is infinitely greater than 1, whence $R^t - 1 = R^t$.
2, 3,	4	$p = \left(\frac{R^t}{rR} a = \right) \frac{a}{r}$.

Cor. 1. $a = pr$.

Cor. 2. $r = \frac{a}{p}$.

PROB. XXXVI.

At what rate of interest will 100 l. amount to 200 l. in $9\frac{1}{2}$ years, at compound interest.

Let	1	$r =$ rate of 1 l. $R = 1 + r$, $t = 9\frac{1}{2}$.
Prob. 32,	2	$100R^{39} = 200$ per quest.

A a

2 ÷	3	$R^{\frac{39}{4}} = 2$
3 \odot 4	4	$R^{39} = 16$
4 lw 39	5	$R = \sqrt[39]{16} = 1.0737$ by logarithms
5 - 1	6	$R - 1 = r = .0737$.
6 \times 100	7	7.37 = rate of interest per cent.

P R O B. XXXVII.

If a principal x be put out at compound interest, for x years, at x per cent. to find the time x , in which it will gain x .

Prob. 32.	1	$pR^t = s$.
per quest.	2	$p = x, r = \frac{x}{100}, R = 1 + \frac{x}{100}, t = x,$ $s = 2x.$
1, 2	3	$x \times 1 + \frac{x}{100}^x = 2x.$
3 ÷ x	4	$1 + \frac{x}{100}^x = 2.$
nature of logs.	5	$x \times 1 + \frac{x}{100} \times M = .3010300$
Prob. 84. cor. 1.	6	$x \times : \frac{x}{100} - \frac{xx}{20000} + \frac{x^3}{3000000}$ $\&c : = \frac{.30103}{M}$
6,	7	$\frac{xx}{100} - \frac{x^3}{20000} + \frac{x^4}{3000000} \&c.$ $= .693147$
by reverf.	8	$x = 8.49824$ years.

PROB. XXXVIII.

Given the rate per cent. for a year (5 l.), to find what the amount of any sum (100 l.), will be at the year's end, at compound interest; supposing it to arise from the principal and interest due every day, &c.

Let	{	1	$r =$ interest of 1 l. for a year.
		2	$n = 365$ the parts of a year.
		3	$\frac{r}{n} =$ interest for 1 day.
3,		4	$1 + \frac{r}{n} =$ money due at 1 day's end.
Prob. 32.		5	$1 + \frac{r}{n}^n =$ money due at the year's end.
by logs.		6	$n \times \log: 1 + \frac{r}{n} = \log: \text{amount}$
			for a year $= .0215694$.
6,		7	$1.0509 =$ amount for a year.
6×100		8	$105.09 =$ amount of 100 l.
or 5		9	$1 + \frac{r}{n}^n = 1 + r + \frac{n \cdot n - 1}{2nn} rr +$
			$\frac{n \cdot n - 1 \cdot n - 2}{2 \cdot 3n^3} r^3 \&c.$ the amount
		10	for a year.
			If the interest is supposed to gain interest every moment, by becoming part of the principal; then n is infinite, and
			$1 + \frac{r}{n}^n = 1 + r + \frac{r^2}{2} + \frac{r^3}{2 \cdot 3} + \frac{r^4}{2 \cdot 3 \cdot 4}$
			&c. the amount at the year's end. But this series is the number belonging to the hyperbolic logarithm r , whence

10,	11	The number belonging to the logarithm .43429448 $r =$ amount of 1 <i>l.</i> for a year $= 1.0513$; and for 100 $= 105.13$, to gain interest continually.
-----	----	--

Schol. If the interest for a day be required, so that it may amount to $1+r$ at the year's end, at compound interest; then the amount at 1 day's end, will be $\sqrt[n]{1+r}$; which is something less than $1 + \frac{r}{n}$.

P R O B. XXXIX.

A man puts out a sum of money at 6 per cent. to continue 40 years; and then both principal and interest is to sink. What is that per cent. to continue for ever?

The question amounts to this; if 100 *l.* be paid for an annuity of 6 *l.* a year for 40 years, what is that per cent?

Put	1	$a=6, p=100, t=40, r =$ rate of 1 <i>l.</i> $R=1+r.$
Prob. 34. cor. 2.	2	$t = \frac{\log:a - \log:a - pr}{\log:R}$
$2 \times$	3	$\text{Log: } R = \frac{\log:a - \log:a - pr}{t}$ $= \frac{1.6 - 1.0 - 100r}{40}$
Suppose	4	$R = 1.05$; then $r = .05$, and $L:R = .019454$; whence $R = 1.046$ which is too little.
Suppose	5	$R = 1.053$, then $r = .053$, and $L:R = .023324$, and $R = 1.055$, too big.

Then

Then by rule 5, Prob. xcii. B. I. you will find $R=1.052$, and the rate $=5.2$ per cent. which may be repeated for more exactness.

P R O B. XL.

If 200 l. be due 3 years hence; and 80 l. 5 years hence; in what time must both be paid together, at 5 per cent.

Let	1	$t =$ the time.
Prob. 32. cor. 1.	2	$\frac{200}{1.05^3} = 172.76$, the present worth of 200 l.
ib.	3	$\frac{80}{1.05^5} = 62.68$, the present worth of 80 l.
2+3	4	235.44, the whole present worth.
Prob. 32. cor. 1.	5	$t = \frac{\log:280 - \log:235.44}{\log:1.05} = 3.5527$ years.

P R O B. XLI.

What must I pay for an annuity of 70 l. to begin 6 years hence, and then to continue for 21 years, at 5 per cent?

Let	1	$a=70, t=21, R=1.05, x=6.$
Prob. 34.	2	$\frac{1 - \frac{1}{R^7}}{r} s =$ present worth of the annuity 7 years hence $=s.$
Prob. 32. cor. 1.	3	$\frac{s}{R^6} = \frac{1 - \frac{1}{R^7}}{rR^6} a =$ present worth of $s, 7$ years hence, $=669.704$ l. the present worth of the annuity in reversion.

S E C T. III.

Arithmetical and geometrical Progression.

P R O B. XLII.

A traveller sets out and goes 9 miles a day; 3 days after, another follows him, who travels the first day 4 miles, the second 5, the third 6, and so on. In what time will he overtake the first?

per quest.	1	$x =$ days the last travelled.
	2	$x - 1 + 4 =$ his last day's travel.
arith. progression.	3	$\frac{x - 1 + 4}{2} \times x =$ his whole journey.
per qu. {	4	$\frac{x + 3}{2} =$ days the first travelled.
	5	$x + 3 \times 9 =$ first man's journey.
$3 = 5$	6	$\frac{xx + 3x}{2} = 9x + 27$
6 reduced	7	$xx - 15x = 54.$
7 extr.	8	$x = 18.$

P R O B. XLIII.

There are three numbers in arithmetic progression, the square of the first together with the product of the other two is 16; and the square of the mean together with the product of the extremes is 17. What are the numbers?

Put	1	$a - e, a, a + e$ for the numbers,
		$b = 16, c = 17.$
per qu. {	2	$2aa - ae + ee = b$
	3	$2aa - ee = c$

Sect. III. ARITHMETICAL PROGRESSION. 359

2 + 3	4	$4aa - ae = b + c = s$ by subst.
4 tran.	5	$ae = 4aa - s$
3 tran.	6	$ee = 2aa - c$
$6 \times aa$	7	$aaee = 2a^4 - caa$
5 \odot 2	8	$aaee = 16a^4 - 8sa^2 + ss$
7 = 8	9	$16a^4 - 8sa^2 + ss = 2a^4 - caa$
9 reduc.	10	$14a^4 - 8s - c.aa + ss = 0$
10 ext.	11	$aa = 9, a = 3.$
$5 \div a$	12	$e = \frac{4aa - s}{a} = 1.$
1,	13	and the numbers are 2, 3, 4.

PROB. XLIV.

There are four numbers in arithmetical progression, whose common difference is 2, and product 3465.

Let	1	$2b = 2, \text{ or } b = 1, p = 3465; a - 3b,$ $a - b, a + b, a + 3b, \text{ the num-}$ bers sought.
per quest.	2	$aa - 9bb \times aa - bb = p$
2 \times	3	$a^4 - 10bba^2 + 9b^4 = p.$
3 extr.	4	$aa = 64, a = 8.$
1,	5	the numbers are 5, 7, 9, 11.

PROB. XLV.

To find five numbers in arithmetic progression, whose sum, and product are given.

Put	1	$a - 2e, a - e, a, a + e, a + 2e$ for the numbers, $b = \text{sum} = 25,$ $p = \text{product} = 2520.$
per qu.	2	$5a - 3e + 3e = 5a = b.$
	3	$a = \frac{b}{5} = m$ by subst.
	4	$a \times aa - 4ee \times aa - ee = p.$

3, 4,	5	$m \times \overline{mm} - 4ee \times \overline{mm} - ee = p.$
5 ×	6	$m \times m^2 - 5mme + 4e^2 = p$
6 ÷ m	7	$4e^2 - 5mme + m^2 = \frac{p}{m}.$
7 extr.	8	$ee = 1, e = 1.$
1	9	and the numbers are 3, 4, 5, 6, 7.

PROBLEM XLVI.

To find three numbers in geometrical progression, where the sum is 20, and the sum of their squares 140.

Let	1	x, y, z be the numbers, $b=20,$ $c=140.$
per qu.	2	$xz=yy$
	3	$x+y+z=b$
	4	$xx+yy+zz=c$
	5	$x+z=b-y$
3-y	5	$x+z=b-y$
5 ⊙ 2	6	$xx+2xz+zz=bb-2by+yy$
6, 2	7	$xx+zz+2yy=bb-2by+yy$
7-yy	8	$xx+zz+yy=bb-2by$
3, 8	9	$bb-2by=c.$
9 ÷	10	$y = \frac{bb-c}{2b} = 6\frac{1}{2}.$
2 × 4	11	$4xz=4yy$
6-11	12	$xx-2xz+zz=bb-2by-3yy$
12 lw 2	13	$x-z = \sqrt{bb-2by-3yy}$
5+13	14	$x = \frac{b-y + \sqrt{bb-2by-3yy}}{2} =$ $\frac{13\frac{1}{2} + \sqrt{13\frac{1}{2}}}{2}.$
5-13	15	$z = \frac{b-y - \sqrt{bb-2by-3yy}}{2} =$ $\frac{13\frac{1}{2} - \sqrt{13\frac{1}{2}}}{2}.$

P R O B. XLVII.

To find four numbers in geometrical progression, whose sum is 15, and the sum of their squares 85.

Let	1	v, x, y, z be the numbers, $b = 15$, $c = 85$.
per qu. }	2	$v + x + y + z = b$.
	3	$v^2 + x^2 + y^2 + z^2 = c$.
2 \odot 2	4	$\frac{v+x+v+y}{2} = \frac{x+y}{2} + \frac{v+z}{2} +$ $2 \times \frac{x+y}{2} \times \frac{v+z}{2} = \frac{xx+2xy+y^2}{2} +$ $\frac{yy+2vy+v^2}{2} + \frac{zz+2xz+z^2}{2} = \frac{bb}{2}$.
3, 4 by propor.	5	$c + 2xy + 2vz + 2 \times \frac{x+y}{2} \times \frac{v+z}{2} = \frac{bb}{2}$.
	6	$vz = xy$.
Put	7	$a = x+y, e = xy = vz$ by proportion.
2, 7	8	$y+z = b-a$.
5, 7, 8	9	$c + 4e + 2a \times \frac{b-a}{2} = \frac{bb}{2}$.
But	10	$v = \frac{xx}{y}, z = \frac{yy}{x}$, by the nature of proportion.
2, 10	11	$\frac{xx}{y} + x + y + \frac{yy}{x} = b$
7, 11	12	$a + \frac{x^2+y^2}{xy} = b$
7 \odot 3	13	$x+y = a^3 = x^3 + 3x^2y + 3xy^2 + y^3$
13 tran.	14	$x^3 + y^3 = a^3 - 3xy \times x + y$
14 \div , 7	15	$\frac{x^3+y^3}{xy} = \frac{a^3}{xy} - 3a = \frac{a^3}{e} - 3a$.
12, 15	16	$a + \frac{a^3}{e} - 3a = b$.
16 $\times e$	17	$a^3 - 2ae = be$
17 tran.	18	$be + 2ae = a^3$

18 ÷	19	$e = \frac{a^3}{b+2a}$
9, 19	20	$c + \frac{4a^3}{b+2a} + 2ab - 2aa = bb$
20 reduc.	21	$baa + ca = \frac{bb-c}{2} b$
21 extr.	22	$a = 6,$
19	23	$e = 8.$
7	24	$y = \frac{e}{x}, a = x + \frac{e}{x}$
24 ×	25	$ax = xx + e$
25 reduc.	26	$xx - ax + e = 0.$
26 extr.	27	$x = 2.$
10, 24	28	$y = 4, v = 1, z = 8.$

P R O B. XLVIII.

To find four numbers in geometrical progression, such that the difference of the means is 100, and the difference of the extremes 620.

Let	1	a, e, u, y be the numbers, $b = 100,$ $c = 620.$
per quest.	2	$y = a - c, u = e - b.$
But	3	$au = ce, ay = eu,$ by the nature of progression.
3, 2	4	$ae - ab = ce, aa - ac = ee - cb.$
4 ÷	5	$a = \frac{ce}{e-b}$
4, 5	6	$aa - ac = \frac{e^2}{e-b} - \frac{ce}{e-b} = ee - cb.$
6 ×	7	$\frac{e^2 - ce}{e-b} \times e - b = \frac{e^3}{e-b}$
7 tran.	8	$3b - c.ce + cb - 3bb.e = -b^3.$
8 red.	9	$ce - be = \frac{b^3}{c-3b}.$
9 extr.	10	$e = 125$
2, 5	11	$a = 625, y = 5, u = 25.$

P R O B.

P R O B. XLIX.

The sum of four quantities in geometrical progression being given, and the sum of the squares of the means, to find the quantities.

Let	1	a^3, a^2e, ae^2, e^3 be the quantities, $b = \text{sum of all, } c = \text{sum of the squares of the means.}$
	2	$a^3 + a^2e + ae^2 + e^3 = b.$
	3	$a^4e^2 + a^2e^4 = c.$
	4	$\frac{a^3 + a^2e + ae^2 + e^3}{a^4e^2 + a^2e^4} = \frac{a^2 + e^2}{a^2e + ae^2} \times \frac{a + e}{a^3c^3} = b.$
Put	5	$y = a^2e + ae^2,$
5 \odot 2	6	$yy = a^4e^2 + 2a^2e^3 + a^2e^4$
6,	7	$a^2e^3 = \frac{yy - c}{2}.$
3, 4, 5	8	$\frac{cy}{yy - c} = b$
8 red.	9	$b yy - 2cy = bc$
7,	10	$ae = \sqrt{\frac{yy - c}{2}} = d$
5,	11	$a + e = \frac{y}{ae} = \frac{y}{d}$
10 $\div a$	12	$e = \frac{d}{a}$
11, 12	13	$a + \frac{d}{a} = \frac{y}{d}.$
	14	$aa - \frac{ya}{d} + d = 0,$ whence $a, e,$ and all the numbers are known.

PROB. L.

There is given the sum of the squares of the extremes (b), of four quantities in geometrical progression; and the sum of the means (c); to find the quantities.

Let	1	a^3, a^2e, ae^2, e^3 be the quantities;
per qu. {	2	$a^6 + e^6 = b.$
	3	$a^4e^2 + a^2e^4 = c.$
2 + 3	4	$a^6 + a^4e^2 + a^2e^4 + e^6 = b + c = d.$
	5	$\frac{a^2 + e^2 \times a^2 + e^2}{2} = \frac{aa + ee}{2}$
5 \times a^4e^4	6	$\frac{a^6 + a^4e^2 \times a^2e^4 + e^6}{2} = \frac{a^2e^2 + a^2e^4}{2}$
Put	7	$y = a^6 + a^4ee$
4 - 7	8	$a^2e^4 + e^6 = d - y$
3, 6, 7, 8	9	$y \times d - y = cc,$
3, 7	10	$a^6 + 2a^4ee + aae^4 = y + c$
10 <i>lw</i> 2	11	$a^3 + aee = \sqrt{y + c} = p$
3, 4, 7,	12	$a^4ee + 2a^2e^4 + e^6 = c + d - y$
12 <i>lw</i> 2	13	$a^2e + e^3 = \sqrt{c + d - y} = q$
11 + 13	14	$a^3 + a^2e + ae^2 + e^3 = p + q.$ Whence the numbers will be found as in the last problem.

Or thus,

Let	1	$\frac{xx}{y}, x, y, \frac{yy}{x}$ be the quantities.
per qu. {	2	$\frac{x^4}{yy} + \frac{y^4}{xx} = b$
	3	$x + y = c,$
3 - x	4	$y = c - x$
2 \times $yyxx$	5	$x^6 + y^6 = bxxyy$
4, 5	6	$x^6 + c - x = bxx \times c - x^2$

6	7	$x^6 + c^6 - 6c^2x + 15c^4xx - 20c^3x^2 +$ $15c^2x^4 - 6cx^5 + x^6 = bccx^2 - 2bcx^3$ $+ bx^4$
reduced	8	$2x^6 - 6cx^5 + 15ccx^4 - 20c^3x^3$ $\quad \quad \quad -b \quad + 2bc$ $+ 15c^4xx - 6c^2x \quad + c^6 = 0.$ $\quad \quad \quad -bcc$

P R O B. LI.

Given the sum of the extremes (b) of five quantities in geometrical progression, and the sum of the three means (c), to find the quantities.

Let	1	$a^4, a^3e, a^2e^2, ae^3, e^4$ be the quantities.
per qu. {	2	$a^4 + e^4 = b.$
	3	$a^3e + a^2e^2 + ae^3 = c.$
	4	$a^2e^2 \times \frac{a^4 + 2a^2e^2 + e^4}{2} = a^2e^2 \times \frac{aa + ce}{2}$ $= a^2e + ae^2.$
Put	5	$y = aae.$
2, 4, 5	6	$y \times \frac{b + 2y}{2} = c - y^2$
6 reduced	7	$yy + by + 2cy = cc.$
2 + 5	8	$a^4 + 2aaee + e^4 = b + 2y$
8 Lw 2 {	9	$aa + ee = \sqrt{b + 2y}$
	10	$aa - ee = \sqrt{b - 2y}.$ Whence $a, e,$ and all the quantities are easily found.

P R O B. LII.

Of five quantities in geometrical progression, there is given the sum of the extrems (b), and the sum of the squares of the three means (c); to find the proportionals.

Let	1	$a^4, a^3e, a^2e^2, ae^3, e^4$, be the quantities.
per qu. {	2	$a^4 + e^4 = b$.
	3	$a^6e^2 + a^4e^4 + a^2e^6 = c$.
	4	$a^3e^2 \times a^4 + e^4 = a^6ee + a^4e^6$
Put	5	$y = aae$
2, 3, 4, 5	6	$y \times b = c - a^4e^4 = c - yy$.
6 + yy	7	$yy + by = c$.
2, 5	8	$a^4 + 2a^2ee + e^4 = b + 2y$
8 lw 2 {	9	$aa + ee = \sqrt{b + 2y}$.
	10	$aa - ee = \sqrt{b - 2y}$.
whence all the rest are found.		

P R O B. LIII.

There are four quantities in geometrical proportion discreet, whose sum is b , sum of the squares c , and sum of their cubes d ; to find the numbers.

Let	1	x, ex, y, ey , be the numbers.
per qu. {	2	$x + ex + y + ey = b$
	3	$x^2 + e^2x^2 + y^2 + e^2y^2 = c$
	4	$x^3 + e^2x^3 + y^3 + e^2y^3 = d$
	5	$x + y = v, x^2 + y^2 = z$.
Put {	6	$1 + e = s, 1 + ee = t, 1 + e^3 = u$
5 & 2 & c.	7	$xy = \frac{vv - z}{2}, x^3 + y^3 = \frac{vv - v^3}{2} v$
2, 5, 6,	8	$sv = b$

3, 5, 6

3, 5, 6

9

$tz=c$

4, 6, 7

10

$\frac{3z-vv}{2}vu=d.$

8, 9

11

$v=\frac{b}{s}, z=\frac{c}{t}$

10, 11

12

$\frac{3bcu}{2st} - \frac{b^3u}{2s^3} = d$

reduced

13

$3bcs^2u - b^3tu = 2s^3td.$

6, 13,

14

and restoring the values of s, t, u ;

$$\text{then } 3bc \times \frac{1+e^3}{1+e} \times \frac{1+e^2}{1+e} -$$

$$b^3 \times \frac{1+e^3}{1+e} \times \frac{1+ee}{1+ee} = 2d \times \frac{1+e}{1+e}$$

 $\times 1+ee$, a 5th power.And e being known all the rest are easily found.

S E C T. IV.

Unlimited Problems.

P R O B. LIV.

How many old guineas at 21 s. 6 d. and pistoles at 17 s. will pay 100 l.; and how many ways can it be done?

Let	1	$a = \text{guineas, } e = \text{pistoles; } 21\text{ s. } 6\text{ d.} = 43 \text{ fix-pences, } 17\text{ s.} = 34 \text{ fix-pences, and } 100\text{ l.} = 4000 \text{ fix-pences.}$
per quest.	2	$43a + 34e = 4000.$
2—	3	$34e = 4000 - 43a.$
3÷	4	$e = 116\frac{1}{4}.$
2÷	5	$a = \frac{4000 - 34e}{43} = wb.$
5 abridg.	6	$\frac{1 - 34e}{43} = wb.$
$\frac{43e}{43} + 6$	7	$\frac{9e + 1}{43} = wb.$
$7 \times (5)$	8	$\frac{45e + 5}{43} = wb.$
$8 \frac{43e}{43}$	9	$\frac{2e + 5}{43} = wb.$
$9 \times (4)$	10	$\frac{8e + 20}{43} = wb.$
$7 - 10$	11	$\frac{e - 19}{43} = wb. = p$
$11 \times (43)$	12	$e = 43p + 19 = 19, 62, 105, \text{ the pistoles.}$
5	13	$a = 78, 44, 10, \text{ the guineas; being three answers in whole numbers.}$

P R O B.

P R O B. LV.

What number is that, which being divided by 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, there will remain 1; but divided by 13, then 0 will remain.

	1	It is plain $5 \times 7 \times 8 \times 9 \times 11$, or 27720 is divisible by any of these numbers.
Put	2	$a =$ some whole number.
per quest.	3	$13e =$ the number sought.
	4	$27720a + 1 = 13e$
4 ÷	5	$e = \frac{27720a + 1}{13} = wb.$
5 abrid.	6	$\frac{4a + 1}{13} = wb.$
6 × (3)	7	$\frac{12a + 3}{13} = wb.$
(1) — 7	8	$\frac{a - 3}{13} = wb. = p.$
8 ×	9	$a = 13p + 3 = 3, 16, \&c.$
5,	10	$e = 6397.$
10 × (13)	11	$13e = 83161$ the number sought.

P R O B. LVI.

A man bought 20 birds for 20 pence; geese at 4 d. quails at half pennies, and larks at farthings. How many did he get of each?

Let	1	$a =$ geese, $e =$ quails, $y =$ larks.
per qu.	2	$a + e + y = 20.$
	3	$4a + e + y = 20.$
2 tr.	4	$y = 20 - a - e$
3, 4	5	$4a + e + 5 - \frac{a + e}{4} = 20$
5 red.	6	$15a + e = 60.$

B b

6,	7	$a = 4.$
2 tran.	8	$e = 20 - a - y$
3, 8	9	$4a + 10 + \frac{1}{2}y - \frac{a+y}{2} = 20$
9 red.	10	$7a - \frac{1}{2}y = 20.$
10	11	$a = 2.$
7, 11	12	$a = 3$
2, 12	13	$e + y = 17$
3, 12	14	$\frac{1}{2}e + \frac{1}{2}y = 8.$
14 \times (2)	15	$e + \frac{1}{2}y = 16$
13 - 15	16	$\frac{1}{2}y = 1, y = 2.$
13,	17	$e = 15.$

P R O B. LVII.

A, B, C, and their wives P, Q, R, went to the market to buy hogs. Each man and woman bought as many hogs as they gave shillings for each hog. A bought 23 hogs more than Q; and B bought 11 more than P. Also each man laid out 3 guineas more than his wife. Which 2 persons were man and wife.

Let	1	$x =$ hogs some man bought,
		$x - y =$ wife's hogs.
per qu.	}	2 the money for the man's $= xx,$
		3 $xx - 2xy + yy =$ wife's money.
4 tr.	4	$xx = xx - 2xy + yy + 63.$
	5	$2xy = 63 + yy$
5 \div	6	$x = \frac{63 + yy}{2y} = wb.$
6 $- y$	7	$x - y = \frac{63 - yy}{2y} = wb.$
But	8	In this case y must be an odd number, $= 1, 3, 5, 7, \&c.$ but it cannot be 5.

8,		9	If $y=1, x-y=31, x=32$
			$y=3, x-y=9, x=12,$
			$y=7, x-y=1, x=8$
per quest.		10	A has 32 hogs, and Q 9, Also B has 12, and P 1.
		11	Whence B and Q } C and P } are man and A and R } wife.

PROB. LVIII.

To find e, y in whole numbers, so that $yy-ee+22e=184.$

		1	To depress the equation, put $y=x+e.$
			2
2 ÷		3	$e = \frac{184-xx}{22+2x} = \frac{92-xx}{11+x} = wb.$ therefore x must be an even number. Therefore
2 ÷			4
4,		5	reduce the equation as low as it can, then $2e = -x + 11 +$ $\frac{63}{11+x} = wb.$
5,			6
Put		7	$11+x=p$
6, 7			8
7, 8		9	then $x = -10, -8, -4, -2, 10, 52$
8,			10
5, 9		11	$2e = 84, 40, 24, 20, 4, -40.$
11 ÷ (2)			12
1,		13	$y = 32, 12, 8, 8, 12, 32$ B b 2

And any pair of these will solve the problem, which are all the possible answers in whole numbers.

P R O B. LIX.

A vintner has wine at 24 d. 22 d. and 18 d. per gallon; of which he would mix 30 gallons, to be sold at 20 d. How much must be take of each?

Let	1	a, e, y be the quantities of each.
per qu. }	2	$a + e + y = 30.$
	3	$24a + 22e + 18y = 600.$
$2 \times (24)$	4	$24a + 24e + 24y = 720.$
$2 \times (22)$	5	$22a + 22e + 22y = 660$
$4 - 3$	6	$2e + 6y = 120.$
$6 \div (2)$	7	$e + 3y = 60. e = 60 - 3y.$
7,	8	$y \sqsupset 20.$
$5 - 3$	9	$-2a + 4y = 60. a = 2y - 30.$
$9 + 2a$	10	$4y = 60 + 2a.$
10,	11	$y \sqsubset 15.$
8, 10,	12	$y = 16, 17, 18, 19.$
9,	13	$a = 2, 4, 6, 8.$
7,	14	$e = 12, 9, 6, 3.$

P R O B. LX.

To find the value of e, y, u, x, z in whole numbers, in the two given equations following.

Given {	1	$e + y + u + x + z = 60.$	
	2	$3e + 4y + 5u + 7x + 9z = 440.$	
	$1 \times (4)$	3	$4e + 4y + 4u + 4x + 4z = 240$
	$1 \times (9)$	4	$9e + 9y + 9u + 9x + 9z = 540$
	$2 - 3$	5	$-e + u + 3x + 5z = 200. e \sqsubset 0.$
	$4 - 2$	6	$6e + 5y + 4u + 2x = 100. e \sqsupset 14\frac{2}{3}$
Suppose	7	$e = 10.$	
	$1 - e$	8	$y + u + x + z = 60 - e = 50$

$2 - 3e$

2—3 ^e	9	$4y+5u+7x+9z=440-3e=410.$
8 × (5)	10	$5y+5u+5x+5z=250.$
8 × (9)	11	$9y+9u+9x+9z=450.$
9—10	12	$-y+2x+4z=160. y \sqsubset 0.$
11—9	13	$5y+4u+2x=40. y \sqsubset 6\frac{1}{2}.$
Suppose	14	$y=4.$
8—y	15	$u+x+z=50-y=46.$
9—4y	16	$5u+7x+9z=410-4y=394.$
15 × (5)	17	$5u+5x+5z=230.$
15 × (7)	18	$7u+7x+7z=322$
16—17	19	$2x+4z=164. z \sqsupset 40\frac{1}{2}.$
16—18	20	$-2u+2z=72. z \sqsupset \text{or } \sqsubset 37.$
Suppose	21	$z=40.$
15—z	22	$u+x=46-z=6.$
16—9z	23	$5u+7x=394-9z=34$
22 × (5)	24	$5u+5x=30$
23—24	25	$2x=4. x=2$
22—x	26	$u=6-x=4.$

And one answer is got, viz. $e=10, y=4, u=4, x=2, z=40.$ for $10+4+4+2+40=60.$
and $3 \times 10 + 4 \times 4 + 5 \times 4 + 7 \times 2 + 9 \times 40 = 440.$

PROB. LXI.

To find a perfect number, or one which is equal to the sum of all its aliquot parts.

Suppose	1	$y^n \cdot x = \text{a perfect number.}$
	2	then $1+y+y^2 \dots \text{to } y^n, +x+$ $xy+xy^2 \dots \text{to } xy^{n-1} = \text{sum of}$ all the aliquot parts.
per quest.	3	$y^n \cdot x = 1+y \dots y^n + x \cdot 1+y \dots y^{n-1}.$

Cor. 3. Pr. 26. propor.	}	4	$1+y\dots y^n = \frac{y^{n+1}-1}{y-1}$
		5	$1+y\dots y^{n-1} = \frac{y^n-1}{y-1}$
3, 4, 5		6	$y^n x = \frac{y^{n+1}-1}{y-1} + \frac{y^n-1}{y-1} x$
6 \times tr.		7	$\frac{y^{n+1}-1}{y-1} \cdot y^n x - \frac{y^n-1}{y-1} x = y^{n+1}-1$
7 \div		8	$x = \frac{y^{n+1}-1}{y-1 \cdot y^n - y^n + 1}$
		9	But that x may be a whole number,
			$\frac{y-1 \cdot y^n - y^n + 1}{y-1 \cdot y^n} = 1$
9 tr.		10	$y-1 \cdot y^n = y^n$
10 \div		11	$y-1 = 1$
11 tr.		12	$y = 2$
1, 12		13	$2^n x =$ a perfect number.
8, 13		14	$2^n \times \frac{2^{n+1}-1}{2-1} =$ perfect number, where $2^{n+1}-1$ must be a prime, as appears by step 2.
		15	If n is an odd number greater than 1, then $2^{n+1}-1$ will be a com- posite number.

S E C T. V.

Rational Squares, Cubes, &c.

P R O B. LXII.

To find two square numbers, whose difference is given.

	1	Let xx and yy be the numbers, a = difference.
Put	2	$\frac{z+v}{2} = x, \frac{z-v}{2} = y$
2 \odot 2	3	$\frac{zz+2zv+vv}{4} = xx$
2 \odot 2	4	$\frac{zz-2zv+vv}{4} = yy$
3-4	5	$zv = xx - yy$
1, 5	6	$zv = a$
	7	Take v at pleasure, then $z = \frac{a}{v}$, whence x and y are known.
	8	If a is a whole number, and x and y are desired in whole numbers; take any two factors that produce a , so they be both even or both odd numbers, if possible. And therefore a must be either an odd number greater than 1, or a number divisible by 4, to have x and y in whole numbers.
If $a=27$.		Take $v=1, z=27$, or $v=3, z=9$
If $a=20$.		$v=2, z=10$.

PROB. LXIII.

To divide a given square into two other squares.

Let	1	$aa, ee =$ the squares required, $bb =$ the given square.
assume {	2	$a = sv$
	3	$e = rv - b$
	4	$aa = ssvv$
2 \odot 2	4	$aa = ssvv$
3 \odot 2	5	$ee = rrvv - 2rbv + bb$
4 + 5	6	$aa + ee = rr + ss.vv - 2rbv + bb$
per quest.	7	$rr + ss.vv - 2rbv + bb = bb$
7 tr.	8	$rr + ss.vv = 2rbv$
8 \div	9	$v = \frac{2rb}{rr + ss}$
2, 9	10	$a = \frac{2rbs}{rr + ss}$
3, 9	11	$e = \frac{rr - ss}{rr + ss} b$

PROB. LXIV.

To find a square number (aa), which multiplied by a given number (n), and a given square (bb) added to it; the sum may be a square.

Let	1	$naa + bb = yy.$
assume	2	$va \text{ } \odot \text{ } b = y$
2 \odot 2	3	$v^2aa - 2bva + bb = yy$
1 = 3	4	$v^2aa - 2bva + bb = naa + bb$
4 tr.	5	$vvaa - naa = 2bva$
5 $\div a$	6	$vv - n.a = 2bv$
	7	$a = \frac{2bv}{vv - n}$, where v may be taken at pleasure.

PROB.

P R O B. LXV.

To find two square numbers, (aa , ee), that their product added to a given number (d), may be a square.

Let	1	$aaee + d = yy$
assume	2	$ae - v = y$
2 \odot 2	3	$aaee - 2aev + vv = yy$
1 = 3	4	$aaee + d = aaee - 2aev + vv$
4 tr.	5	$2aev = vv - d$
	6	$e = \frac{vv - d}{2av}$, where a and v may be taken at pleasure.

P R O B. LXVI.

To find three such numbers x , y , z ; so that $yy = xz$; and $x + y$, and $z + y$, may be two squares.

Assume	1	$x + y = aa$
	2	$z + y = ee$
1 \rightarrow y	3	$x = aa - y$
2 \rightarrow y	4	$z = ee - y$
3 \times 4	5	$xz = aa - y \times ee - y = yy$ per quest.
5 \times	6	$yy = a^2e^2 - aay - eey + yy$.
6 tr.	7	$a^2e^2 = aa + ee.y$
	8	$y = \frac{aaee}{aa + ee}$, where a , e , may be taken at pleasure.
3, 8	9	$x = aa - \frac{aaee}{aa + ee} = \frac{a^4}{aa + ee}$
4, 8	10	$z = ee - \frac{aaee}{aa + ee} = \frac{e^4}{aa + ee}$

P R O B.

P R O B. LXVII.

To find a number, from which two given numbers
(a, b) being severally subtracted; the remainders
shall be two squares.

Let	1	$x =$ the number sought.
per qu.	}	2 $x - a = yy$
		3 $x - b = zz$
		4 $x = a + yy$
3, 4	5	$a + yy - b = zz$
assume	6	$v - y = z$
6 \odot 2	7	$vv - 2vy + yy = zz$
5 = 7	8	$a - b + yy = vv - 2vy + yy$
8 tr.	9	$2vy = vv + b - a$
9 $\div 2v$	10	$y = \frac{vv + b - a}{2v}$
4, 10	11	$x = a + \frac{vv + b - a}{4vv}$

P R O B. LXVIII.

To find three numbers (x, y, z), whose sum shall be a
square, and also the sum of any two to be a square.

Suppose	}	1	$x + y = rr$
		2	$x + z = ss$
		3	$y + z = tt$
		4	$x + y + z = vv$
4 \times (2)	5	$2vv = 2x + 2y + 2z$	
1 + 2 + 3	6	$2x + 2y + 2z = rr + ss + tt$	
5 = 6	7	$2vv = rr + ss + tt$	
Put	8	$s = p - v, t = q - v$	
7, 8	9	$2vv = rr + pp - 2pv + vv$ $+ qq - 2qv + vv$	
9 tr.	10	$2pv + 2qv = rr + pp + qq$	

$$11 \quad v = \frac{rr + pp + qq}{2p + 2q}, \text{ where } r, p, q \text{ are taken at pleasure; whence } s \text{ and } t \text{ are known (step 8).}$$

$$4-3 \quad 12 \quad x = vv - tt$$

$$4-2 \quad 13 \quad y = vv - ss$$

$$4-1 \quad 14 \quad z = vv - rr$$

PROB. LXIX.

To find three squares in arithmetic proportion.

Suppose	}	1	$xx - y = vv$
		2	xx
		3	$xx + y = zz$
1+3		4	$2xx = vv + zz$
assume	}	5	$v = s - x$
		6	$z = t - x$
4, 5, 6		7	$2xx = ss - 2sx + xx$ $+ tt - 2tx + xx$
7 tr.		8	$2s + 2t \cdot x = ss + tt$
8 ÷		9	$x = \frac{ss + tt}{2s + 2t}$
5, 9		10	$v = s - x = \frac{ss + 2st - tt}{2s + 2t}$
6, 9		11	$z = t - x = \frac{2st + tt - ss}{2s + 2t}$

PROB. LXX.

To find two numbers (x, y) so that $xy + x$, and $xy + y$, may be squares.

per qu.	}	1	$xy + x = vv$
		2	$xy + y = \text{a square.}$
1 ÷ x		3	$y + 1 = \frac{vv}{x}$

3—1	4	$y = \frac{vv-x}{x}$
2, 4	5	$\frac{vv-x}{x} \times \overline{x+1} = \square$
5,	6	to effect this, let $x+1$ be the side, then $\frac{vv-x}{x} = x+1$
6 red.	7	$vv = xx + 2x = \text{square.}$
Let	8	$r-x = \text{side,}$
7, 8	9	$rr - 2rx + xx = xx + 2x$
9 red.	10	$x = \frac{rr}{2r+2}$
4, 6, 10	11	$y = 1 + \frac{rr}{2r+2}$, where r may be taken at pleasure.

PROB. LXXI.

To find two numbers, whose sum and difference, shall be two squares.

Let	1	a, e , be the numbers.
per qu. {	2	$a+e = yy$
	3	$a-e = \text{a square.}$
	4	$a = yy - e$
2—e	5	$yy - 2e = \text{a square.}$
Put	6	$r-y = \text{root of it.}$
5, 6	7	$rr - 2ry + yy = yy - 2e$
7 tr.	8	$2ry = rr + 2e$
8 ÷ 2r	9	$y = \frac{rr+2e}{2r}$, where r, e may be taken at pleasure; then
4,	10	$a = yy - e.$

PROB.

P R O B. LXXII.

To find three numbers (a, e, y), that the sum of their squares may be a square.

per quest.	1	$aa + ee + yy = vv$
assume	2	$v = d + y$
2 \odot 2	3	$v^2 = dd + 2dy + yy$
1 = 3	4	$aa + ee + yy = dd + 2dy + yy$
4 \div	5	$y = \frac{aa + ee - dd}{2d}$, where a, e , and d are taken at pleasure.

P R O B. LXXIII.

To divide a number into two parts, so that the sum of the squares may be a square.

Let	1	$s =$ the number ; a, e , the parts.
per qu. {	2	$s = a + e$
	3	$aa + ee = vv.$
2 \odot 2	4	$aa + 2ae + ee = ss.$
4 —	5	$aa + ee = ss - 2ae$
2 — a	6	$e = s - a$
3, 6, 5	7	$aa + ee = ss - 2as + 2aa = vv$
assume	8	$ra - s = v.$
7, 8	9	$ss - 2as + 2aa = rraa - 2rsa + ss$
9 reduc.	10	$a = \frac{r-1}{rr-2} \times 2s$
6, 10	11	$e = \frac{r-2}{rr-2} \times rs.$

PROB. LXXIV.

To find two numbers in the ratio of b to c , so that either of them added to the square of the other, shall make two squares.

Let	1	$ba, ca,$ be the numbers.
per qu. {	2	$bbaa + ca =$ a square.
	3	$ccaa + ba =$ a square.
Put	4	$bbaa + ca = \overline{ba - v}^2 = bbaa - 2bv + vv$
4 tr.	5	$2bva + ca = vv.$
5 ÷	6	$a = \frac{vv}{2bv + c}$
3, 6	7	$ccaa + ba = \frac{ccvv}{2bv + c} + b \times \frac{vv}{2bv + c}$
7,	8	$ccaa + ba = \frac{ccvv + 2bbv + bc \times \left(\frac{vv}{2bv + c}\right)^2}{2bv + c} =$ a square.
8 ÷ □	9	$ccvv + 2bbv + bc =$ a square $= zz.$
Put	10	$z = cv - r.$
9, 10	11	$ccvv + 2bbv + bc = ccv - 2cvr + rr$
11 reduc.	12	$v = \frac{rr - bc}{2bb + 2cr}.$

PROB. LXXV.

To find a number, to which adding a given cube number, the sum shall be a cube; and subtracting another cube number, the remainder shall be a cube.

Let	1	x be the number; b^3, c^3 the two cubes.
per qu. {	2	$x + b^3 =$ a cube
	3	$x - c^3 =$ a cube

assume

assume	4	$x+b^3 = \overline{b + \frac{cc}{bb}a}^3 = b^3 + 3c^2a + \frac{3c^4}{b^3}aa + \frac{c^6}{b^6}a^3.$
4,	5	$x = 3c^2a + \frac{3c^4}{b^3}aa + \frac{c^6}{b^6}a^3.$
assume	6	$x-c^3 = \overline{a-c}^3 = a^3 - 3a^2c + 3acc - c^3$
6,	7	$x = a^3 - 3a^2c + 3acc$
5=7	8	$a^3 - 3a^2c + 3acc = 3acc + \frac{3c^4}{b^3}aa + \frac{c^6}{b^6}a^3.$
8 ÷	9	$a - 3c = \frac{2c^4}{b^3} + \frac{c^6}{b^6}a.$
9 reduced	10	$a = \frac{b^3+c^3}{b^6-c^6} \times 3cb^3 = \frac{3cb^3}{b^3-c^3}$

P R O B. LXXVI.

To divide the sum of two given cubes into two other cubes.

Let	1	x^3, y^3 be the cubes sought; b^3, c^3 the given cubes.
assume	2	$x = b + v$
	3	$y = c - \frac{bb}{cc}v$
	2 ⊙ 3	4
3 ⊙ 3	5	$y^3 = c^3 - 3bbv + \frac{3b^4}{c^3}v^2 - \frac{b^6}{c^6}v^3$
4+5	6	$x^3 + y^3 = b^3 + c^3 + 3bv^2 + \frac{3b^4}{c^3}v^2 + v^3 - \frac{b^6}{c^6}v^3 = b^3 + c^3, \text{ per quest.}$

6 tr.	7	$3bv^2 + \frac{3b^4}{c^3}v^2 = \frac{b^6}{c^6}v^3 - v^3.$
$7 \times \frac{c^6}{vv}$	8	$3bc^6 + 3b^4c^3 = b^6v - c^6v.$
8 ÷	9	$v = \frac{b^3 + c^3}{b^6 - c^6} \times 3bc^3 = \frac{3bc^3}{b^3 + c^3}.$

PROB. LXXVII.

To find three such cube numbers, whose sum may be both a square and a cube number.

Let	1	a, e, y be their roots; x^6 the sum of their cubes.
per quest.	2	$a^3 + e^3 + y^3 = x^6$
$2 - y^3$	3	$e^3 + y^3 = x^6 - a^3$
assume	4	$\frac{x^4}{aa}v - a = e$
	5	$xx - v = y$
	6	$a^3 = \text{a cube}$
4 \odot 3	7	$+\frac{x^{12}}{a^6}v^3 - \frac{3x^3}{a^3}v^2 + 3x^4v - a^3$
5 \odot 3	8	$+x^6 - 3x^4v + 3x^2v^2 - v^3$
6, 7, 8,	9	$\frac{x^{12}}{a^6}v^3 - \frac{3x^3}{a^3}vv + 3x^2v^2 - v^3 = 0.$
$9 \times \frac{a^6}{vv}$	10	$x^{12}v - a^6v = x^8 - x^2a^3 \times 3a^3.$
Suppose	11	$v = \frac{x^6 - a^3}{x^{12} - a^6} \times 3a^3x^2 = \frac{3a^3xx}{x^6 + a^3},$ where
or suppose	x and a may be taken at pleasure; then v being known, e and y are known by step 4, and 5.	
	$x = 1, a = 1,$ then $v = \frac{1}{3},$ whence	
	$e = \frac{5}{8}, y = \frac{2}{3},$ and the numbers are	
	$\frac{1}{8} + \frac{125}{512} + \frac{8}{27} = 1,$	
	$x = 2, a = 1,$ then $v = \frac{13}{63},$ and $e = \frac{137}{273},$	
	$y = \frac{248}{273};$ and the numbers	
	$1 + \frac{2048383}{274625} + \frac{15252992}{274625} = 64.$	

S E C T. VI.

Geometrical Problems.

HAVING hitherto, in all the foregoing sections, kept an account of the whole process by registering the several steps at length in the margin; so that the reader may see at once how each step is derived from the rest; and by this means become acquainted with the manner of proceeding, in any operation. It may be presumed, that by this time, he will be able to see the connection of the several parts of the process in any solution, without such a formal explanation. Therefore, for brevity's sake, in what follows, I shall not tie myself to this method, but generally write down the process after a shorter way, without notifying all these particulars; and content myself with mentioning only such deductions as are less obvious.

P R O B. LXXVIII.

In the triangle CAD; there are given AC, AD; and the lines CE, DB, drawn to the given points E, B; to find the point of intersection F. 23.

Put $AB=r$, $CB=m$, $AE=p$, $AD=d$, $CE=f$, $DB=g$; and the line sought $CF=a$. Draw EI \parallel to DB .

By the similar triangles CBF, CIE (Geom. II.

12,) $a : f - a :: m : \frac{mf - ma}{a} = BI$, and $d : p :: r :$

$\frac{pr}{d} = AI$. Then $AI + BI = AB$, that is, $\frac{pr}{d} +$

Fig.
23.

+ $\frac{mf - ma}{a} = r$, and multiplying by da , $pra + dmf - dma = dar$, and $dra + dma - pra = dmf$, whence

$$a = \frac{dmf}{dr + dm - pr}.$$

P R O B. LXXIX.

24. To divide a triangle ABC in a given ratio, by a line drawn through a given point P.

Through P draw ED parallel to BA, and put $AB = b$, $AC = d$, $BC = f$, $BE = g$, $EP = p$, $BF = x$, and the ratio as m to n , and $m + n = s$.

By similar triangles, $g + x : p :: x : \frac{px}{g + x} = BI$; then (Geom. II. 19.) $BI \times BF : BA \times BC :: m : m + n$, that is,

$\frac{p x x}{g + x} : b f :: m : s$; then $b m f = \frac{s p x x}{g + x}$, and $s p x x = b m f x + b m f g$: by which equation x is found.

P R O B. LXXX.

25. To divide a triangle into two equal parts, by a line of a given length.

Let BD be perpendicular to AC, KH the given line, and HL parallel to BD. Put $AC = a$, $BC = b$, $HK = c$, $CD = d$, $CK = x$; then (Geom. II. 19.) $AC \times BC = 2KC \times HC$, or $ab = 2x \times CH$, and

$CH = \frac{ab}{2x}$, and by similar triangles (Geom. II. 13)

$b : d :: \frac{ab}{2x} : \frac{abd}{2bx} = CL$; and $KL = KC - CL = x$

$\frac{abd}{2bx} = \frac{2bx - abd}{2bx}$. But (Geom. II. 21. cor. 5.)

HK =

Sect. VI. PROBLEMS.

387

$$HK^2 - KL^2 = CH^2 - CL^2, \text{ or } \frac{4bbccxx - 4bbx^2 - a^2b^2d^2 + 4ab^2dx^2}{4bbxx} = \frac{a^2b^2 - a^2b^2d^2}{4bbx}$$

Fig. 25.

or $4bbx^2 - 4bbccx^2 - 4adbb = -a^2b^2$, whence x will be found.

PROB. LXXXI.

To find the inaccessible distance AB, by help of the triangle ACD; CAB being one right line. 26.

Through B draw BEF, and draw EG parallel to CD. Put $AC = a$, $AD = b$, $CD = c$, $AE = d$, $CF = f$, and $AB = x$. Then, by similar triangles, $AD (b) :$

$$CD (c) :: AE (d) : EG = \frac{cd}{b}; \text{ and } AD (b) : CA (a)$$

$$:: AE (d) : AG = \frac{ad}{b}.$$

Then $GB = \frac{ad+bx}{b}$. And by the similar triangles BGE, BCF; $CF (f) : CB (a+x) :: EG (\frac{cd}{b}) : GB (\frac{ad+bx}{b})$. Therefore $\frac{adf+bfx}{b} = \frac{cda+cdx}{b}$, $bfx - cdx = cda - fda$, and $x = \frac{c-f}{bf-cd} da$.

PROB. LXXXII.

If the line EFB be drawn from the angle E, perpendicular to the diagonal AD of a right-angled parallelogram, and BF, FD are given. To find the sides of the parallelogram. 27.

Let $AF = x$, $EF = y$, $BF = b$, $DF = c$. The triangles AFB, AFE, and DFE are similar. Therefore

Fig. 27. fore $b : x :: x : \frac{xx}{b} = FE = y$, and $b : x :: y$ or $\frac{xx}{b} : c$.

Whence $\frac{x^3}{b} = bc$, and $x^3 = bbc$, and $x = \sqrt[3]{bbc}$.

Then $AE = \sqrt{xx + yy}$, and $ED = \sqrt{cc + yy}$.

PROB. LXXXIII.

28. To describe a square in the given triangle ATE.

Draw TC perpendicular to AE, and let BFGD be the square. Put $AE = b$, $AC = c$, $CE = d$, $TC = p$, BF or $BD = x$, $AB = y$. Then

The triangles ABF, ACT are similar, and $y : x :: c : p$, whence $cx = py$. Also the triangles EDG, ECT are similar, and $ED = b - x - y$, whence $b - x - y : x :: d : p$, and $dx = pb - px - py = pb - px - cx$. Whence $x = \frac{pb}{d + c + p} = \frac{pb}{b + p}$.

PROB. LXXXIV.

29. Six equal circles of 2 inches diameter are inscribed in an equilateral triangle, touching one another and the sides of the triangle. To find the side of the triangle.

Draw AF perpendicular to BC, and from the centers O, S, draw OD, SE perpendicular to AB, and let $DO = r$, $AB = x$.

The triangles ABF, ADO, ESB are similar, and (Geom. II. 39. cor.) $AF = AB \sqrt{\frac{3}{4}}$. Then BF

$(\frac{1}{2}x) : AF (x\sqrt{\frac{3}{4}}) :: DO (r) : AD = \frac{rx\sqrt{\frac{3}{4}}}{\frac{1}{2}x} = 2r\sqrt{\frac{3}{4}}$

$= EB$, and $DE = 4r$, whence AB or $x = 4r + 4r$

$\sqrt{\frac{3}{4}} = \frac{2 + \sqrt{3}}{1} \times 2r = 4 + 2\sqrt{3}$.

PROB.

PROB. LXXXV.

There are two circles BDA and BFC touching in B, and if DE be perpendicular to BA at the center E; then there is given AC and DF; to find the diameters. 30.

Let radius $BE = a$, $DF = b$, $CA = d$; then $FE = a - b$, $EC = a - d$, then $FE^2 = BE \times EC$ (Geom. IV. 17), that is, $aa - 2ab + bb = aa - ad$, and $2ba - da = bb$, and $a = \frac{bb}{2b - d}$, whence $BC = 2a - d$.

PROB. LXXXVI.

In the triangle ABC, there are given the three perpendiculars, from the angles upon the opposite sides; to find the sides. 31.

Let $AO = a$, $CP = b$, $BR = c$, and $AB = y$.

Then twice the area $= by = AC \times c = CB \times a$, whence $AC = \frac{by}{c}$, and $CB = \frac{by}{a}$. And (Geom.

II. 21.) $\frac{bby}{cc} - bb = AP^2$; and (Geom. II. 23. cor.)

$$AP = \frac{AC^2 + AB^2 - CB^2}{2AB} = \frac{\frac{bby}{cc} + yy - \frac{bby}{aa}}{2y} =$$

$$\frac{bby}{2cc} + \frac{1}{2}y - \frac{bby}{2aa}; \text{ whence } \frac{bby}{2cc} + \frac{1}{2}y - \frac{bby}{2aa} =$$

$$\sqrt{\frac{bby}{cc} - bb}. \text{ That is, } aabby + aaccy - bbccy =$$

$$2aac \sqrt{bby - bbcc}; \text{ put } aabb + aacc - bbcc = d, \text{ then}$$

$$dy = 2aac \sqrt{bby - bbcc}; \text{ and by reduction,}$$

$$y = \frac{aabc}{\sqrt{a^2bbcc - dd}}$$

P R O B. LXXXVII.

32. In the triangle ABC, there is given the rectangle of the sides; the rectangle of the segments of the base, made by a perpendicular; and the area: to find the rest.

Let the area $=b$, $AD \times DC = c$, $AB \times BC = d$, and $BD = z$, $2y =$ difference of the segments AD, DC .

Then $\frac{2b}{z} = AC$, and $\frac{b}{z} + y = DC$, $\frac{b}{z} - y =$

DA. Whence $\frac{bb}{zz} - yy = c$, and $\sqrt{z^2 + \frac{b}{z} + y} \times$

$\sqrt{z^2 + \frac{b}{z} - y} = d$; and squaring all the quantities,

and putting $\frac{bb}{zz} - c$ for yy , and v for $zz + \frac{2bb}{zz} - c$,

and then $v + \frac{2by}{z} \times v - \frac{2by}{z} = dd = vv -$

$\frac{4bby}{zz} =$ (restoring the values of v and y) $z^4 + \frac{4b^4}{z^4}$

$+ cc + 4bb - 2czz - \frac{4cbb}{zz} - \frac{4b^4}{z^4} + \frac{4bbc}{zz}$, or $z^4 -$


$2czz + 4bb + cc = dd$. Whence z is known, and

$y = \sqrt{\frac{bb}{zz} - c}$, and then AD, DC , and AB, BC will be found.

P R O B. LXXXVIII.

33. In the right-angled triangle ABD, there is given the perpendicular, on the hypotenuse; and the radius of the inscribed circle: to find the sides.

Put the perpendicular $BQ = p$, radius $CR = r$, $AD = a$, $AB = e$, $BD = y$. Then (Geom. II. 21.)



[Faint, illegible text, likely bleed-through from the reverse side of the page.]

$aa = ee + yy$. And (II. 20. cor. 2.) $pa = ey$. But Fig. AD or $AF + FD = AR + DI$, and $AD + 2CR = 33$.

$AB + BD$, that is, $a + 2r = e + y$; whence $aa + 2pa = ee + 2ey + yy = e + y = a + 2r = aa + 4ra + 4rr$,

therefore $2pa - 4ra = 4rr$, and $a = \frac{2rr}{p - 2r}$. Also

$aa - 2pa = ee - 2ey + yy = e - y$, and $\frac{e - y}{2} =$

$\frac{\sqrt{aa - 2pa}}{2}$. Whence $e = \frac{a + 2r + \sqrt{aa - 2pa}}{2}$,

and $y = \frac{a + 2r - \sqrt{aa - 2pa}}{2}$.

P R O B. LXXXIX.

There is an isosceles triangle, in which two circles are inscribed, touching one another and the sides of the triangle; their diameters are 8 and 12: to find the sides of the triangle. 34.

From the centers D, F, draw DG, FH \perp to BC, and FO \parallel to CB \dots draw CFDA.

Put $DG = r$, $FH = s$, $DO = r - s = c$, $FD = r + s = b$. Then $FO = \sqrt{bb - cc} = d$, and $CB = a$.

The triangles DFE and BCA are similar, whence $b : d :: a : \frac{da}{b} = AC$, and $c : b :: r : \frac{rb}{c} = CD$,

then $r + \frac{rb}{c} = AC = \frac{da}{b}$, whence $a = \frac{bbr}{cd} +$

$\frac{br}{d} = \frac{360}{\sqrt{96}}$.

P R O B. XC.

35. *There is given AD, and CD the radius of the semi-circle CEG, to find the radius of a circle inscribed between AC, the tangent AE, and the circle CE.*

Draw from the center O, the line OI perpendicular to AC; through O, draw AOF bisecting the angle DAE, and put radius $DE=r$, $AD=d$, $OI=a$, then $AE=\sqrt{dd-rr}=b$.

Then (Geom. II. 25.) $AD:AE::DF:EF$, and $AD+AE:AE::DE:EF$; that is, $d+b:$

$b::r:\frac{br}{d+b}=EF$, $DO=a+r$, and $DI=$

$\sqrt{DO^2-OI^2}=\sqrt{rr+2ra}$, and $AI=d-\sqrt{rr+2ra}$, and by the similar triangles AEF and AOI,

$d-\sqrt{rr+2ra}:a::b:\frac{br}{b+d}$. Then $ba=\frac{dbr}{b+d}$

$-\frac{br}{b+d}\sqrt{rr+2ra}$. Put $b+d=c$, and reducing,

$$\begin{array}{r} cca - 2r^2a = r^4 \\ -2drc \qquad - drr. \end{array}$$

P R O B. XCI.

36. *Through a given point B, to draw the right line BDC, so that the part DC comprehended between the two lines AC, AH, equidistant from B, may be of a given length.*

Produce CA to E, and compleat the rhombus EABH; make the angle $CDF=CAF$, and let $CD=a$, AE or AH= b , BA= d , AC= x , AF= y .

The triangles CAD, CEB are similar, therefore

$$CA(x):CD(a)::AE(b):DB=\frac{ab}{x}.$$

Since $\angle FDC = FAC$, therefore their supplements $FDB=$

FDB=CAB, and so the triangles BAC and BDF Fig. are similar, whence BA (d) : AC (x) :: DB 36.

$$\left(\frac{ab}{x}\right) : DF = \frac{ab}{d}.$$

But the triangles FAD and FDB are similar; for $\angle BDF = \angle FAD$, (for $\angle BAD = \angle BAE = \angle FAC$, add $\angle DAC$, then $\angle CAB = \angle FAD$, that is, $\angle FDB = \angle FAD$;) and $\angle F$ is common; therefore AF (y) : DF ($\frac{ab}{d}$) :: DF : $\frac{aabb}{ddy} = FB = d + y$, which reduced is $ddy + dy = aabb$; whence y will be had.

Again, the triangles DAF and BAC are also similar, and $CB = a + \frac{ab}{x}$, then DF ($\frac{ab}{d}$) : AF (y) :: CB : CA (x); whence $\frac{abx}{d} = \frac{aby}{x} + ay$, reduced $abxx - adyx = abdy$, whence x is had. Then

$$CE (b+x) : EB (b) :: AC (x) : AD = \frac{bx}{b+x}.$$

PROB. XCII.

Through a given point B, to draw the right line BDC, 36_a so that the part DC, included between the lines AC, AH, may be given.

Through B, draw BH, BE parallel to EAC and AH, and put $CD = a$, $AE = b$, $AH = c$, $AB = d$, $PH = f$, BP being perpendicular to AH, and $AC = x$.

The triangles CAD and CEB are similar, and $CE (x+b) : EB (c) :: CA (x) : AD = \frac{cx}{b+x}$;

and $DH = c - \frac{cx}{b+x} = \frac{bc}{b+x}$. And (Geom. II.

$$22.) BD = \sqrt{bb + \frac{ccbb}{bb + 2bx + xx} + \frac{2bcf}{b+x}}, \text{ and } b :$$

Fig.

36.

$$b : \sqrt{bb + \frac{ccbb}{bb + 2bx + xx} + \frac{2bcf}{b+x}} :: x : a;$$

$$\text{whence } bbaa = bbxx + \frac{ccbbxx}{bb + 2bx + xx} + \frac{2bcfxx}{b+x}.$$

$$\text{which reduced is } x^4 + 2bx^3 + bbx^2 - 2baax - bbaa = 0.$$

$$+ \frac{2cf}{b} - aa$$

P R O B. XCIII.

37. *The difference of the height of two hills being given, and their distance; to find their heights.*

Let BA, ED be the hills, put radius $r = CR = 698000$, $DE - BA = b = 119$, $AB = a$, $BE = c = 6$. Then $CB = r + a$, $CE = r + b + a$. Then (Geom.

II. 21.) $BR = \sqrt{2ra + aa}$, $RE =$

$\sqrt{bb + 2br + 2ra + 2ba + aa}$, whence $BE =$

$\sqrt{2ra + aa} + \sqrt{bb + 2br + 2ra + 2ba + aa} = c$, and

$\sqrt{bb + 2br + 2ra + 2ba + aa} = c - \sqrt{2ra + aa}$, and by

squaring, $bb + 2br + 2ra + 2ba + aa = cc + 2ra + aa -$

$2c\sqrt{2ra + aa}$, and $2c\sqrt{2ra + aa} = cc - bb - 2rb -$

$2ba = dd - 2ba$ (by substitution); and when squared

$8ccra + 4ccaa = d^2 - 4ddba + 4bbaa$, and when re-

duced, $aa + 2ra = \frac{cc - bb - 4rb}{4} + \frac{rrbb}{cc - bb}$ and

$a = 164,69$.

P R O B. XCIV.

38. *Three lines drawn from the three angles of a triangle to the middle of the opposite sides, being given; to find the sides.*

Put $AD = b = 18$, $E = c = 24$, $BF = d = 30$, $CB = x$, $AB = y$, $AC = z$.

Then (Geom. II. 28.) $yy + zz = 2bb + \frac{1}{2}xx$, $yy +$
 $xx = 2dd + \frac{1}{2}zz$, $zz + xx = 2cc + \frac{1}{2}yy$; and adding these

these three equations, $2xx + 2yy + 2zz = 2bb + 2cc$ Fig. 38.

$$+ 2dd + \frac{1}{2}xx + \frac{1}{2}yy + \frac{1}{2}zz, \text{ and } xx + yy + zz = \frac{4}{3}bb$$

$$+ \frac{4}{3}cc + \frac{4}{3}dd, \text{ from this subtract the first equa-}$$

$$\text{tions, then } xx = \frac{4}{3}dd - \frac{2}{3}bb + \frac{4}{3}cc - \frac{1}{2}xx, \text{ or } 9xx$$

$$= 8cc + 8dd - 4bb, 9yy = 8bb + 8dd - 4cc, 9zz = 8bb + 8cc - 4dd, \text{ whence } x = 34,176; y = 28,844; z = 20.$$

PROB. XCV.

ABC is an equilateral triangle, O a point in it equidistant from A, B, C. If the sides, and the line BO be all produced till they cut the line PD in D, E, R, P; then there is given DE, ER, RP; to find the side of the triangle ABC, and the area. 39.

Draw EF, EG parallel to BP, BR; and put $DE = a = 304$; $ER = b = 121.6$; $RP = c = 159.6$; and $DR = d$, $DP = s$, CL or $AL = x$, CG or $FG = y$. Then (Geo. II. 39. cor.) $BL = x\sqrt{3}$, $EG = y\sqrt{3}$. The triangles DEF and DPA are similar, whence

$$a : 2y :: s : \frac{2sy}{a} = AP, \text{ and } PB = 2x + \frac{2sy}{a}.$$

Since $\angle PBR = \angle EBR$, therefore (Geo. II. 25) $2x + \frac{2sy}{a} : 2x + 2y :: c : b$, and $2bx + \frac{2bsy}{a} = 2cx +$

$$2cy, \text{ whence } cx - bx = \frac{bsy}{a} - cy, \text{ and } y = \frac{ca - ba}{bs - ac} x$$

$\frac{fx}{g}$, by substitution.

Again $DE (a) : EG (y\sqrt{3}) :: DR (d) : \frac{dy}{a}\sqrt{3}$

$$= RL = \frac{dfx}{ag}\sqrt{3}, \text{ and } RB = x\sqrt{3} + \frac{dfx}{ag}\sqrt{3}, \text{ and}$$

$$PB = 2x + \frac{2sfx}{ag}, \text{ and } BE = 2x + 2y = 2x + \frac{2fx}{g}.$$

But

Fig. But (Geom. II. 26.) $BR^2 + PR \times RE = PB \times EB$,

39. that is, $3xx \times 1 + \frac{df}{ag} + bc = 4xx \times 1 + \frac{f}{g} \times 1 + \frac{sf}{ag}$

And by reduction, $1 + \frac{4f}{g} + \frac{4sf - 6df}{ag} + \frac{4saff - 3ddf}{aagg} \times$ into $xx = bc$. Whence $x = 78.4$,

$y = 40$, and the area $ABC = 10646.16$.

P R O B. XCVI.

40. In the triangle ABC, there is given the base, and difference of the sides and the area: to find the triangle.

Let the area $= f = 796$; difference of the sides CA, $CB = b = 10$; base $AB = d = 50$; perpendicular $CD = \frac{2f}{d} = p = 31.84$, and $AD = a$. Then

$AC = \sqrt{aa + pp}$, and $CB = \sqrt{d - a + pp}$; therefore by the question $\sqrt{aa + pp} + b =$

$\sqrt{dd - 2da + aa + pp}$, which squared is $aa + bb + pp + 2b\sqrt{aa + pp} = dd - 2da + aa + pp$, and $2b\sqrt{aa + pp} = dd - bb - 2da$, and squaring both sides $4bbpp = d^2 + b^2 - 2ddb - 4d^2a + 4bbda + 4ddaa$. Which

reduced is $aa - da = \frac{bbpp}{dd - bb} - \frac{dd - bb}{4}$, whence

$a = 16.739$, $AC = 36$, $BC = 46$, $BD = 33.261$.

P R O B. XCVII.

41. There is given the side of a rhombus, and the side of its inscribed square; to find the area.

Let $AB = BD = d = 4\frac{1}{2}$, $CO = CE = s = 3$, $BC = x$. Then $DC = d - x$, and $AC = d + x$.

The

The triangles ACE and CDO are similar, and Fig. 41.

$d+x : s :: d-x : \frac{d-x}{d+x} s = DO$. And (Geom. II. 41.

$$21.) \quad ss + ss \times \frac{d-x}{d+x} = \overline{d-x}^2; \text{ that is } 2ss \times \overline{d+x}$$

$$= \overline{dd-xx}^2; \text{ reduced, } \begin{array}{r} x^4 - ddx + d^4 \\ - 2ss \quad - 2ssd \end{array} = 0:$$

Whence $x = \sqrt{ss+dd} \pm \sqrt{ss+4dd} = \frac{5}{8}$, and

AC=5, AE=4, DO=2 $\frac{1}{2}$, DQ=5 $\frac{1}{2}$, area = 73 $\frac{1}{2}$,
QA=7.

PROB. XCVIII.

Given the four sides of a trapezium inscribed in a circle; to find the diagonals, and diameter of the circle. 42.

Let AB=a, BC=b, CD=c, AD=d. BE=x, the triangles ABE, and CED are similar; for $\angle ABE = \angle ECD$ (Geom. IV. 12. cor. 2.); and the angles at E are vertical; therefore AB (a) : BE (x) :: DC (c) : CE = $\frac{cx}{a}$; also the triangles AED

and BEC are similar, and BC (b) : CE ($\frac{cx}{a}$) ::

AD (d) : DE = $\frac{dcx}{ab}$. And BC (b) : BE (x) ::

AD (d) : AE = $\frac{dx}{b}$. Then BD = $x + \frac{dcx}{ab}$,

and AC = $\frac{dx}{b} + \frac{cx}{a}$. Then (Geom. IV. 32.) AC

\times BD = AB \times CD + AD \times BC, or $\frac{dxx}{b} + \frac{ddcx}{abb} + \frac{cxx}{a}$

+ $\frac{dcxx}{baa} = ac + bd$, whence x is had, and then AC

and BD are known.

Then

Fig. Then suppose a perpendicular from A upon BD, 42. then (Geom. II. cor. 23.) the distance of the per-

pendicular from D is $= \frac{AD^2 + DB^2 - AB^2}{2BD} = f.$

And $\sqrt{AD^2 - ff} =$ the perpendicular $= p$; and (Geom. IV. 28.) $p : AD :: AB : \text{diameter of the}$
circumscribing circle $= \frac{AD \times AB}{p}.$

P R O B. XCIX.

43. *The three semicircles HFG, HEJ, and GOJ touch one another in H, G, and I; to draw a fourth circle FOE to touch all the rest.*

From the centers A, B, C draw the lines ADE, BD, and CD; and DP perpendicular to AC, and let $AG = a$, BE or BI $= b$, CG $= c$, and $DE = x$. Then $AD = a + x$, $BD = b - x$, $CD = c + x$, $AC = a + c$, $BC = b - c$.

In the triangle ADC (Geom. II. 22. cor.)
 $PC = \frac{c+a + c+x - a+x}{2c+2a}$, and in the triangle
BDC, $PC = \frac{b-c + c+x - b-x}{2b-2c}.$

Whence

$$\begin{array}{r} cc + 2ca + aa \\ + cc + 2cx + xx \\ - aa - 2ax - xx \\ \hline c + a \end{array} = \begin{array}{r} bb - 2bc + cc \\ + cc + 2cx + xx \\ - bb + 2bx - xx \\ \hline b - c \end{array}$$

That is,

$$\frac{2cc + 2ca - 2cx - 2ax}{c + a} = \frac{2cc - 2bc + 2cx + 2bx}{b - c};$$

And

And multiplying alternately,

$$\begin{array}{r} 2bcc + 2bca + 2bcx - 2abx = 2c^3 - 2bcc + 2ccx + 2bcx \\ - 2c^3 - 2acc - 2ccx + 2cax \qquad + 2cca - 2bca + 2cax \\ \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad + 2bax \end{array}$$

And

$$4abx + 4ccx = 4bcc + 4bca - 4c^3 - 4cca,$$

Whence $x = \frac{a+c \times b-c}{ab+cc} c.$

PROB. C.

In the triangle ACB, there is given the sides AC, CB; and the length and breadth of the inscribed rectangular parallelogram DEHF; to find the rest. 44.

Draw CP perpendicular to AB, and let CA = b, CB = c, DE or GP = p, DF = a, CP = z, AB = y; and let p = b + c, q = b - c.

The triangles CDF and CAB are similar, and z : y :: z - p : a; whence za = zy - py, and zy - za = py; therefore z = $\frac{py}{y-a}$.

Again, (Geom. II. 24.) y : p :: q : $\frac{pq}{y}$ = diff. segments of the base. Therefore AP = $\frac{1}{2} y + \frac{pq}{2g}$.

But (Geom. II. 21.) bb = $zx + \frac{1}{2} y + \frac{pq}{2y}$ = $\frac{ppy}{y-a} + \frac{1}{4} yy + \frac{1}{2} pg + \frac{ppqq}{4yy}$. Which equation reduced is

$$\begin{array}{r} y^6 - 2ay^5 + aay^4 + 8abby^3 + ppqqy^2 - 2ap^2q^2y \\ + 4pp - 2apq \quad + 2aapq \\ + 2pq \qquad \qquad - 4aabb \\ - 4bb \\ + aappqq = 0. \end{array}$$

PROB.

Fig.

P R O B. CI.

45. *In the right-angled triangle AVP, KD is drawn parallel to the base, and there is given the base AP, and the segments VD, AK; to find the rest.*

Let $AK=b=200$, $AP=c=400$, $VD=d=260$, and $DP=a$; then (Geom. II. 12.) $a : b :: d :$
 $\frac{bd}{a} = VK$, and $VA = b + \frac{bd}{a} = \frac{b}{a} \times \overline{a+d}$. But

$$AV^2 - VP^2 = AP^2, \text{ or } \frac{bb}{aa} \times \overline{a+d}^2 - \overline{a+d}^2 = cc,$$

or $\overline{bb-aa} \times \overline{a+d}^2 = aacc$, that is,

$$\begin{aligned} a^4 + 2da^3 + ccaa - 2bbda &= bdd. \\ + dd \\ - bb \end{aligned}$$

And $a=141,727$, and $a+d$ or $VP=401,727$.

P R O B. CII.

46. *In the figure CAFD; CA, BP, DF are perpendicular to AF; and the sides of the triangle being produced, there is given HA, AC, CB, and BD, DF, FT; to find the sides of the triangle HBT.*

Let $HA=n$, $AC=p$, $CB=f$, $BD=d$, $DF=c$, $FT=b$, and $TH=a$, $TP=z$, $PH=v$, $BP=x$, $BT=y$, $e=HB$.

The triangles TBP and TGA are similar, and $y : x :: f+y : p$, and $py = fx + yx$, or $py - xy = fx$, whence $y = \frac{fx}{p-x}$. The triangles HBP and HDF are similar, and $e : x :: e+d : c$, or $ce = ex + xd$, and $ce - xc = dx$, whence $e = \frac{dx}{c-x}$. Likewise $z : x ::$
 $a+n$

$a+n : p$, and $pz = \overline{a+n} \times x$, and $z = \frac{a+n}{p} x$.

Fig.
46.

Likewise $v : x :: a+b : c$, and $cv = \overline{a+b} \times x$, and
 $v = \frac{a+b}{c} x$.

But $v+z=a = \frac{a+n}{p} x + \frac{a+b}{c} x$, whence $pca =$
 $cax + cnx + pax + pbx$, and $pca - cxa - pxa = cnx +$
 pbx ; therefore $a = \frac{cnx + pbx}{pc - cx - px}$.

Again, $TC = f + \frac{fx}{p-x} = \frac{pf}{p-x}$, and (Geom. II.

$$21.) \sqrt{\frac{ppff}{p-x}} - pp = n+a = n + \frac{cnx + pbx}{pc - cx - px} =$$

$$\frac{pcn - ncx - npx + ncx + pbx}{pc - cx - px} = \frac{pcn + rpx}{pc - sx}, \text{ (putting}$$

$r = b - n$, and $s = c + p$); and by squaring,

$$\frac{ppff - p^2 + 2p^2x - pp^2x}{pp - 2px + xx} = \frac{ppccnn + 2ppcnrx + rrpp^2x}{ppcc - 2pcsx + ssxx},$$

which reduced is

$$\begin{aligned} pprrx^2 + 2ppcnrx^2 + ppccnrx^2 - 2p^2cennx + p^2ccnn &= 0 \\ + ppss - 2p^2rr &- 4p^2cnr + 2p^2cnr + p^2cc \\ - 2p^2ss &+ p^2rr - 2cp^2s - ccp^2ff \\ - 2p^2cs &+ p^2ss + 2cp^2ffs \\ &- ppffss - 2ccp^2s \\ &+ 4p^2cs \\ &+ p^2cc \end{aligned}$$

PROB. CIII.

Given the sides and area of a trapezium; to find the diagonal. 47.

Draw the perpendiculars BE, DF upon the diagonal AC; and put $AB = a = 4$, $BC = b = 6$, $CD = c = 7$, $DA = d = 5$, and $f =$ the area.

D d

Then

Fig.
47.Then (Geom. II. 23. cor.) $CE = \frac{yy+bb-aa}{2y}$,

$$\text{and } BE = \sqrt{bb - \frac{yy+bb-aa}{2y}} =$$

$$\sqrt{\frac{4bbyy - y^4 - 2yy \times bb - aa - bb - aa}{4yy}} =$$

$$\sqrt{\frac{-y^4 + yy \times 2bb + 2aa - bb - aa}{4yy}}. \text{ In like man-}$$

$$\text{ner } DF = \sqrt{\frac{-y^4 + yy \times 2cc + 2dd - cc - dd}{4yy}}.$$

$$\text{Put } \frac{2aa+2bb}{2} \times yy - \frac{bb-aa}{2} = pyy - qq = v.$$

$$\text{and } \frac{2cc+2dd}{2} \times yy - \frac{cc-dd}{2} = ryy - ss = z.$$

$$\text{Then } \frac{1}{2}y \times \overline{BE + DF} = f, \text{ and } 2y \times \overline{BE + DF}$$

$$= 4f, \text{ that is, } \sqrt{v-y^4} + \sqrt{z-y^4} = 4f, \text{ and by squa-}$$

$$\text{ring } v+z-2y^4 + 2\sqrt{vz} - \frac{v}{z}y^4 + y^8 = 16ff, \text{ and}$$

$$\sqrt{vz} - \frac{v}{z}y^4 + y^8 = 8ff - \frac{v+z}{2} + y^4, \text{ and by squar-}$$

$$\text{ing, } vz - \frac{v}{z}y^4 + y^8 = 64f^4 - 8ff \times \frac{v+z}{2} + \frac{v+z}{4}$$

$$+ 16ff - \frac{v-z}{2} \times y^4 + y^8. \text{ And } vz = 64f^4 - 8ff \times$$

$$\frac{v+z}{2} + \frac{v+z}{4} + 16ffy^4 = 64f^4 - 8ff \times \frac{v+z}{2} +$$

$$\frac{vv}{4} + \frac{2vz}{4} + \frac{zz}{4} + 16ffy^4, \text{ or } 16ffy^4 - 8ff \times \frac{v+z}{2}$$

$$+ \frac{vv-2vz+zz}{4} + 64f^4 = 0; \text{ that is, } 64ffy^4 -$$

$$32ff \times \frac{v+z}{2} + \frac{v-z}{2} + 256f^4 = 0, \text{ and restor-}$$

ing the values of v and z , we shall have

+ 64f

P R O B. C V.

49. *There are given the three sides of the triangle ABC, and the angles A, and B, are bisected by the lines AD, BE; to find the length of one as AD, and also the distance AF to the point of intersection F.*

Put $AB=a$, $BC=b$, $AC=c$, and $AD=x$, $AF=y$.
Then (Geom. II. 25.) $AB:AC::BD:DC$, and
 $AB+AC:AB::BD+DC:BD$; that is, $a+c:$

$a::b:\frac{ab}{a+c}=BD$; likewise $a+c:c::b:\frac{bc}{a+c}=CD$.
But (Geom. II. 26.) $AD^2+BDC=BAC$
that is, $xx+\frac{abbc}{a+c}=ac$, and $xx=ac-\frac{bb}{a+c}$ ×

$$ac=ac \times \frac{a+c-bb}{a+c} = ac \times \frac{a+c+b \times a+c-b}{a+c}$$


whence $x = \frac{\sqrt{ac \times a+c+b \times a+c-b}}{a+c} = AD$.


Again, $AB+BD:AD::AB:AF$, that is,
 $a+\frac{ab}{a+c}:\frac{\sqrt{ac \times a+c+b \times a+c-b}}{a+c}::a:$

$$y = \frac{a\sqrt{ac \times a+c+b \times a+c-b}}{a+c \times a + \frac{ab}{a+c}} =$$

$\frac{\sqrt{ac \times a+c+b \times a+c-b}}{a+c+b}$, that is,

$$y = \sqrt{\frac{ac \times a+c-b}{a+c+b}} = AF.$$





117-107

PROB. CVI.

The diameters of three circles being given, which are 50.
described from the angular points of a triangle, as
centers, whose three sides are given; to find the ra-
dius of a fourth circle to touch all the three.

Let ABC be the given triangle, D the center of
the circle required; on AB let fall the perpendi-
culars DE, CK, and draw DF perpendicular to
AC. And put $AB=b$, $AC=c$, $CB=d$, and
 $AO=r$, $BR=s$, $CT=t$; and $AK=g$, $KC=b$;
and $AE=x$, $AF=y$, $OD=a$. In the triangle
ADB, (Geom. II. 23.) $aa+2as+ss=aa+2ar+rr$
 $+bb-2bx$. Whence $2bx=rr+bb-ss-2as+2ar$,
and $x=\frac{rr+bb-ss-2as+2ar}{2b}$. And in the tri-
angle ADC, $aa+2at+tt=aa+2ar+rr+cc-2cy$,
and $2cy=rr+cc-tt-2ta+2ra$, and
 $y=\frac{rr+cc-tt-2ta+2ra}{2c}$.

The triangles ACK, AFG are similar, and
 $g:c::y:\frac{cy}{g}=AG$, then $x-\frac{cy}{g}=GE$. Also the
triangles DGE (AGF,) and ACK are similar;
whence $b:g::x-\frac{cy}{g}:\sqrt{aa+2ar+rr-xx}=DE$.
Whence $b\sqrt{aa+2ar+rr-xx}=gx-cy$.

Put $l=rr+bb-ss$, $f=2s-2r$, $m=rr+cc-tt$,
 $n=2t-2r$, $p=lg-bm$, $q=bn-fg$. Then $x=\frac{l-fa}{2b}$, and $b\sqrt{aa+2ar+rr-\frac{l-fa}{2b}}=\frac{lg-fga}{2b}$
 $-\frac{m-na=p}{2} \frac{qa}{2b}$. Which squared is $bbaa+2rbba$

Then to expunge y (by Prob. liv. rule 2) we have Fig. 52.

$$A = -2cf, B = -2cfxx - 4bcx + 2cbb.$$

$$D = mfx + 2afx - pf \\ - mfx - 2mbx - mbb.$$

Whence $AB + DD = 0$, that is,

$$\left. \begin{array}{l} 4ccfxx + 8bcfx - 4bbccf \\ + 4aaff - 4apff + pff \\ - 4ambf - 4ambb + 2mbbpf \\ + 4mmbb + 4mbpf + mmb^2 \\ + 4mmb^2 \end{array} \right\} = 0.$$

PROB. CVIII.

In a triangle, there is given a perpendicular, the difference of the sides, and the difference of the segments of the base; to find the sides. 53.

Let the perpendicular $CD = a$, $CB - CA = c$, and $BD - DA = b$, $DA = x$. Then $CA = \sqrt{aa + xx}$, and $CB = \sqrt{aa + xx} + c$, and $AB = 2x + b$. Then (Geom. II. 24.), $b + 2x : 2\sqrt{aa + xx} + c :: c : b$; whence $bb + 2bx = 2c\sqrt{aa + xx} + cc$, and $2bx + bb - cc = 2\sqrt{aa + xx}$, and squared is $4bbxx + 4bx \times bb - cc + bb - cc = 4aa + 4xx \times cc$;

reduced

$$\begin{array}{r} 4bbxx + 4b^2x = 4aacc. \\ - 4cc \quad - 4bcc \quad - b^2 \\ \quad \quad \quad + 2bbcc \\ \quad \quad \quad - c^2 \end{array}$$

Fig.

P R O B. CIX.

54. *There is given the perpendicular in a triangle, and the two differences between the least side, and the other two; to find the sides.*

Let the perpendicular $AD = a$, $BC - BA = b$, $AC - AB = c$, $AB = x$; then $BC = b + x$, $AC = c + x$, and $BD = \sqrt{xx - aa}$, and $DC = b + x - \sqrt{xx - aa}$; and (Geom. II. 24.) $BC (b + x) : AC + AB (c + 2x) :: AC - AB (c) : DC - DB (b + x - 2\sqrt{xx - aa})$; whence $\frac{b + x}{2b + 2x} \times \sqrt{xx - aa} = \frac{cc + 2cx}{bb + 2bx + xx - cc - 2cx}$; and $\frac{b + x}{2b + 2x} \times \sqrt{xx - aa} = \frac{bb + 2bx + xx - cc - 2cx}{bb + 2bx + xx - cc - 2cx}$

Put $bb - cc = d$, $2b - 2c = f$; then

$\frac{4bb + 8bx + 4xx \times xx - aa}{4bb + 8bx + 4xx \times xx - aa} = \frac{xx + fx + d^2}{xx + fx + d^2}$, which multiplied and reduced is

$$\begin{aligned} 3x^4 + 8bx^3 + 4bbx^2 - 8baax - 4bbaa &= 0. \\ - 2f &- 4aa &- 2df &- dd \\ &- ff \\ &- 2d \end{aligned}$$

P R O B. CX.

55. *Having all the sides of a right-angled triangle ACB; to find either segment of the base AD, the perpendicular CD, the area, and the radius of the inscribed circle, &c.*

1. Let $AC = a$, $CB = c$, $AB = b$, $z = \frac{a + b + c}{2}$; then (Geom. II. 20. cor. 1.) $aa = b \times AD$, and $AD = \frac{aa}{b}$. But $aa = bb - cc = \overline{b + c} \times \overline{b - c}$; therefore $AD = \frac{\overline{b + c} \times \overline{b - c}}{b}$, that is,

the

the segment $AD = \frac{aa}{b} = \frac{\overline{b+c} \times \overline{b-c}}{b}$.

2. For the perpendicular CD .

$$CD^2 = AC^2 - AD^2 = aa - \frac{a^4}{bb} = \frac{aa \times \overline{bb-aa}}{bb}, \text{ and}$$

$$CD = \frac{a}{b} \sqrt{\overline{bb-aa}} = \frac{a}{b} \sqrt{\overline{b+a} \times \overline{b-a}}; \text{ or}$$

$$CD = \frac{a}{b} \sqrt{cc} = \frac{ac}{b}. \text{ Therefore the perpendicular}$$

$$\text{lar } CD = \frac{ac}{b} = \frac{a}{b} \sqrt{\overline{bb-aa}} = \frac{a}{b} \sqrt{\overline{b+a} \times \overline{b-a}}.$$

3. For the area.

$$\frac{AB \times CD}{2} = \text{area}; \text{ that is, the area} = \frac{ac}{2} =$$

$$\frac{a \sqrt{\overline{b+a} \times \overline{b-a}}}{2}. \text{ And since } aa + cc = bb, \text{ add } 2ac,$$

then $aa + 2ac + cc$ or $\overline{a+c}^2 = \overline{bb} + 2ac$, and $2ac =$

$$\overline{a+c}^2 - \overline{bb}, \text{ and } \frac{ac}{2} \text{ or the area} = \frac{\overline{a+c}^2 - \overline{bb}}{4} =$$

$$\frac{\overline{a+c+b} \times \overline{a+c-b}}{4} = z \times z - b. \text{ And since } \overline{a+c}^2$$

$$+ \overline{c}^2 = 2aa + 2cc = 2bb, \text{ therefore } \overline{a+c}^2 - \overline{bb} = \overline{bb}$$

$$- \overline{a-c}^2. \text{ Therefore the area} = \frac{\overline{bb} - \overline{a-c}^2}{4} =$$

$$\frac{\overline{b+a-c} \times \overline{b-a+c}}{4} = \overline{z-a} \times \overline{z-c}. \text{ Hence the}$$

$$\text{area} = \frac{ac}{2} = \frac{a \sqrt{\overline{b+a} \times \overline{b-a}}}{2} = \frac{\overline{a+c+b} \times \overline{a+c-b}}{4}$$

$$= \frac{\overline{b+a-s} \times \overline{b-a+c}}{4} = z \times \overline{z-u} = \overline{z-a} \times \overline{z-c}.$$

4. For

4. For the radius of the inscribed circle.

The area $= \frac{a+b+c}{2} =$ radius of the inscribed circle (Geom. IV. 30. cor.) $= \frac{a+b+c}{2} \times r$, putting r for the radius; then $r = \frac{2 \text{ area}}{a+b+c} = \frac{a+c-b}{2} = z-b$; or the radius $= \frac{ac}{a+b+c}$, that is, the radius of the inscribed circle is $= \frac{ac}{2z} = \frac{a+c-b}{2} = z-b$.

5. For the circumscribing circle.

The radius of the circumscribing circle $= \frac{1}{2} b$ (Geom. IV. 14.)

6. For the tangents.

The tangent, or the distance from A to the point of contact of the inscribed circle $= \frac{a+b-c}{2} = z-c$, (Geom. IV. 30).

And the distance from the right angle C is $= r$ the radius.

7. For the distance of the center.

The distance from A $= \sqrt{rr + \frac{a+b-c}{2}} = \sqrt{\frac{a+c-b}{2} + \frac{a+b-c}{2}}$ (by Art. 4. and 6) $= \sqrt{\frac{a+b-c}{4} + \frac{a-b-c}{4}} = \sqrt{\frac{a+d}{4} + \frac{a-d}{4}}$ (putting

$$\begin{aligned} \text{(putting } d=b-c) &= \sqrt{\frac{2aa+2dd}{4}} = \sqrt{\frac{aa+b-c^2}{2}} \quad \text{Fig. 55.} \\ &= \sqrt{\frac{aa+bb-2bc+cc}{2}} = \sqrt{\frac{2bb-2bc}{2}} = \sqrt{bb-bc} \\ &= \sqrt{b \times b - c}; \text{ that is,} \end{aligned}$$

the distance of A from the center of the inscribed circle is $= \sqrt{b \times b - c}$.

PROB. CXI.

Having the sides of an oblique triangle ABC; to find the perpendicular CD, the segments of the base, and the area. 56.

Let AC=a, AB=b, CB=c, $\frac{a+b+c}{2} = z$,
 $a+c=s$, $a-c=d$.

1. For the segments.

AB : AC+CB :: AC-CB : dif. segments
 (Geom. II. 24.); that is, $b : s :: d : \frac{ds}{b} =$
 AD-DB. Then $\frac{1}{2}b + \frac{ds}{2b}$ or $\frac{bb+ds}{2b} =$ great-
 er segment AD, and $\frac{bb-ds}{2b} =$ BD the lesser seg-
 ment.

2. For the perpendicular.

$$CD = \sqrt{aa - \frac{bb+ds}{2b}} = \sqrt{\frac{4bbaa - b^4 - 2bbds - ddss}{4bb}}$$

but $2a=s+d$, and $4aa=s+d$, and $4bbaa=bs+bd$,
 therefore

CD =

Fig.
56.

$$CD = \frac{\sqrt{bbss + 2bbds + bbdd - b^4 - 2bbds - ddss}}{4bb} =$$

$$\frac{\sqrt{ss \times bb - dd - bb \times bb - dd}}{4bb} = \frac{\sqrt{bb - dd} \times \sqrt{ss - bb}}{4bb}$$

$$= \frac{\sqrt{b+d} \times \sqrt{b-d} \times \sqrt{s+b} \times \sqrt{s-b}}{2b}; \quad \text{But } s+b=a$$

+c+b, s-b=a+c-b, and b+d=a+b-c, b-d=b+c-a, therefore

$$CD = \frac{\sqrt{a+b+c} \times \sqrt{a+c-b} \times \sqrt{a+b-c} \times \sqrt{b+c-a}}{2b}$$

$$\text{But } z = \frac{a+b+c}{2}, z-a = \frac{b+c-a}{2}, z-c =$$

$$\frac{a+b-c}{2}, z-b = \frac{a+c-b}{2}. \quad \text{Therefore}$$

$$CD = \frac{\sqrt{2z \times 2z-a \times 2z-c \times 2z-b}}{2b} =$$

$$\frac{2\sqrt{z \times z-a \times z-b \times z-c}}{b}, \quad \text{that is, the perpendicular}$$

$$CD = \frac{\sqrt{a+b+c} \times \sqrt{a+b-c} \times \sqrt{a-b+c} \times \sqrt{b+c-a}}{2b}$$

$$= \frac{\sqrt{b+d} \times \sqrt{b-d} \times \sqrt{s+b} \times \sqrt{s-b}}{2b} =$$

$$\frac{2\sqrt{z \times z-a \times z-b \times z-c}}{b}.$$

3. For the area.

Since the area is = $\frac{1}{2} AB \times CD$ (Geom. II. 10. cor. 2.), and CD was found by the last article, let $CD=p$; since $AD = \frac{aa+bb-cc}{2b}$,

therefore.

$$\text{therefore } CD = \sqrt{\frac{aa+bb-cc}{2b}} = \frac{\sqrt{2aabb+2aacc+2bbcc-a^2-b^2-c^2}}{4bb}; \text{ therefore}$$

$$\text{we have the area of the triangle } ACB = \frac{1}{2}pb =$$

$$\frac{1}{2}\sqrt{2aabb+2aacc+2bbcc-a^2-b^2-c^2} =$$

$$\frac{1}{4}\sqrt{a+b+c \times a+b-c \times b+c-a \times a+c-b}$$

$$= \frac{1}{4}\sqrt{b+d \times b-d \times s+b \times s-d} =$$

$$\sqrt{z \times z+a \times z+b \times z+c}$$

PROB. CXII.

Having the sides of an oblique triangle; to find the 57.
radius of the inscribed circle, &c.

1. In the triangle ABC, bisect the two angles A, B, by the lines AF, BE to intersect in O the center of the inscribed circle. From O, C, let fall the perpendiculars OD, CP, upon the base AB. And put $AB=b$, $AC=a$, $CB=c$, $AP=d$, $PB=f$, $CP=p$, and $DO=x$, $DP=y$; then $AD=d-y$, $BD=f+y$.

Then (Geom. II. 25.) $CA:AP::CS:SP$, and $CA+AP:AP::CP:SP$, that is, $a+d:$

$d::p:\frac{pd}{a+d}=SP$. Likewise $c+f:f::p:\frac{pf}{c+f}=LP$. The triangle APS, ADO are similar, and

$d:\frac{pd}{a+d}::d-y:\frac{pd-py}{a+d}=DO=x$. Also the

triangles BPL and BDO are similar, and

$f:\frac{pf}{c+f}::f+y:\frac{pf+py}{c+f}=x=\frac{pd-py}{a+d}$; then multi-

plying

Fig. 57. plying, $apf + dpf + apy + dpy = cpd + fpd - cpy - fpy$,
and transposing, $apy + dpy + fpy + cpy = cpd - apf$;
that is, because $d + f = b$, $apy + bpy + cpy = cpd - apf$,

$$\text{and } y = \frac{cd - af}{a + b + c}. \text{ Whence } x = \frac{pd - py}{a + d} = \frac{pd}{a + d}$$

$$\frac{pcd - pat}{a + d \times a + b + c} = \frac{pda + pdb + pdc - pdc + paf}{a + d \times a + b + c} = \frac{pba + pbd}{a + d \times a + b + c} = \frac{pb}{a + b + c}.$$

And since p may be had various ways, from the last problem; therefore we shall have the radius of the inscribed circle

$$= \frac{bp}{a + b + c} = \frac{1}{2} \sqrt{\frac{a - c + b \times a + c - b \times b + c - a}{a + b + c}}$$

$$= \frac{1}{2} \sqrt{\frac{b + n \times b - n \times s - b}{s + b}} = \sqrt{\frac{z - a \times z - b \times z - c}{z}}$$

$$\text{Where } z = \frac{a + b + c}{2}, s = a + c, n = a - c.$$

2. For the tangent AD.

$$\text{We have } AD = d - y = d - \frac{cd - af}{a + b + c} = \frac{ad + bd + cd - cd + af}{a + b + c} = \frac{ad + bd + a \times b - d}{a + b + c} = \frac{ba + bd}{a + b + c}$$

But (Geom. II. 23.) $bd = \frac{aa + bb - cc}{2}$, therefore

$$AD = \frac{2ab + aa + bb - cc}{2 \times a + b + c} = \frac{a + b - c}{2 \times a + b + c} = \frac{a + b - c}{2}$$

$$\text{the tangent } AD = \frac{a + d}{a + b + c} b = \frac{a + b - c}{2} = z - c.$$

3. For the central distance.

$$AO^2 = AD^2 + DO^2 = \frac{a+d}{a+b+c} \cdot bb + \frac{pp}{a+b+c} \cdot bb =$$

$$\frac{aa + 2ad + dd + aa - dd}{a+b+c} \cdot bb = \frac{2aa + 2ad}{a+b+c} \cdot bb = \frac{2abb}{a+b+c}$$

$\times a+d$. But $d = \frac{aa+bb-cc}{2b}$; therefore $AO^2 =$

$$\frac{2abb}{a+b+c} \times \frac{2ab+aa+bb-cc}{2b} = ab \times \frac{a+b-c}{a+b+c} =$$

$$\frac{a+b+c}{a+b+c} \times \frac{a+b-c}{a+b+c} \times ab = \frac{a+b-c}{a+b+c} \times ab = \frac{z-c}{z} ab.$$

That is, the distance of the center from the angle

A is $= \sqrt{\frac{a+b-c}{a+b+c} \times ab} = \sqrt{\frac{z-c}{z} \times ab}.$

P R O B. CXIII.

Having the sides of a triangle to find the radius of the circumscribing circle. 58.

Let ABC be the triangle, draw the diameter CF of the circumscribing circle, and let CP be perpendicular to AB. Put $AC=a$, $AB=b$, $CB=c$, $CP=p$, $z = \frac{a+b+c}{2}$, $s=a+c$, $d=a-c$, CH or $HF=R$.

Then (Geom. IV. 28), $p : a :: c : 2R$, and $R = \frac{ac}{2p}$. Now since we have the value of p various ways by problem cxi. we shall have the value of R so many ways. Hence R the radius of the circumscribing circle $= \frac{ac}{2p}$.

$$\begin{aligned}
 \text{Fig. 58.} &= \frac{abc}{\sqrt{a+b+c} \times a+b-c \times a-b+c \times b+c-a} \\
 &= \frac{abc}{\sqrt{b+d} \times b-d \times s+b \times s-b} \\
 &= \frac{4\sqrt{z \cdot z-a \cdot z-b \cdot z-c}}{abc} \\
 &= \frac{abc}{4 \times \text{area}}.
 \end{aligned}$$

Cor. Hence r the radius of the inscribed circle :
to R the radius of the circumscribed circle : :

As $z-a \times z-b \times z-c$:
to $\frac{1}{4}abc$.

PROB. CXIV.

59. Given the base of a triangle, and the diameters of the inscribed and circumscribed circles; to find the sides.

Let QRW be the triangle, $QDWB$ the circumscribing circle, DB (perpendicular to QW) its diameter. Draw BR , which will bisect the angle R . Let QS bisect the angle Q , then S is the center of the inscribed circle. Through S draw ASV parallel to QW . Then AP is the radius of the inscribed circle. Draw BV , BW .

Let $BD=a$, $QW=b$, $AP=c$, $BP=v$, $BR=x$,
then $av-vv = \frac{1}{4}bb$, and $v = \frac{a + \sqrt{aa-bb}}{2}$. Let
 $BW=d = \sqrt{av + \frac{1}{4}bb} = \sqrt{av}$. $BA = v + c = s$,
 $BV=p = \sqrt{BA \times BD}$.

The triangles BRD , BPT and BAS are similar, whence $x : a :: v : \frac{av}{x} = BT$; and $x : a ::$

$s : \frac{as}{x} = BS$. But (Geom. IV. 17. cor.) $as = pp$,
and

Sect. VI. PROBLEMS.

417

and $av = dd$; therefore $BT = \frac{dd}{x}$, and $BS = \frac{pp}{x}$. Fig. 59.

Then $RS = x - \frac{pp}{x}$, and $TS = \frac{pp - dd}{x}$. But

(Geom. II. 25.) $TS : SR :: TQ : QR$; and the triangles TQR and BWR are similar (Geom. IV. 12. cor. 2.), and $TQ : QR :: BW : BR$; whence

$TS : SR :: BW : BR$, that is, $\frac{pp - dd}{x} : \frac{xx - pp}{x} :$

$d : x$, whence $ppx - ddx = dxx - dpp$, and $dxx +$

$ddx = dpp + pp^2$, or $d + x \times dx = d + x \times pp$, and

$dx = pp$, whence $x = \frac{pp}{d}$. Then $BR (x) : DR$

$(\sqrt{aa - xx}) :: BP (v) : PT = \frac{v\sqrt{aa - xx}}{x}$;

whence QT, TW are known. Then $BW (d) :$

$BR (x) :: QT : QR :: TW : WR$, the two sides of the triangle.

PROB. CXV.

There is given the base of a triangle, the line that bisects the vertical angle, and the diameter of the circumscribing circle; to find the sides. 60.

Let $AB = b$, EO or $OF = r$, $CD = d$, $HD = x$, $FD = y$.

Then $AD = \frac{1}{2}b + x$, $DB = \frac{1}{2}b - x$, $FH = \sqrt{yy - xx}$.

And (Geom. IV. 20. cor. 2.) $ADB = CDF$, or $\frac{1}{4}bb - xx = dy$. The triangles FDH, FEC are similar, and $y : \sqrt{yy - xx} :: 2r : v + d$, and $yy + dy =$

$2r\sqrt{yy - xx} = 2r\sqrt{y^2 + dy - \frac{1}{4}b^2}$. Which squared is,
E e y² +

Fig. $y^4 + 2dy^3 + ddy = 4rryy + 4rrdy - rrb$, and reduced $y^4 + 2dy^3 + ddy - 4rrdy + rrb = 0$. Then

$$x = \sqrt{\frac{1}{4}bb - dy}.$$

Also $BF = \sqrt{yy - xx + \frac{1}{4}bb}$, and the triangle ADF, CDB are similar, and $AD (\frac{1}{2}b + x) : AF$ or $BF (\sqrt{yy - xx + \frac{1}{4}bb}) :: CD (d) : CB = \frac{d\sqrt{yy - xx + \frac{1}{4}bb}}{\frac{1}{2}b + x}$.

Also the triangles ADC, BDF are similar, and $BD (\frac{1}{2}b - x) : BF (\sqrt{yy - xx + \frac{1}{4}bb}) :: CD (d) : CA = \frac{d\sqrt{yy - xx + \frac{1}{4}bb}}{\frac{1}{2}b - x}$.



S E C T. VII.

Problems in Plain Trigonometry.

P R O B. CXVI.

In the triangle ABC, there is given the angle B, the side AB; and the sum of the sides BC, AC; to find the sides. 61.

LET $AB=d$, $BC+AC=b$, sine $\angle B=s$,
 cof. $=c$, $AC=x$, then $CB=b-x$.

By plain Trigonometry rad. (1) : $AB (d)$: :
 S.PAB or cof. B (c) : $PB=cd$.

Then (Geo. II. 22.) $xx=dd+bb-2bx+xx+2cdx$
 $b-x$, reduced $2bx+2cdx=bb+dd+2bcd$, and
 $x = \frac{bb+dd+2bcd}{2b+2cd}$.

P R O B. CXVII.

In the triangle ACB, there is given the two segments AD, DB, made by the perpendicular, and the angle ACB; to find the rest. 62.

Make $DE=DB$, and draw CE , then put
 $BD=b$, $AD=d$, CB or $CE=y$, $S.ACB=s$,
 $S.ACE=x$; then $AE=d-b$, $AB=d+b$.

By Trigonometry, (in the triangle ACB) AB
 $(b+d)$: $S.ACB (s)$: : $CB (y)$: $\frac{sy}{b+d} = S.CAB$.

Also (in the triangle ACE), $CE (y)$: $S. CAE$
 $(\frac{sy}{b+d})$: : $AE (d-b)$: $S.ACE (x)$, and $x = \frac{d-b}{d+b} s$.

Fig. 62. Then $\frac{ACE+ACB}{2} = ACD$, and $\frac{ACB-ACE}{2} = BCD$. Then $S.ACD : AD :: \text{rad} : AC$. And $S.BCD : BD :: \text{rad} : CB$.

P R O B. CXVIII.

63. In the triangle ABC there is given AB, and the angle B, and the ratio of AC to BC, to find the sides.

Let fall AD on BC (produced); and put $AB=b$, $AC=a$, the ratio of AC to CB as 1 to r , then $CB=ra$, and $\text{cos. } ACB=c$. Then $\text{rad. } (1) : AC (a) :: S.DAC (c) : ca=DC$. (Geom. II. 22.) $bb=aa+rrea+2craa$, whence $aa = \frac{bb}{rr+1+2cr}$, and $a = \frac{b}{\sqrt{rr+1+2cr}}$.

P R O B. CXIX.















64. In the triangle CAB, there is given two sides and the included angle; to find the area.


Let CA, AB and the angle A be given; draw CF perpendicular to AB, and let $AB=b$, $AC=d$, $S. \angle A=s$. Then in the triangle ACF, $\text{rad. } (1) : AC (d) :: S.A (s) : sd=CF$. Then $\frac{CF \times AB}{2} = \text{area}$, or $\frac{sd b}{2} = \text{area}$; that is, half the rectangle of the sides multiplied by the sine of the included angle, gives the area.

P R O B. CXX.

65. Given all the sides of a trapezium, and two opposite angles; to find the area.

Let the angles B, D be given, and through the other two angles A, C, draw the diagonal AC. Let





Let $AB=b$, $BC=c$, $CD=d$, $DA=f$, $S.\angle B=p$, Fig. S.D= q . Then by the last problem, the area of 65.

the triangle $CBA = \frac{bcp}{2}$, and the triangle $CDA = \frac{dfq}{2}$; therefore the trapezium $= \frac{bcp+dfq}{2}$.

PROB. CXXI.

In the triangle WNE , there is given the segment SE , 66. the angle WNE , and the ratio of NE to NW ; to find the sides.

Let WP be perpendicular to EN , and suppose NE to WN as 1 to p , $S.N=s$, $\text{cof. } N=c$, $SE=b$, $NE=x$, then $WN=px$. In the triangle WNP , $\text{rad.}(1) : WN(px) :: S.PWN(c) : PN=pcx$, and by the similar triangles ENS , EWP ; $ES : EN :: EP : EW$, or $b : x ::$

$$x+pcx : \frac{xx+pcxx}{b} = EW. \text{ But (Geom. II. 22.)}$$

$$\frac{x^4+2pcx^2+ppccx^2}{bb} = xx+ppxx+2pcxx, \text{ or}$$

$$\frac{1+2pc+ppcc}{b} \times xx = bb \times \frac{1+pp+2pc}{1+pp+2pc}, \text{ and } x =$$

$$\frac{1+p}{1+p} \sqrt{1+pp+2pc}.$$

Or thus,

Let $t = \text{tang. of } \frac{EWN+E}{2}$, then $WN+NE$

$$(px+x) : WN-NE (px-x) :: t : \frac{tpx-tx}{px+x} =$$

$\frac{p-1}{p+1} t = \text{tang. } \frac{1}{2} \text{ diff. of the angles } W \text{ and } E.$

Whence the angles W , E are known. Then as $\text{cof. } E : b :: \text{rad.} : x$, required.

Fig.

P R O B. CXXII.

67. In the right-angled triangle ABC, there is given, the sum of AB and BC, the angle CDB; likewise $\angle ACD = \angle DCE$ are given; to find CB, &c.

Let $S.ACB = s$, $S.CAB = c$, $CB = x$, $AB + BC = b$, and $BA = b - x$. Then $s : c :: b - x : x$, and $sx = cb - cx$, whence $x = \frac{cb}{s+c}$. Then CB, BA are known. Let m , n be the tangents of BCE, BCD; then $1 : x :: m : BE :: n : BD$.

P R O B. CXXIII.

68. In the triangle ADC, there is given AB, BC; and the angles ADB, BDC; to find AD, DC.

Let $S.ADB = s$, $S.BDC = t$, $S.ADC = p$, $\text{cotang. ADC} = q$, $AB = b$, $BC = c$, $AC = d$, $AD = x$.

By plain Trigonometry, $b : s :: x : \frac{sx}{b} = S.ABD$ or CBD. And $t : c :: \frac{sx}{b} : \frac{csx}{tb} = CD$. Then $AD + CD \left(x + \frac{csx}{tb} \right) : AD - CD \left(x - \frac{csx}{tb} \right) :: \tan. \frac{A+C}{2} (q) : \frac{tbq - csq}{tb + cs} = \text{tang. } \frac{A-C}{2}$. Then $\angle sA$ and C are known; then $p : d :: S.C : AD :: S.A : CD$.

P R O B. CXXIV.

69. In the triangle ABC, there is given AB, the angle C, and CD, which is drawn to the middle of AB; to find the sides.

Draw AF perpendicular to CB, and put AD or $DB = b$, $CD = d$, $S.\angle C = s$, $\text{col.} C = c$, $AC = x$, $BC = y$. Then

Then (Geom. II. 28.) $xx + yy = 2bb + 2dd$. By Fig. Trigonometry $1 : x :: s : sx = AF$, and $1 : x :: 69$.

$c : cx = CF$. Then $BF = y - cx$, and $ssxx + yy - 2cyx + ccxx = 4bb$, subtract this from the first equation, then $xx - ssxx + 2cyx - ccxx = 2dd - 2bb$, that is (because $ss + cc = 1$) $2cyx = 2dd - 2bb$, and

$$2yx = \frac{2dd - 2bb}{c}; \text{ therefore } yy + 2yx + xx = 2bb$$

$$+ 2dd + \frac{2dd - 2bb}{c}, \text{ and } y + x =$$

$$\sqrt{2bb + 2dd + \frac{2dd - 2bb}{c}} = m; \text{ in like manner}$$

$$y - x = \sqrt{2bb + 2dd - \frac{2dd - 2bb}{c}} = n. \text{ Then}$$

$$y = \frac{m+n}{2}, \quad x = \frac{m-n}{2}.$$

PROB. CXXV.

Given the angles of altitude BCA , BDA , the horizontal angle BCD , and the line of station CD ; to find the height AB . 70.

Let $b = \cotang. BCA = 47 : 30$; $c = \cotang. BDA = 40 : 12$; $d = S. BCD = 87.5$; $CD = f = 283.274$ feet; $AB = x$.

Then in the triangle ABC , $1 : x :: b : bx = BC$, and in the triangle ABD , $1 : x :: c : cx = BD$; and

in the triangle BCD , $cx : d :: bx : \frac{db}{c} = S. BDC$

$= 50.39$; whence $DBC = 42.16$; let $n = S. BDC$; then $n : f :: d : BD = cx$, and $ncx = fd$, and

$$x = \frac{fd}{nc} = 355.458.$$

Fig.

P R O B. CXXVI.

71. Given the sum of the sides of a triangle, and all the angles severally; to find the sides.

Let S.A= s , S.B= n , S.C= t , AB+BC+CA= b , AC= x . Then by Trigonometry, $n : x :: s :$

$\frac{sx}{n} = CB$, and $n : x :: t : \frac{tx}{n} = AB$. And $x +$

$$\frac{sx}{n} + \frac{tx}{n} = b, \text{ whence } x = \frac{nb}{n+s+t}.$$

P R O B. CXXVII.

72. In the right-angled triangle VAB, there is given the perpendicular AB, the segment VC, and the angle VAC; to find CB.

Let AB= b , VC= c , tang. VAC= t , BC= a .

Then by plain trigonometry, $b : 1 :: a : \frac{a}{b} =$

tang. BAC, and (trig. viii.) $1 - \frac{ta}{b} : 1 :: t + \frac{a}{b}$

$: \frac{bt+a}{b-ta} = \text{tang. BAV}$. Whence $1 : b :: \frac{bt+a}{b-ta}$

$: a+c = \frac{bbt+ba}{b-ta}$, and multiplying, $ba+bc -$

$taa-tca=bbt+ba$, reduced $aa+ca = \frac{bc}{t} - bb$.

P R O B. CXXVIII.

73. In the right-angled triangle ABC, BE=EC, and $\angle ABD = \angle CBD$; and there is given BD and $\angle CAE$; to find the sides.

Draw DF parallel to CB; then in the triangle DFB, the angles at B, D are 45° , and BD being given,

given, DF and FB its equal, are given; and since Fig. CE=EG, therefore DG=GF. Let DG or GF=b, 73.

S.DAG=s, AF=x; then AG= $\sqrt{bb+xx}$, AD= $\sqrt{4bb+xx}$. And by plain Trig. AG $\sqrt{(bb+xx)}$

: rad. (1) :: AF (x) : $\frac{x}{\sqrt{bb+xx}}$ = S.G. Also

S. G ($\frac{x}{\sqrt{bb+xx}}$) : AD ($\sqrt{4bb+xx}$) :: S.DAG

(s) : DG (b); whence $\frac{bx}{\sqrt{bb+xx}} = s\sqrt{4bb+xx}$,

reduced $xx^2 + 5bbxx + 4b^4 = 0$.

$$\frac{bbxx}{ss}$$

P R O B. CXXIX.

From the point B, to draw the lines BC, BD, BA, 74.
so that CD, DA, and the angles CBD, DBA,
may be given.

Draw CF perpendicular to AB, and put CD=b,
DA=c, CA=s, and S.CBD=f, S.DBA=d,
S.CBA=m, cof. CBA=n, and CB=x.

Then by plain Trigonometry 1 : x :: n : nx = BF,

and b : f :: x : $\frac{fx}{b}$ = S.D, and d : c :: $\frac{fx}{b}$: $\frac{cfx}{bd}$ =

BA. But in the triangle CBA, (Geom. II. 23.)

ss = xx + $\frac{ccffxx}{bbdd}$ - 2nx \times $\frac{cfx}{bd}$, which reduced

is $x = \frac{bds}{\sqrt{bbdd + ccff - 2bdjnc}}$

Fig.

P R O B. CXXX.

75. In an oblique triangle there is given, the base, and perpendicular, and angle opposite to the base; to find the sides.

Draw AD perpendicular to CB; and put $AB=b$, $S.ACB=s$, $\text{cof. } ACB=c$, $\text{perp. } CF=p$, $AC=x$, $CB=y$.

In the triangle ACD, $1 : x :: s : sx = AD$, and $1 : x :: c : cx = CD$. The triangles ABD and CBF are similar, and $y : b :: p : sx = AD$, whence $pb = sxy$. In the triangle ABC (Geom. II. 23.),

$$bb = xx + yy - 2cxy; \text{ but } xy = \frac{pb}{s}, \text{ and } y = \frac{pb}{sx};$$

$$\text{therefore } bb = xx + \frac{ppbb}{ssxx} - \frac{2cpb}{s}; \text{ which reduced is}$$

$$x^4 - \frac{bbxx}{s} + \frac{bbpp}{ss} = 0.$$

Otherwise,

Let $AC + CB = x$, $AC - CB = y$, the rest as before; then $AD = s\sqrt{xx + y}$, $CD = c\sqrt{xx + y}$, $\frac{pb}{s} = \sqrt{xx - yy}$.

Then in the triangle ABC, $bb = x + y + \frac{x - y}{s} - 2c \times \sqrt{xx - yy}$; that is, $2xx + 2yy - 2c\sqrt{xx + y} + 2cy = bb$,

and putting $\sqrt{xx - \frac{pb}{s}}$ for yy , we have $2xx + 2\sqrt{xx - \frac{pb}{s}} -$

$$\frac{2pb}{s} - 2c\sqrt{xx + y} + 2cy = bb, \text{ or } 4xx - \frac{2pb}{s} -$$

$$\frac{2cpb}{s} = bb, \text{ whence } x = \sqrt{\frac{1}{4}bb + \frac{1+c}{2s} \times bp} = m,$$

$$\text{and } y = \sqrt{\frac{1}{4}bb + \frac{c-1}{2s} bp} = n. \text{ Then } AC = \frac{m+n}{2}$$

$$BC = \frac{m-n}{2}.$$

Or

Or thus,

Fig.

Let $f = \text{cofine of the sum of the angles } A, B;$
 $v = \text{cof. of their difference; the rest as before.}$
 Then (Trig. II. 10.) $v - f : s :: p : \frac{1}{2}b$, and $v - f =$
 $\frac{2ps}{b}$, and $v = \frac{2ps}{b} + f$. Then the angles A and B will
 be known, and consequently their opposite sides.

75.

PROB. CXXXI.

Given all the sides of a triangle, to find the center of
 the circumscribed circle. 76.

On the middle of AB, AC, erect the perpendi-
 culars DO, FO, the point of interfection O, is the
 center of the circumscribing circle. From O draw
 OI, OG, parallel to AC, AB; and put $AB = b,$
 $AC = d,$ $S. \angle A = s,$ $\text{cof. } A = c,$ $AI = x,$ $IO = y.$
 The $\angle CGO = \angle CAB = \angle OID;$ and in the right-
 angled triangles OI, OGF, it will be $1 : y ::$
 $c : cy = ID,$ and $1 : y :: s : sy = OD.$ Al-
 so $1 : x :: c : cx = GF,$ and $1 : x :: s : sx =$
 $FO.$ Then $x = \frac{1}{2}b - cy,$ and $y = \frac{1}{2}d - cx,$ and
 $cy = \frac{1}{2}b - x = \frac{1}{2}cd - cex,$ and $x - cex = \frac{1}{2}b - \frac{1}{2}cd,$
 or $sx = \frac{b - cd}{2},$ whence $x = \frac{b - cd}{2ss},$ and $y = \frac{d - cb}{2ss}.$

Therefore $DO = \frac{d - cb}{2s},$ and $FO = \frac{b - cd}{2s}.$ Like-

wise $AO = \frac{\sqrt{bb + dd - 2bcd}}{2s}.$

PROB. CXXXII.

In the given triangle ABC, the angles AOC, COB, 77.
 $\angle BOA,$ about the point O, are given; to find the
 distances AO, BO, CO.

Produce CO to D, and BO to E. And let
 $AB = b,$ $AC = d,$ $CB = f,$ $S. A = s,$ $\text{cof. } A = c,$
 $S. B = q,$

Fig. S.B=q, cof. B=n, S.AOE=g, cof. AOE=m,
77. S.AOD=b, S.DOB=p, AO=x.

Then by Trigonometry, AB (b) : S.O (g) ::
AO (x) : $\frac{gx}{b}$ = S.ABO, and $\sqrt{1 - \frac{ggxx}{bb}}$ = cof.
ABO=y; and (Trig. I. 6. cor.) $\frac{ggx}{b} + my$ = cof.
OAB; and (I. 6.) $gy - \frac{mgx}{b}$ = S.OAB. Also
(Trig. I. 6.) $\frac{sggx}{b} + smy - cgy + \frac{cmgx}{b}$ = S.CAO;
and (I. 6.) $qy - \frac{gnx}{b}$ = S.CBO. And by Trigonometry,
 $p : f :: qy - \frac{gnx}{b} : \frac{fqy}{p} - \frac{gnfx}{bp}$ = CO, and
 $b : d :: S.CAO : CO = \frac{sggd}{bb} + \frac{dsmy}{b} - \frac{dcgy}{b} +$
 $\frac{dcmgx}{bb} = \frac{fqy}{p} - \frac{gnfx}{bp}$, and multiplying, $sggdpx +$
 $bpdsmy - bpdscy + pdcmgx = bbfqy - gnfbx$; and transposing,
 $bbfqy + bpdscy - bpdsmx = sggdx + pdcmgx +$
 $gnfbx$, and $y = \frac{sggd + pdcmg + gnfb}{bbfq + bpdcg - bpdsm} x = \frac{rx}{t}$ by substitution,
that is, $\sqrt{1 - \frac{ggxx}{bb}} = \frac{rx}{t}$, and $1 - \frac{ggxx}{bb}$
 $= \frac{rrxx}{tt}$, and $bbtt - ggttxx = bbxx$, reduced
 $x = \frac{bt}{\sqrt{bb + ggtt}}$.

Or thus,

78. Make the angle BAF = supplement of BOC,
and $\angle ABF = \text{sup. } \angle AOC$; through A, B, F describe the circle AOBF, to intersect CF in O, the point required.

Calculation.

Sect. VII. PROBLEMS.

429

Calculation. In the triangle ABF, all the angles are given, and the side AB, to find AF. Fig. 78.

In the triangle CAF; CA, AF, and $\angle CAF$ are given; to find $\angle ACF$.

In the triangle ACO, there is given AC, and all the angles; to find AO, CO.

PROB. CXXXIII.

In the right-angled triangle ABC, there is given BA, and angle CBD; also $AT=TD$, and $\angle ABT = \angle CBT$; to find AC. 79.

Let $BA=b=95.23$, $\text{tang. } DBC=t = T. 15^\circ$, AT or $DT=a$, then $AD=2a$. By trigonometry,

$b : a :: 1 : \frac{a}{b} = \text{tang. } TBA$, and $b : 2a : 1 : \frac{2a}{b} = \text{tang. } DBA$. Then (Trig. I. 2. Schol.) the tang.

$2TBA$ or $\text{tang. } ABC = \frac{2ab}{bb-aa}$. Also (Trig. I. 8.)

$1 - \frac{2ta}{b} : 1 :: t + \frac{2a}{b} : \frac{bt+2a}{b-2ta} = \text{tang.}$

$\frac{ABD + DBC}{b-2ta} = \text{tang. } ABC$. Whence $\frac{2ab}{bb-aa} =$

$\frac{bt+2a}{b-2ta}$, reduced $2a^2 - 3btaa = bt^2$, and $a = 65.13$,

and $\frac{2ab}{bb-aa} \times b = AC$.

PROB. CXXXIV.

Upon a horizontal plane, there stands a tall pine-tree leaning towards the south. A man standing on the north side of it 50 yards from the foot, finds the tree to subtend an angle of 39° . Afterwards going directly west 73 yards, it subtends an angle of 46° . What is the tree's length?

Let AC be the tree; E, F, the two stations. Draw AB perpendicular to the horizon, and AD perpen- 80.

Fig. perpendicular to FC produced; draw BD, AC. The triangles ABD, ABC, ADC, BDC, CEF, DAF, BAE, are all right-angled. Put $EF=b$, $CE=d$, $CF=c$, $CD=y$, $\text{tang. AEB}=s$, $\text{tang. AFC}=t$.

By Trigonometry, $1 : t :: c+y : c+y \times t = AD$,
whence $AC = \sqrt{yy + tt \times c+y}$. The triangles

FCE, BCD are similar, and $d : c :: y : \frac{cy}{d} = BC$,

and $d : y :: b : \frac{by}{d} = BD$. Then $AB =$

$$\sqrt{yy + tt \times c+y} - \frac{ccy}{dd} = \sqrt{tt \times c+y} - \frac{bby}{dd}$$

And in the triangle ABE, $d + \frac{cy}{d} : 1 ::$

$$\sqrt{tt \times c+y} - \frac{bby}{dd} : s, \text{ whence } \frac{dds + csy}{d} =$$

$$\sqrt{tt \times c+y} - \frac{bby}{dd}; \text{ and squared } d^4ss + 2cddssy +$$

$$ccssyy = dd \times ttcc + 2ttcy + tyy - bbyy, \text{ and reduced}$$

$$ccssyy + 2cddssy = ddtcc.$$

$$-ddt - 2cdatt - d^4ss$$

$$+ bb$$
















P R O B. CXXXV.


81. Given two altitudes and two azimuths of a cloud in motion; to find the point of the wind.

Let A be the first, B the second place of the cloud, O the place of observation, ABC the plane of the cloud's motion; AB its line of direction. Let AD, BE, CO be perpendicular to the horizon, then DEO is equal and parallel to ABC, and MDE is the path of the cloud on the earth. Let OM be the meridian.

Put $p = \text{tang. AOD}$, $q = \text{tang. BOE}$, $t = \text{cotang. } \frac{1}{2} \text{DOE}$, $OD = x$. In the triangle AOD, $1 : x :: p : px = AD = BE$; and in the triangle BOE,

$q :$





PLATE

$q : px : 1 : \frac{px}{q} = OE$. In the triangle DOE , (Trig. Fig. 81.

II. 6.) $\frac{px}{q} + x : \frac{px}{q} - x :: t : \frac{p-q}{p+q} t = \text{tang.}$

$\frac{ODE - OED}{2}$, and the sum and difference of ODE , OED being had; ODE , and OED will be known.

In the triangle ODM , there is given ODM and DOM , therefore OMD is known, which is the way of the cloud or of the wind.

PROB. CXXXVI.

On a clear day, the wind standing N. N. E. I observed a small cloud W. by S. whose altitude was 41° , and whilst the shadow of the cloud moved over 1230 yards upon a horizontal plane, the cloud itself moved through an angle of $9^\circ : 37'$ as I observed it with an instrument. What was the cloud's height?

Let E be the place of observation, CA the tract of the cloud, FG its projection upon the horizon; AF, CG, BE being perp. to the horizon. Let BD be perp. to ACD , and AK to ECK . 82.

Let AC or $GF = d = 1230$, $S.DCB = c = 5$ points, $\text{cof. } DCB = n$, $S.GEC = b = 41^\circ$, $\text{cof. } GEC = s$, $\text{tang. } AEC = t = 9^\circ : 37'$; $CG = x$. By Trigonometry, in the triangle CGE , $b : x :: 1 : \frac{x}{b} = CE$, and $b : x$

$:: s : \frac{sx}{b} = CB$, and in the triangle BCD , $1 : \frac{sx}{b} :: c$

$: \frac{csx}{b} = DB$, and $1 : \frac{sx}{b} :: n : \frac{nsx}{b} = CD$. Then DE

$= \sqrt{xx + \frac{ccssxx}{bb}} = \frac{x}{b} \sqrt{bb + ccss} = \frac{x}{b} p$, by substitution.

The

Fig. The triangles CED and CAK are similar, and

$$82. \quad \frac{x}{b} (CE) : \frac{px}{b} (DE) :: (AC) d : pd = AK; \text{ and } \frac{x}{b}$$

$$(CE) : \frac{snx}{b} (CD) :: d (AC) : snd = CK. \text{ And in the}$$

$$\text{triangle AKE, } snd + \frac{x}{b} (EK) : 1 (\text{rad.}) :: pd (AK)$$

$$: t (\text{tang. AEK}), \text{ whence } sndt + \frac{tx}{b} = pd, \text{ and } tx =$$

$$bpd - sndtb, \text{ and } x = \frac{bpd - sndtb}{t} = \frac{bdp}{t} - sndb.$$

P R O B. CXXXVII.

83. In the triangle ACB, there is given AC, CB, the segment AD, and the angle DCB; to find the rest.

Draw AF perpendicular to CB; and put AC = a, DB = b, CB = d, S.ACD = s, cof. ACD = c, and S.CDB = x.

In the triangle CDB, $d : x :: b : \frac{bx}{d} = \text{S.DCB}$, and

in the triangle CAD, $x : a :: s : \frac{as}{x} = \text{AD}$, then AB

$$= b + \frac{as}{x} = \frac{bx + as}{x}. \text{ But (Trig. I. 5. cor. 1.)}$$

$$c \sqrt{1 - \frac{bbxx}{dd}} - \frac{sbx}{d} = \text{cof. ACB} = cz - \frac{sbx}{d} \text{ (putting}$$

$$z = \sqrt{1 - \frac{bbxx}{dd}}); \text{ and in the triangle ACF, } 1 : a$$

$$:: cz - \frac{bsx}{d} : cza - \frac{absx}{d} = \text{CF}. \text{ But (Geom. II.}$$

$$23.) \frac{bx + as}{xx} = aa + dd - 2d \times caz - \frac{absx}{d}, \text{ and}$$

multiplying

S E C T. VIII.

Problems in spherical Trigonometry.

P R O B. CXXXVIII.

*Given the latitude of the place and the sun's longitude;
to find the ascensional difference.*

84. **L**ET $b = S.$ greatest declination, $c = S.$ sun's longitude AD, $x = S.$ declination BD, $t =$ tang. latitude PCH.

Then in the r^t \angle spherical triangle ADB, rad.
(1) : S.AD (c) :: S.A (b) : $bc = S.$ BD, and (Trig.

I. 1. schol.) tang. BD = $\frac{bc}{\sqrt{1-bbcc}}$; and in the tri-
angle CBD, rad. (1) : cotang. C (t) :: tang. BD

$\left(\frac{bc}{\sqrt{1-bbcc}}\right)$: S.CB (x), and $x = \frac{bct}{\sqrt{1-bbcc}}$

P R O B. CXXXIX.

84. *Given the sun's declination, and the sum of the latitude
and amplitude; to find each of them.*

Let $a =$ sine of half the sum of CD and PH, $b =$
the cosine, $d = S.$ declination, $s = S.$ sum of CD and
PH, $x = S.$ half their difference, $y = S.$ whole diffe-
rence. Suppose PH greater than CD. Then
(Trig. I. 6. schol.) $b\sqrt{1-xx-ax} = \text{cos. PH}$; and
 $a\sqrt{1-xx-bx} = S.CD$. But in the triangle CBD,
cos. C ($b\sqrt{1-xx-ax}$) : SBD (d) :: rad. (1) :
S.CD

Sect. VIII. SPHERICAL TRIG. PROBLEMS.

435

S.CD $(a\sqrt{1-xx}-bx)$. Whence $ab \times \sqrt{1-xx} -$ Fig.
 $aa\sqrt{1-xx} - bb\sqrt{1-xx} + abxx = d$; 84.

or $ab - x\sqrt{1-xx} = d$ (because $aa+bb=1$),

But (Trig. I: 2. schol.) $2x\sqrt{1-xx}=y$, and $2ab$
 $=s$. Whence $s-y=2d$, and $y=s-2d$.

P R O B. CXL.

Given the sun's altitude at six, and also when west; to find the latitude. 85.

R is the sun's place when west, and O at six a clock. Let $t=S.RC$ the altitude west, $s=S.OI$ the altitude at six, $x=S$. latitude.

In the triangle CIO, S.C (x): S.OI (s) :: rad.

(1): $\frac{s}{x} = S.CO$ the declination. In the triangle

DCR, S.C (x): S.DR ($\frac{s}{x}$) :: rad. (1): S.CR

(1), and $tx = \frac{s}{x}$, or $xx = \frac{s}{t}$, and $x = \sqrt{\frac{s}{t}}$.

P R O B. CXLI.

AEC, BCD are two triangles right-angled at E and D, and standing on the great circle ECD; also $AC=CB$, and EC, CD, and the angle ACB are given; to find the angles and sides. 86.

Let $a=tang. DC$, $b=tang. CE$, $s=S$. half the sum of the angles, BCD, ACE; $c=cosine$, $x=$ sine, $y=cosine$ of half the difference. Then $sx+cy=cos.$ lesser ACE, and $cy-sx=cos.$ greater BCD. And in the triangle ACE, $cy+sx:1::b$

$\frac{a}{cy+sx} = tang. AC$. And in the triangle BCD,

F f z

cy-

Fig. 86. $c y - s x : 1 :: a : \frac{a}{c y - s x} = \text{tang. CB.}$ Whence

$$\frac{a}{c y - s x} = \frac{b}{c y + s x}, \text{ and } a c y + a s x = b c y - b s x, \text{ and } a s x$$

$$+ b s x = b c y - a c y, \text{ and } \frac{x}{y} = \frac{b c - a c}{b s + a s} = \text{tang. half the}$$

difference of the angles BCD, ACE; whence the angles themselves are had.

P R O B. CXLII.

87. *Given the sun's amplitude, and altitude at six; to find the latitude, and declination.*

Let P be the pole, Z the zenith, CB the amplitude, AP the altitude at six.

Let S.CB = b , p = S.AP, x = S. lat. PO, y = S. twice the latitude. In the triangle CBD, rad. (1) : S.C ($\sqrt{1 - x x}$) :: S.CB (b) : $b \sqrt{1 - x x}$ = S. BD or AC. And in the triangle CAP, rad. (1) : S.AC ($b \sqrt{1 - x x}$) :: S.C (x) : S.AP (p); there-

fore $b x \sqrt{1 - x x} = p$, and $2 x \sqrt{1 - x x} = \frac{2 p}{b}$,

but $y = 2 x \sqrt{1 - x x}$; whence $y = \frac{2 p}{b}$; then x will be known, and $b \sqrt{1 - x x}$, the declination.

P R O B. CXLIII.

88. *Given two altitudes and two azimuths of the sun; to find the latitude.*

Let Z be the zenith, P the pole; S, O two places of the sun. Let s, f = sine and cosine of ZS; p, q = sine and cosine of ZO; m = cof. PZS, n = cof. PZO; x, y = sine and cosine of PH. Then (Trig. II. 38.) cof. SP = $s y m + f x$, and cof. OP =

Sect. VIII. PROBLEMS.

437

OP = $pyn + qx$. Whence $ym + fx = pyn + qx$, and Fig. 88.
 $fx - qx = pyn - ym$; therefore $\frac{x}{y} = \frac{pn - sm}{f - q} =$
 tang. PH the latitude.

PROB. CXLIV.

Given the latitude of the place; and the sun's altitude is equal to his azimuth from the south, and equal to the hour from noon; to find any of them.

89.

Z is the zenith, P the pole, ZP is given, and $\angle ZPO = AZO = DO$. Let $s = S.ZP$, $c = \cos ZP$,
 $y = S.ZPO = S.AZO$. Then $\sqrt{1 - yy} = S.ZO$,
 $y : \sqrt{1 - yy} :: y : \sqrt{1 - yy} = S.OP$, and $y = \cos$
 OP. But (Trig. III. 38.) $s\sqrt{1 - yy} \times \sqrt{1 - yy} +$
 $cy = y$, or $s - yy + cy = y$, and $yy + y = s$, or

$$yy + \frac{1 - c}{s} y = 1.$$

PROB. CXLV.

There are two places, whose latitudes are the complements of each other to 90° , and the sun's declination being given, he rises an hour sooner in one place than the other; to find the latitudes.

87.

Let $t = \text{tang. declination}$, $b = \text{tang. ascensional difference}$, $x = \text{tang. one latitude}$, then $\frac{1}{x} = \text{tang. the other latitude}$. In the triangle CDB, rad. (1) : cotang. DCB (x) :: T.DB (t) :: $tx = S.DC$ the ascensional difference in the first latitude, and $1 : \frac{1}{x} :: t : \frac{t}{x} =$ the ascensional difference in the other latitude. But (Trig. I. 9) $1 + t : 1 ::$

Fig. 89. $tx - \frac{t}{x} : b$, and $tx - \frac{t}{x} = b + btt$, $txx - t = bx + bttx$, and $xx - \frac{btx}{t} = 1$.

$$-\frac{b}{t}$$

P R O B. CXLVI.

The stile of an horizontal dial being turned down, fell upon the hour line of 8; query, the latitude it was made for?

Let $t = \text{tang. of } 4 \text{ hours or } 60^\circ$, answering to 8 a clock, $x = \text{S. latitude}$; then $\frac{x}{\sqrt{1-xx}} = \text{tang. latitude} = \text{hour angle of } 8$, by the question. Whence by the known proportion of dialling,

$$1 : x :: t : \frac{x}{\sqrt{1-xx}}, \text{ and } tx = \frac{x}{\sqrt{1-xx}}, \text{ or,}$$

$$t\sqrt{1-xx} = 1, \text{ and } \sqrt{1-xx} = \frac{1}{t} = \text{cof. lat.}$$

P R O B. CXLVII.

To find in what latitude, an erect south declining dial may be made, so that the declination of the plane, the distance of the substile from the meridian, and the stile's height, are all equal.

90. Let ABC be the right-angled spherical triangle, in which are found all the requisites; viz. $AB = \text{co-lat.}$ $\angle A = \text{co-declination}$, $\angle B = \text{plane's distance from the meridian}$, $CB = \text{stile's height}$, $AC = \text{substile's distance from the meridian}$.

Let $x = \text{S. AB}$, $y = \text{S. BC}$. Then (by the properties of right triangles) $1 : \text{tang. BC} :: \text{cotang. A} : \text{S. AC}$, or $\text{S. AC} = \text{tang. BC} \times \text{cotang. A}$. But by

by the question, $BC=AC=\text{comp. } A$. Therefore Fig.

$\text{tang. } BC \text{ or } \text{cotang. } A = \frac{y}{\sqrt{1-yy}}$. Whence 90°

$$y (\text{S.AC}) = \frac{y}{\sqrt{1-yy}} \times \frac{y}{\sqrt{1-yy}} (\text{tang. } BC \times$$

$$\text{cotang. } A) = \frac{yy}{1-yy}, \text{ and } 1-yy=y, \text{ or } yy+y=1,$$

$$\text{whence } y = \frac{\sqrt{5-1}}{2}.$$

Again, in the same triangle, $1 : \text{cos. } AC ::$
 $\text{cos. } BC : \text{cos. } AB$, whence $\text{cos. } AB = \text{cos. } AC \times$
 $\text{cos. } BC$; that is, $\sqrt{1-xx} = \sqrt{1-yy} \times \sqrt{1-yy}$
 $= 1-yy$, therefore $\sqrt{1-xx}=y$. And since AC
 $= BC$, therefore $\angle B = \angle A$. Hence all these five
 are equal; 1. Plain's declination. 2. Distance of
 the substile from the meridian. 3. Stile's height.
 4. Latitude of the place. 5. Comp. of the plane's
 diff. longitude; and each of them = twice the
 sine of $18^\circ = .618034$; whence the latitude and
 declination = $38^\circ : 10\frac{1}{3}$.

PROB. CXLVIII.

*Given the sun's meridian altitude, and also his altitude
 at two; to find the latitude.*

Let Z be the zenith, P the pole; B, O two 91.
 places of the sun. Let $a, b = \text{fine and cosine of}$
 BZ ($PO - PZ$); $x, y = \text{fine and cosine of } PO +$
 PZ , $c = \text{cos. } P$, $d = \text{cos. } ZO$. Then (Trig. I. 6.
 schol.) $ay + bx = \text{S.PO}$, $by - ax = \text{cos. } PO$, $ay - bx$
 $= \text{S.PZ}$, $by + ax = \text{cos. } PZ$. And in the triangle
 OPZ (Trig. III. 38.) $ay + bx \times ay - bx \times c = by - ax$
 $\times by + ax = d$; that is, $caay - cbbx + bby - aax = d$,
 but $yy = 1 - xx$, therefore $caa - caax - cbbx + bb$
 $- bbbx - aax = d$, but $aa + bb = 1$, whence $caa -$
 $cx + bb - xx = d$, and $xx - cx = caa + bb - d$, and

Fig.

91. $x = \sqrt{\frac{caa + bb - d}{1 - c}}$. Whence PO (or PB), and PZ are had.

P R O B. CXLIX.

92. In the spherical triangle VAC, there is given the perpendicular AB, the angle A, and the base VC; to find the segments.

Put $b = S.AB$, $t = \text{tang. VAC}$, $c = \text{tang. VC}$, $x = \text{tang. BC}$. In the triangle BAC, $b : 1 :: x :$

$\frac{x}{b} = \text{tang. BAC}$, and (Trig. I. 9.) $1 + cx : 1 ::$

$x : \frac{c - x}{1 + cx} = \text{tang. VB}$, and $1 + \frac{tx}{b} : 1 ::$

$x : \frac{tb - x}{b + tx} = \text{tang. VAB}$. And in the tri-

angle VAB, $1 : b :: \frac{tb - x}{b + tx} : \frac{c - x}{1 + cx}$. Whence

$\frac{c - x}{1 + cx} = \frac{tb - x}{b + tx} b$. And multiplying,

$bc - bx + tcx - txx = tb^2 - bx + bbctx - cbxx$, and reduced, $t - bc.xx + bbct - tc.x = bc - tbb$.

P R O B. CL.

Travelling in an unknown part of the world, I found by chance an old horizontal dial, whose hour lines were so decayed by length of time, that I could only discover those of 4 and 5; whose distance I found just 21 degrees; to find the latitude of the place.

Let $b = \text{tang. } 60$ the hour arch of 4.
 $d = \text{tang. } 75$ the hour arch of 5.
 $t = \text{tang.}$ of their difference 21.
 $x = S.$ latitude.

Then

Then by the known proportion of dialling, rad: Fig. S. lat. :: tang. hour arch : tang. hour angle, that is,

$$1 : x :: b : bx = \text{tang. hour } \angle \text{ of } 4.$$

$$1 : x :: d : dx = \text{tang. hour } \angle \text{ of } 5.$$

But (Trig. I. 8.) $1 - bix : 1 :: 1 + bx : dx$, whence

$$1 + bx = dx - bdtxx, \text{ and } bdtxx + b - d \cdot x + 1 = 0.$$

In numbers $2.48133x^2 - 2x = -.383864$.

and $x = .49084 = \text{S. } 29 \text{ } 24 \text{ the lat.}$

or $x = .31518 = \text{S. } 18 \text{ } 22 \text{ the lat.}$

PROB. CLI.

There are given the latitudes of three places lying in the arch of a great circle; and the diff. longitude are equal, between the middle one and each of the extrem places; to find their distances.

Let P be the pole, AE the equinoctial; G, V, M, the three places. Put $b = \text{tang. MD}$, $c = \text{tang. VB}$, $d = \text{tang. CG}$, $x = \text{S. AB}$, $y = \text{S. CB}$ or BD .

Then (Trig. I. 5.) $x\sqrt{1-y} + y\sqrt{1-xx} = \text{S. AD}$; and (Trig. I. 6.) $x\sqrt{1-y} - y\sqrt{1-xx} = \text{S. AC}$.

Then (Trig. III. 27. cor. 1.) $x : c :: x\sqrt{1-y} + y\sqrt{1-xx} : b$; and $x : c :: x\sqrt{1-y} - y\sqrt{1-xx} : d$. Whence $\frac{bx}{c} = x\sqrt{1-y} + y\sqrt{1-xx}$, and

$$\frac{dx}{c} = x\sqrt{1-y} - y\sqrt{1-xx}.$$

Then adding and subtracting these last equations; $\frac{bx + dx}{c} = 2x\sqrt{1-y}$, and $\frac{bx - dx}{c} = 2y\sqrt{1-xx}$; by the

former $\sqrt{1-y} = \frac{b+d}{2c} = \text{cos. CB or BD}$. Hence y is known. Therefore in the triangles GBV, VPM,

Fig. two sides and the included angle are given, to find
93. GV, VM, the distances required.

Or thus,

(By Trig. III. 44. cor.) As rad: $\frac{1}{2}$ tang. PV ::
tang. GC + tang. MD : cof. GPV or MPV.

P R O B. CLII.

94. In the spherical ABC, we have given the angles ADC,
CDB, BDA, about the point D; and the triangle
ABC itself; to find the distances AO, BO, CO.

Let s, c be the sine and cosine of A ; p, q = sine
and cof. B , $a = S.CDB$, $f = S.CB$, $b = S.CDA$, $d =$
 $S.AC$, $h = \text{cof. } AB$, $n = \text{cof. } ADB$, x, v = sine and
cof. DAB ; y, z = sine and cof. DBA .

Then (Trig. I. 6.) $sv - cx = S.CAD$, and $px -$
 $qy = S.CBD$, and in the triangles $CAD, BAD, b:$

$$d :: sv - cx : \frac{dsv - dcx}{b} = S.CD ; \text{ and } a : f :: pz -$$

$$qy : \frac{fpz - fqy}{a} = S.CD ; \text{ therefore } adsv - adcx =$$

$$bfpz - bfqy.$$


In the triangle ADB (Case 10. Trig.) $bxy - vx = n$.


But $v = \sqrt{1 - xx}$, $z = \sqrt{1 - yy}$, therefore

$$ads\sqrt{1 - xx} - adcx = bfp\sqrt{1 - yy} - bfqy.$$

$$\text{and } bxy - \sqrt{1 - xx} \times \sqrt{1 - yy} = n.$$

From these two equations, the roots may be
easiest found by problem xcv; otherwise it will
ascend to a high equation: or if you please you
may proceed by rule 5, prob. xcii.





PROB. CLIII.

Given the difference of the azimuths of three known stars; to find their altitudes.

Let Z be the zenith; A, B, C, the stars. Since their places are given, the triangle ABC will be given. Put $s, c = \text{fine}$ and $\text{cos. } \frac{1}{2}ABC$, $a = S.AB$, $b = S.BC$, $n = \text{cos. } AC$, $q = \text{cos. } AZC$, $d = S.AZB$, $e = S.CZB$; $x, y = \text{fine}$ and $\text{cos. } \frac{CBZ - ABZ}{2}$. Then 95
 $sy + cx = S.CBZ = v$, and $sy - cx = S.ABZ = z$. And in the triangles ABZ, CBZ, $d : a :: z : \frac{az}{d} = S.AZ$, and $b : e :: v : \frac{ve}{b} = S.CZ$; and in the tri-

angle AZC (Trig. III. 38.) $\frac{qazve}{db} + \sqrt{1-zz} \times \sqrt{1-vv} = n$; and transposing, $\sqrt{1-zz} \times \sqrt{1-vv} = n - \frac{qazve}{bd}$; and squaring, $1-zz-vv + vvzz = nn - \frac{2nqae}{bd} vx + \frac{qqaee}{bbdd} v^2z^2$; and transposing, $1-nn = zz + vv - \frac{2nqae}{bd} vx + \frac{qqaee - bbdd}{bbdd} v^2z^2$, but $z^2 + v^2 = 2ssy^2 + 2ccxx$, and $vx = ssyy - ccxx$. Therefore $1-nn = 2ssyy + 2ccxx - \frac{2nqaess}{bd} yy + \frac{2nqaecc}{bd} xx + \frac{qqaee - bbdd}{bbdd} \times$

$ssy^4 - 2s^2c^2x^2y^2 + c^4x^4$. Let $1 - nn = mm$, $2 - \frac{2nqae}{bd} = f$, $2 + \frac{2nqae}{bd} = g$, $\frac{qqaee}{bbdd} - 1 = p$; then $mm = fssyy + ccgxx + ps^4y^4 - 2pc^2s^2y^2x^2 + pc^4x^4$, but $yy = 1 - xx$; therefore expunging y , and reducing, $px^4 + gcc - fss - 2ps^2xxx = mm - fss - ps^4$.

P R O B. CLIV. 9

Given the sun's declination, the difference of two altitudes, the difference of azimuths, and the difference of times; to find the altitudes, and the latitude.

96. Let P be the pole, Z the zenith; A, B, the places of the sun. Put $d = S.$ sun's declination, $s = S.$ $\frac{1}{2}APB$, $c = \text{cos. } \frac{1}{2}AZB$; $p, q = \text{fine and cosine } BZ - AZ$; $x, y = \text{fine and cos. } \frac{BZ + AZ}{2}$; then (Trig. I. 6. schol. 2.) $qx + py = S.ZB$, $qx - py = S.ZA$, $qy - px = \text{cos. } ZB$, $qy + px = \text{cos. } ZA$. Then in the triangle API, $1 : d :: s : ds = S.$ half AD, and (Trig. I. 2. schol.) $1 - 2dds = \text{cos. } AB$; and in the triangle AZB (Trig. III. 38.) $qqxx - ppyy \times c + qqyy - ppxx = 1 - 2ddss$; but $pp + qq = 1$, and $xx + yy = 1$; therefore $cqqxx - cpp + cppxx + qq - qqxx - ppxx = 1 - 2ddss$, or $cqqxx + cppxx - qqxx - ppxx = 1 - 2ddss + cpp - qq$; that is, $cx - xx = 1 - 2ddss + cpp - 1 + pp = cpp + pp - 2ddss$, whence $x = \frac{c + 1 \cdot pp - 2ddss}{c - 1}$. Then the sides ZA, ZB are known. In the triangle PAI, find the $\angle A$; and in the triangle ZAB, find the angle A, and their difference ZAP; then in the triangle ZAP, there's given two sides and the included angle A; to find ZP, the co-latitude.

P R O B. CLV.

Given two altitudes of the sun or a star, and the times of observation; to find the declination, and latitude.

97. Let Z be the zenith, P the pole; B, A, the places of the sun or star. Let $c = \text{cos. } BZ$, $f =$

$f = \text{cof. AZ}$, $b = \text{cof. ZPB}$, $d = \text{cof. ZPA}$, $x = \text{cof. BP} - \text{ZP}$, $y = \text{cof. BP} + \text{ZP}$. Fig. 97.

Then (Trig. III. 42. cor. 1.) $x - y : 2 :: x - c : 1 - b$, and $x - y : 2 :: x - f : 1 - d$; therefore $x - c : 1 - b :: x - f : 1 - d$; therefore $x - dx - c + cd = x - f - bx + bf$; and $bx - dx = c - f + bf - cd$,

whence $x = \frac{bf - cd + c - f}{b - d}$. Also $1 - b \times x - y = 2x - 2c$, or $by - y = x + bx - 2c$, and $y = \frac{x + bx - 2c}{b - 1} = \frac{2c}{1 - b} - \frac{1 + b}{1 - b} x = \frac{bf - cd + f - c}{b - d}$.

PROB. CLVI.

Given two altitudes of the sun, the difference of times, and difference of azimuths; to find the latitude and declination.

Suppose P the pole, Z the zenith; B, A, the places of the sun. Put $c = \text{cof. BZA}$, $s = \text{cof. BPA}$, $y = \text{S. BP}$ or PA ; $b, d = \text{fine}$ and cof. ZB ; $f, p = \text{fine}$ and cof. ZA . 98.

Then (Trig. III. 38.) in the triangle BZA, $bcf + dp = \text{cof. BA}$; and in the triangle BPA, $yy + \sqrt{1 - yy} \times \sqrt{1 - yy} = \text{cof. BA}$; therefore $yy + 1 - yy = bcf + dp$, and $s - 1 - yy = bcf + dp - 1$, and $y = \frac{\sqrt{1 - bcf - dp}}{1 - s} = \text{S. BP}$ the declination.

Then in the triangle BPA, find the angle B, and in the triangle ZBA find the angle B, and then the difference ZBP. Then in the triangle ZBP there are given two sides ZB, BP, and the included angle B; to find ZP, the co-latitude.

Fig.

P R O B. CLVII.

Given the sun's declination, two altitudes, and the time between them; to find the latitude.

98. Let a, b = fine and cof. PA; d, f = fine and cof. PB; n = cof. ZB, m = cof. ZA; s, c = fine and cof. fine $\frac{1}{2}$ BPA; y, z = fine and cof. $\frac{ZPA + ZPB}{2}$; v, x = fine and cof. ZP.

Then (Trig. I. 6. schol.) $cz - sy = \text{cof. ZPA}$ and $cz + sy = \text{cof. ZPB}$, and (Trig. III. 38.) $aczv - asyv + bx = m$, and $dczv + dsyv + fx = n$; and

$$cz - sy = \frac{m - bx}{av}, \text{ and } cz + sy = \frac{n - fx}{dv}, \text{ and by}$$

adding and subtracting, $2cz = \frac{n - fx}{dv} + \frac{m - bx}{av}$

$$= \frac{an - afx + dm - dbx}{adv}, \text{ also } 2sy = \frac{n - fx}{dv} - \frac{m - bx}{av}$$

$$= \frac{an - afx - dm + dbx}{adv}. \text{ Whence } z =$$

$$\frac{an + dm - afx - dbx}{2cadv}, \text{ and } y = \frac{an - dm + dbx - afx}{2sadv}$$

$$\text{Put } \frac{an + dm}{2cad} = p, \frac{an - dm}{2sad} = q, \frac{db + af}{2cad} = r;$$

$$\frac{db - af}{2sad} = t. \text{ Then } z = \frac{p - rx}{v}, \text{ and } y = \frac{q + tx}{v};$$

But $z = \sqrt{1 - yy}$, and $v = \sqrt{1 - xx}$; then $\sqrt{1 - yy}$

$$= \frac{p - rx}{z}, \text{ and } 1 - yy = \frac{p - rx}{vv}, \text{ and transposing}$$

$$yy = 1 - \frac{p - rx}{vv} = \frac{1 - xx - p - rx}{vv}. \text{ Also } yy =$$

 $q + tx$

$\frac{q+tz}{vv}$. Whence $1-xx-pp+2prx-rrxx=qq+98$.

$2qtx+ttxx$; and reduced

$$\begin{array}{r} ttxx + 2tqx + qq = 0. \\ + rr \quad - 2pr + pp \\ + 1 \quad \quad - 1 \end{array}$$

Or thus,

In the triangle BPA, 2 sides and the included angle are given, to find $\angle B$, and BA. In the triangle BZA, all the sides are given, to find the angle B, and from thence ZBP. Then in the triangle ZPB, two sides and the included $\angle B$, are given; to find ZP the comp. of the latitude.

PROB. CLVIII.

Given three altitudes of the sun in one day, and the times between them; to find the latitude, &c.

Let $s, c = \text{fine and cof. APB}$; $t, b = \text{fine and cof. ABC}$, $d = \text{cof. AZ}$, $f = \text{cof. BZ}$, $g = \text{cof. CZ}$, $x = \text{S.ZPA}$, $y = \text{S.AP}$, BP or CP ; $z = \text{S.ZP}$.

Then (Trig. I. 5. cor. 1.) $c\sqrt{1-xx} - sx = \text{cof. ZPB}$, and $b\sqrt{1-xx} - tx = \text{cof. ZPC}$. And (Trig. III. 38.) $zy\sqrt{1-xx} + \sqrt{1-zz} \times \sqrt{1-yy} = d$,

$czy\sqrt{1-xx} -szyx + \sqrt{1-zz} \times \sqrt{1-yy} = f$, and $byz\sqrt{1-xx} + tyzx + \sqrt{1-zz} \sqrt{1-yy} = g$. By

transposition, $\sqrt{1-zz} \times \sqrt{1-yy} = d - zy\sqrt{1-xx}$. Then $czy\sqrt{1-xx} -szyx + d - zy\sqrt{1-xx} = f$, and

$bzy\sqrt{1-xx} -txyx + d - zy\sqrt{1-xx} = g$. From the former of these two last equations we get $zy =$

$$\frac{d-f}{1-c\sqrt{1-xx}+sx}$$

$$\frac{d-g}{1-b\sqrt{1-xx}+tx}$$

; and making these last equations

tions

Fig.
99

tions equal, and reducing, we have $\frac{x}{\sqrt{1-xx}}$

$$\frac{a-f \times 1-b - d-g \times 1-c}{d-g \times s - d-f \times t} = \text{tang. ZPA. Then}$$

x will be known, and also zy .

But $\sqrt{1-zz} \times \sqrt{1-yy} = d-zy\sqrt{1-xx}$. Put $zy=r$, then $1-yy-zz+zzzy=dd-2dzy\sqrt{1-xx}+zzyy-zzyxx$, and transposing, $yy+zz=1-dd+2dr\sqrt{1-xx}+rrxx=p$ by substitution; then $yy+2yz+zz=p+2r$, and $yy-2yz+zz=p-2r$, and $y+z=\sqrt{p+2r}$, and $y-z=\sqrt{p-2r}$, whence $y = \frac{\sqrt{p+2r} + \sqrt{p-2r}}{2}$, and $z = \frac{\sqrt{p+2r} - \sqrt{p-2r}}{2}$.

P R O B. CLIX.

Having at one instant the altitudes of two known stars; to find the latitude.

100. Let Z be the zenith, P the pole; F, A, the stars. In the triangle APF, there is given the sides AP, FP the co-declinations, and angle P, the diff. right ascensions, to find AF, and $\angle F$. Then in the triangle ZFA, all the sides are given, to find the angle F; then $\angle ZFP$ will be known. Let $c = \text{cof. ZFP}$; $a, b = \text{fine and cof. ZF}$; $d, f = \text{fine and cof. FP}$; $x = \text{cof. ZP}$. Then (Trig. III. 38.) $adc + bf = x = \text{S. latitude}$.

P R O B. CLX.

If there be two known stars in one azimuth, and having the altitude of either given; to find the latitude of the place.

101. Let Z be the zenith, P the pole; F, A, the two stars in the azimuth circle ZFA, and ZF is given.

given. In the triangle APF, two sides and the included angle P, are given; to find the angle F. Put $c = \text{cof. } \angle ZFP$; $a, b = \text{fine and cofine of } ZF$; $d, f = \text{fine and cofine of } FP$; $x = \text{cof. } ZP$. Then in the triangle ZFP (Trig. III. 38.) $adc + bf = x$ the fine of the latitude.

If ZA is given, you must find the angle FAP, and put $a, b = S.$ and $\text{cof. } ZA$; $d, f = S.$ and $\text{cof. } AP$, &c.

Otherwise thus,

Let the altitude of A be given; $s, c = \text{fine and cof. } APF$; $d, f = \text{fine and cof. } AP$; $m, n = \text{fine and cof. } FP$; $a, b = \text{fine and cof. } ZA$, $x = \text{cof. } ZP$. Then (Trig. case 7. spherical triangles) co-

tang. A = $\frac{dn - mfc}{ms}$, and tang. A = $\frac{ms}{dn - mfc} = t$,

by substitution; and (Trig. I. 1. schol.) $\text{cof. } A =$

$\frac{1}{\sqrt{1 + tt}}$. And (Trig. III. 38.) $\frac{ad}{\sqrt{1 + tt}} + bf = x$

the S. latitude.

This is a useful problem.



S E C T. IX.

Geometrical Loci, and Problems relating thereto.

102. **I**F the right line AP be drawn from a given point A, and any number of right lines PM, PM, &c. be drawn thereon, parallel to one another, or making any given angle with AP. And if the relation of the indetermined quantities AP, PM be denoted in general by some equation; and if the lengths of PM be every where, such as that equation gives; then the curve passing through all the points M, is called the *Locus* of the points M, or the locus of that equation. And that equation declares the nature of the curve MM.

The *degrees* of the *Loci* are denominated from the degrees of the equations, by which they are denoted. Thus a locus, of the first degree is that where the indetermined quantities rise to one dimension; of the second degree, when they arise to two dimensions; of the third degree, when they rise to three dimensions, &c.

Right lines are said to be given in position, when they make given angles with one another.

P R O B. CLXI.

103. *If the pole AC or BC revolves about the center C, and the weight D, and string BD hangs at the end of it; to find the nature of the curve GD, described by the weight D.*

Take AG, and CF, =BD. Then since CF, BD are equal and parallel, therefore (Geom. I. 5. cor.

cor. 3.) CB, FD, are equal, or $FD=CB=CA$. Fig. 103.
 And since $FC=GA$, add CG, and then $FG=CA$;
 whence $FD=FG$. Therefore GD is a circle whose
 center is F, the same with AB, but in a lower po-
 sition.

P R O B. CLXII.

Suppose ACD, *acd*, &c. to be right-angled triangles, 104.
 one of whose angles falls upon the fixed point A, the
 other in the line AE; and if the segments BD, *bd*,
 be given; to find the nature of the curve passing
 through all the right angles C, *c*, &c.

Let $AB=x$, $BC=y$, $BD=a$; then in the right-
 angled triangle ACD; $AB(x) : BC(y) :: BC$
 $(y) : BD(a)$; whence $ax=yy$, for the nature of
 the curve. Therefore the curve Cc is a parabola,
 whose latus rectum is BD or *bd*, and A the
 vertex.

P R O B. CLXIII.

A is a fixed point, AB a given line, ABD a given 105.
 angle; then suppose the curve AMB to be generated
 after such a manner that drawing any line AC, it
 may be, as BC to BP :: as *r* to *s*: to find the
 nature of the curve, or the locus of all the points M.

Draw MP, *mp* parallel to DB, and let $AP=x$,
 $PM=y$, $AB=a$. Then by similar triangles AP

$(x) : PM(y) :: AB(a) : BC = \frac{ay}{x}$. And by

the problem, $r : s :: BC \left(\frac{ay}{x}\right) : BP(a-x)$.

Therefore $\frac{say}{x} = ra - rx$, and $y = \frac{r}{sa} \times \frac{a - xx}{r}$, for
 the nature of the curve; and it passes through A;
 because when x is $=0$, y is $=0$.

PROB. CLXIV.

106. Given the triangle ABC, and drawing PD parallel to BC; suppose it be always $PM^2 = PD^2 - BC^2$; to find the nature of the curve BM.

Put $AB = a$, $BC = b$, $AP = x$, $PM = y$, then by similar triangles, $a : b :: x : \frac{bx}{a} = PD$, and by the question, $\frac{bbxx}{aa} - bb = yy$, or $aayy = bbxx - bbaa$, and $y = \frac{b}{a} \sqrt{xx - aa}$. And the curve passes through B.

PROB. CLXV.

107. The triangle ABC is given being right-angled at B, and drawing FM parallel to CB, it be every where $PF^2 + PM^2 = BC^2$; to find the nature of the curve BM.

Let $AB = a$, $BC = b$, $AP = x$, $PM = y$. Then by similar triangles, $a : b :: x : \frac{bx}{a} = PF$. And by the question, $\frac{bbxx}{aa} + yy = bb$, whence $yy = bb - \frac{bbxx}{aa}$, and $y = \frac{b}{a} \sqrt{aa - xx}$, the equation for all the points M. And in A, where x is 0, $y = b$, or $AD = CB$.

P R O B. CLXVI.

If AB, CF, AC be right lines given in position; 108.
and PD, pd be always parallel to AC; and if
PM be every where equal to CD; to find the
locus of the point M.

Since AC, PD are parallel; AP will be to CD
in a given ratio (Geom. II. 12. cor. 2.); put
 $AP=x$, $PM=y$, and let $a:b::AP(x):$
 $\frac{bx}{a}=CD$; therefore $y=\frac{bx}{a}$. Therefore AMm is
a right line, passing through A.

P R O B. CLXVII.

If the three lines CA, CB, AB be given in position; 109.
and PM be always drawn parallel to AB; and it
be every where $AP \times PD = PM^2$; to find the nature
of the curve Mm.

Let $CA=a$, $AB=b$, $AP=x$, $PM=y$; then by
similar triangles, $a:b::a+x:\frac{a+x}{a}b=PD$, then
by the question, $\frac{a+x}{a}bx = yy$, and $y =$

$\sqrt{\frac{b}{a} \times ax + xx}$; and the curve passes through A,
since both x and y are 0, at once.

But if CB be parallel to CA, then $PD=AB$,
and $bx=yy$, or $y=\sqrt{bx}$. And if C lie on the o-
ther side of A, then $PD = \frac{a-x}{a}b$, and $y =$

$\sqrt{\frac{b}{a} \times ax - xx}$. In which case, when $x=a$,
then $y=0$, and the curve passes there through
the axis CA.

Fig. 109. When x is greater than a , y is the square root of a negative quantity, which is impossible; and therefore the curve goes no further than A.

PROB. CLXVIII.

110. There is given the right-angled triangle ABD; and drawing PM, pm always parallel to BD; and making PM every where equal to BF; to find the nature of the curve DMM.

Put $AB=a$, $BD=b$, $BP=x$, $PM=y$. Then by similar triangles, $a : b :: a+x : \frac{a+x}{a} b$


$$= PF, \text{ and } BF^2 = xx = \frac{bb \times a+x}{aa} = \frac{aaax + bbaa + 2bbax + bbxx}{aa}; \text{ therefore}$$


$$y = \sqrt{bb + \frac{2bbx}{a} + \frac{aa+bb}{aa}xx}, \text{ for the nature of the curve.}$$

PROB. CLXIX.

111. BA is a given line, draw PM perpendicular to BAP; and let AM be always a mean proportional between AB and AP; to find the nature of the curve AmM.

$$\text{Let } AB=a, AP=x, PM=y; \text{ then } AM = \sqrt{ax+yy}, \text{ and per quest. } a : \sqrt{ax+yy} :: \sqrt{ax+yy} : x; \text{ and } ax = xx+yy, \text{ whence } y = \sqrt{ax-xx}.$$





The right page of the manuscript is mostly blank, with faint horizontal lines visible, suggesting it was part of a ruled notebook or a page intended for text. There is no legible text or other markings on this page.

P R O B. CLXX.

The line CF and point A being given; from any 112.
point D in that line, through A, draw DAM,
and make $DA \times AM$ always equal to a given
square; to find the locus of M.

Draw BAP perpendicular to CF, and PM per-
pendicular to AP, and put $AB = a$, $AP = x$,
 $PM = y$, then $AM = \sqrt{xx + yy}$, bb = the given
square. By similar triangles $x : \sqrt{xx + yy} :: a :$
 $\frac{a\sqrt{xx + yy}}{x} = AD$. Then per quest. $\frac{a}{x} \times \sqrt{xx + yy}$
 $= bb$, and $xx + yy = \frac{bbx}{a}$, whence $y = \sqrt{\frac{bbx}{a} - xx}$.

P R O B. CLXXI.

AD is a circle, C its center; draw AB perpendicu- 113.
lar to CAP. Then draw any line CB, and BM
parallel to AP; and make always $BM = DB$; to
find the nature of the curve mM.

Draw MP perpendicular to CP, and let $CA = r$,
 $AP = x$, $PM = y$; then $CB = \sqrt{rr + yy}$, and
 $BD = \sqrt{rr + yy} - r$. But AP or $BM = BD$, that
is, $x = \sqrt{rr + yy} - r$, and $\sqrt{rr + yy} = r + x$, and
squaring, $rr + yy = rr + 2rx + xx$, whence $y =$
 $\sqrt{2rx + xx}$.

PROB. CLXXII.

114. CAD is a given angle, CD a given line, M a given point in it. Let this line so move in the angle CAD, that the ends D, C may always touch the sides AD, AC; to find the curve described by the point M.

Draw MP, MB parallel to AC, AD, and put $DM = a$, $CM = b$, $\text{cof. } \angle A = c$, $AP = x$, $PM = y$. By similar triangles, $CM (b) : BM (x)$

$$:: DM (a) : DP = \frac{ax}{b}. \text{ But in the triangle}$$

$$\text{MPD (Trig. case 5.) } yy + \frac{aaxx}{bb} - \frac{2acxy}{b} = aa,$$

or $yy = aa + \frac{2acxy}{b} - \frac{aaxx}{bb}$, for the equation of the curve.

PROB. CLXXIII.

115. The line CA is perpendicular to AP, ABM is a square whose sides CB, BM are given; to find the curve described by the point M, whilst the $\angle B$, and end C, move along PA, AC.

Draw MP perpendicular to AP, $CB = a$, $BM = b$, $AP = x$, $PM = y$; the triangles CAB, BMP are similar, and $b : y :: a : \frac{ay}{b} = AB$. And $BP =$

$$\sqrt{bb - yy}, \text{ therefore } \frac{ay}{b} + \sqrt{bb - yy} = x, \text{ and}$$

$$\sqrt{bb - yy} = x - \frac{ay}{b}, \text{ and squaring } bb - yy = xx -$$

$$\frac{2axy}{b} + \frac{aayy}{bb}. \text{ And reduced } \frac{aa + bb}{bb} yy = bb$$

$$+ \frac{2ayx}{b} - xx.$$

PROB.

P R O B. CLXXIV.

The lines AV, AB, AE are given in position, the point V is given; if the line VE be always drawn through V, and the part intercepted CE be divided in a given ratio at M; to find the locus of the points M, m. 116.

Draw MP parallel to AB, and DM, BE parallel to AP, and put $AP=x$, $PM=y$, $AV=a$. r to s as CM to ME, $S.\angle BAE=p$, $S.\angle EAP$ or $BEA=q$.

The triangles VAC, VPM, DCM, BCE are similar, and $a+x : y :: a : \frac{ay}{a+x} = AC$, and $a+x$

$: y :: (DM) x : \frac{xy}{a+x} = DC$. Also $r : s :: CM$

$: ME :: CD \left(\frac{xy}{a+x} \right) : DB = \frac{sxy}{ra+rx}$. And CM

$: CE :: r : r+s :: x : \frac{r+s}{r}x = BE$, and in the tri-

angle BAE, $p : q :: BE \left(\frac{r+s}{r}x \right) : BA =$

$\frac{r+s}{pr}qx = y + \frac{sxy}{r \times a+x}$, and $r+s \times qx \times a+x$

$= pry \times a+x + psxy$, which reduced is (putting

$r+s=d$) $dqxx + dqax = pray + dpxy$, and $y =$

$\frac{dqxx + dqax}{pra + dpx} = \frac{dq}{p} \times \frac{ax+xx}{ar+dx}$.

P R O B. CLXXV.

BCD is a given angle, D a fixed point, BM parallel to CP; and BM to MD are always in a given ratio; to find the locus of M. 117.

Draw MP parallel to BC; and put $CA=a$, $AD=b$, $\text{cof. } \angle P=c$, $AP=x$, $PM=y$; then $PD=$

Fig. $PD=b-x$, and $BM=a+x$; then (Trig. case 5.)

117. $MD = \sqrt{yy + b-x^2 - 2cy \times b-x}$. Whence it

is, $a : b :: a+x : \sqrt{yy + b-x^2 - 2cy \times b-x}$;
 which squared and multiplied, $bbaa + 2bbax + bbxx$
 $= aayy + aabb - 2aabx + aaxx - 2caaby + 2caayx$, and
 reduced $aayy + 2caayx - 2caaby = bbxx + aaxx +$
 $2bbax + 2aabx$.

PROB. CLXXVI.

118. *C, D are two fixed points in the line CD; and CM square is every where to MD square, in the given ratio of r to s. To find the locus of M.*

Draw MP perpendicular to CD, and let $CA=a$,
 $AD=d$, $AP=x$, $PM=y$; then $CP=a+x$,
 $PD=b-x$. Therefore it is $r : s :: aa : bb ::$
 $(CM^2) yy + a+x^2 : (MD^2) yy + b-x^2$; there-
 fore $bbyy + bbaa + 2bbax + bbxx = aayy + aabb -$
 $2aabx + aaxx$. Whence $aayy - bbxx = bbxx - aaxx$
 $+ 2bbax + 2aabx$; and $yy = \frac{a+b}{aa-bb} \times 2abx -$
 $xx = \frac{2ab}{a-b} x - xx$.

PROB. CLXXVII.

119. *The lines AB, AD are given by position; the points P, B, and angles CPD, CBD are given. Now if the angles CPD, CBD move about the centers P, B, whilst the intersection D (of the sides PD, BD) runs along the line AD; to find the curve which the intersection C, of the other sides, describes.*

Draw CS, DF perpendicular to AB; and put
 $AP=a$, $Pb=b$, $\text{tang. } \angle PAD=t$, $\text{tang. } \angle CPD=p$,
 $\text{tang. } \angle CBD=q$.

tang. CBD = q , PF = v , PS = x , SC = y , and Fig. 119.
 BF = $b - v$, BS = $b - x$.

Then by Trigonometry, $1 : a + v :: t : ta + tv$
 = DF.

And $v : 1 :: ta + tv : \frac{ta + tv}{v}$ = tang. DPF.

Also $x : 1 :: y : \frac{y}{x}$ = tang. CPS.

And $b - v : 1 :: ta + tv : \frac{ta + tv}{b - v}$ = tang. DBF.

And $b - x : 1 :: y : \frac{y}{b - x}$ = tang. CBS.

But (Trig. I. 8.) $1 - \frac{ta + tv}{vx} y : 1 :: \frac{ta + tv}{v} + \frac{y}{x} : p$.

And $1 - \frac{ta + tv}{b - v} \times \frac{y}{b - x} : 1 :: \frac{ta + tv}{b - v} + \frac{y}{b - x} : q$.

and multiplying the extremes and means,

$$px - ptay - ptvy = tax + tvx + vy.$$

And $q \times b - v \times b - x - taqy - tqvy = ta + tv \times b - x + by - vy$; that is, $qb \times b - x - qv \times b - x - taqy - tqvy = tab - tax + tv \times b - x + by - vy$; and transposing, $tv + qv \times b - x + tqvy - vy = qb \times b - x - taqy + tax - tab - by$. By this and the former equation, $v = \frac{qbb - qbx + taqy - tab + tax - by}{qb - qx + tqy + tb - tx - y} =$

$\frac{tax + ptay}{px - tx - ptv - y}$. And substituting for the known compound quantities,

$$\frac{cx + dy + f}{-gx + hy + l} = \frac{tax + tapy}{nx - sy}; \text{ reduced}$$

$$\begin{array}{r} tapby - tagxx - tapgy + talx + tapy = 0. \\ +sd - cn + tab - fu + sf \\ +sc \\ -dn \end{array}$$

Fig.

P R O B. CLXXVIII.

120. To find the figure for the section of a groin, the bases of the two solids being AFL a semicircle, and ABC, a right-angled isocetes triangle.

Groining in joinery is fitting two prismatic solids, to join at right angles, so that the surfaces of both may coincide, no part of one being higher than the other, and the ends of both of them must be cut away to a certain figure, or else they can never join truly.

Let the perpendicular sections AFL, ABC of the solids be perpendicular to the plane LACD, on which the figure is to be drawn. And suppose AMD to be the figure; draw MI, MP parallel to AC, AL; at I, P, draw the ordinates IF, PO, perpendicular to AL, AC. Now the nature of the groin requires that the lines FI, and PO, which are to coincide, must be equal. Therefore compute FI, OP in both figures, and put them equal to one another.

Let AL or AC = a , AP = x , PM = y . Then $IF = \sqrt{AI \times IL} = \sqrt{a-y} \times y$; and since ABC is a right angle and AB = BC, OP will = AP; therefore OP = x , whence $x = \sqrt{ay - yy}$. Whence AM is an arch of a circle equal to AF. And for the same reason, the part at D of AMD is a like arch, and the whole curve AMD consists of two quadrants of the circle AFL, meeting in the middle, and turning contrary ways. Therefore if the ends of the two solids, be cut into the figures ELDMAF, and BCDMAB; they will exactly fit one another.

P R O B.

P R O B. CLXXIX.

To find the figure of a groin, when the bases or ends 121.
of the bodies are AFL a semicircle, and ABC the
segment of a circle.

Let AMD be the curve; draw MP, MI parallel to AL, AC; and IF, PO perpendicular to AL, AC.

Put $AL = a$, $AC = b$, $AP = x$, $PM = y$. Then because the figures APL and ABC must always be of equal height, therefore $\frac{1}{2}a =$ the height of ABC. Then to find the diameter of ABC, we shall have $\frac{1}{2}bb + \frac{1}{2}aa$ divided by $\frac{1}{2}a$, for the diameter; put $D =$ diameter, then $D - a =$ the distance of the cord AC from the center, put $D - a = c$, and PO or $IF = v$. Then by the nature of the circle (in the figure ABC), $2cv + vv = bx - xx$; and $ay - yy = vv$, in the figure AFL. Therefore $2cv + ay - yy = bx - xx$, and $2cv = yy + bx - xx - ay$, also $v = \sqrt{ay - yy}$, and $2cv = 2c\sqrt{ay - yy} = yy - ay + bx - xx$; which squared and reduced gives an equation of the fourth power for the locus of M.

P R O B. CLXXX.

To find a general equation to the ellipsis, referred to any line as an axis; which ellipsis will therefore be the locus of that equation.

Let BFDG be the ellipsis, C the center: Let 122.
the point A be given, and any line AL, given in position, for the axis. Take the angle KAL at pleasure, and through C, draw the diameter BD, parallel to AK, and FCG the conjugate to it, and AN, PM, LK parallel to FG. Put BC or $CD = t$,

Fig. CD=t, p= parameter belonging to BD, AL=a,
 122. LK=b, AK=c, CN=f, AN=n, AP=x,
 PM=y.

By the similar triangles ALK, API; $a : b :: x$
 $:\frac{bx}{a}=PI$; and $a : c :: x : \frac{cx}{a}=AI$. Then PR

$$= n + \frac{bx}{a}, RM=y-n-\frac{bx}{a}, CR=\frac{cx}{a}-f, BR$$

$=t-f+\frac{cx}{a}$, RD=t+f- $\frac{cx}{a}$. And by the pro-
 perty of the ellipsis $2t : p :: BRD : RM^2 :: t-$

$$ff + \frac{2cfx}{a} - \frac{ccxx}{aa} : yy-2ny + nn + \frac{bbxx}{aa} -$$

$\frac{2bxy}{a} + \frac{2bnx}{a}$; and multiplying extremes and
 means, and reducing,

$$2aatyy-4abixy+2tbbxx-4aatny+4abtnx+2aatnn=0$$

$$+pcc \qquad -2apcf-paatt$$

$$\qquad \qquad \qquad +paaff$$

An equation to the ellipsis FD referred to the axis
 AL. Where note, yy and xx have the same sign.
 And if xy is in the equation, the square of half its
 coefficient is less than the coefficient of xx multi-
 plied by the coefficient of yy. And if xy be want-
 ing, xx and yy have the same sign.

PROB. CLXXXI.

To find a general equation to the hyperbola, referred
 to any line as an axis; and which hyperbola will
 consequently be the locus of that equation.

123. Let DM be a hyperbola, C the center, AL
 any line drawn from the given point A. Make
 LAK any given angle; and through C draw the
 diameter BD, parallel to AK, and FCG its con-
 jugate,

jugate, and draw AN, PM, LK parallel to FG, Fig. Put BC or CD = t , p = parameter of BD, AL = a , LK = b , AK = c , CN = f , AN = n , AP = x , PM = y .

From the similar triangles ALK, API, we shall get (as in the last problem,) $PI = \frac{bx}{a}$, $AI = \frac{cx}{a}$, whence $PR = n + \frac{bx}{a}$, $RM = y - n - \frac{bx}{a}$, and $CR = \frac{cx}{a} - f$, and $DR = \frac{cx}{a} - f - t$, and $BR = \frac{cx}{a} - f + t$. And by the nature of the hyperbola,

$$2t : p :: (BR \times DR) \frac{ccxx}{aa} - \frac{2cfx}{a} + ff - tt :$$

$$(MR^2) yy - 2ny + nn - \frac{2bxy}{a} + \frac{2bnx}{a} + \frac{bbxx}{aa}.$$

And the means and extremes multiplied, and then reduced,

$$2taayy - 4btaxy + 2tbbxx - 4tnaay + 4tnbax + 2taann = 0.$$

$$- pcc \qquad + 2acpf \quad - paaff$$

$$\qquad \qquad \qquad + paatt$$

Note, when xy is not in the equation, yy and xx have different signs. And if xy be there, the square of half its coefficient is greater than the coefficient of xx multiplied by the coefficient of yy .

S E C T. X.

Mechanical Problems.

P R O B. CLXXXII.


124. *If the weight P break the beam DE, when supported loose at A, B; to find what weight will break it, when the ends D, E, are fixed, that they cannot rise.*


SUPPOSE $DA=AC$, and $BE=BC$. Suppose the beam cut through at C, and let $\frac{1}{2}P$ be laid upon D, whilst $\frac{1}{2}P$ remains at C; then the pressure at A will be $=P$, therefore the beam will also break at A, having the same stress there as it had at C. For the same reason, if $\frac{1}{2}P$ be applied to E, CE will break at B. Consequently, if $2P$ be applied to C, the beam being whole; and the ends D, E fixed; the beam will break at A, C, and B; and therefore it bears twice the weight or $2P$, at C, before it breaks.

P R O B. CLXXXIII.

125. *The strength of a beam AB, being given; to find its strength when a hole (ac) is cut out of the middle, and also an equal one (rn) in the side.*

By the principles of mechanics, the strength of the beams whose thicknesses are db , da , dc , will be as db^2 , da^2 , and dc^2 . Now as the strength of all the particles between b and d , is denoted by db^2 , and the strength of all the particles between a and





17439. X. P.

SECTION II
Geometrical Problems

PROBLEM I

1. To find the locus of the intersection of two straight lines, one of which is fixed, and the other moves so that its intercepts on two fixed lines are reciprocals.

Let the fixed line be AB , and the two fixed lines be AC and AD . Let the moving line be EF , where E is on AC and F is on AD . Let P be the intersection of EF and AB . The condition is that $AE \cdot AF = k$, where k is a constant. The locus of P is a hyperbola with AB as its axis.

PROBLEM II

2. To find the locus of the intersection of two straight lines, one of which is fixed, and the other moves so that its intercepts on two fixed lines are in a constant ratio.

Let the fixed line be AB , and the two fixed lines be AC and AD . Let the moving line be EF , where E is on AC and F is on AD . Let P be the intersection of EF and AB . The condition is that $AE/AF = k$, where k is a constant. The locus of P is a straight line parallel to AB .

and d , by ad^2 ; therefore the strength of all the particles between b and a , (the point D being fixed) will be $db^2 - da^2$, add the strength between c and d , which is cd^2 ; and the strength of ba and cd , that is, the strength of the hollow beam is $db^2 - da^2 + cd^2$. But at the section r the strength is fn^2 .

Whence if $nr = ac$, the strength at b to the strength at r is as $db^2 - da^2 + cd^2$ to $db - ca^2$; that is, as $db^2 - 2dc \times ca - ca^2$ to $db^2 - 2db \times ca + ca^2$. Therefore if db^2 be the strength of the whole beam, $2dc + ca \times ca$ will be the defect of strength of the hollow beam, when it breaks at b ; and $2ab - ca \times ca$, the defect of strength when it breaks at n or f , which is greater than the former. And for the same reason the defect of strength to break at d , will be $2ba + ac \times ca$.

P R O B. CLXXXIV.

To support a long prismatic body horizontally by two props A, B ; that it shall as soon break in A or B as in C .

Let $DA = AF = GB = BE = y$, $CF = CG = x$, $DC = CE = n$, then $n = 2y + x$.

The parts AF and BG lay no stress upon C , being balanced by the contrary weights DA, BE , equal to them. Therefore the stress at C , arises from the weight FG ; and must be equal to the stress at A , arising from the weights AD, AF .

The stress at A by the weight DF is $\frac{1}{2}DF \times DF$ or $2yy$, (Mechan. 70. and cor.) and the stress (by FG suspended) at C is $AB \times FG$, or $2y + 2x \times 2x$. But (ib. cor. 5.) $2AC(2y + 2x) : AF + AC(2y + x) ::$ stress at C , by FG suspended at $C(2y + 2x \times 2x) : \text{the stress at } C, \text{ in the position } FG = \frac{2y + x \times 2x}{2y + x}$.

H h

Therefore

Fig. Therefore $\frac{2xy}{2y+x} = \frac{2x}{2y+x} \times 2x$. Or $yy = 2yx + xx =$
 126. $nx = n \times n - 2y$, and $yy + 2xy = nm$. Whence $y =$
 $n \times \sqrt{2-1}$. And $x = n \times 3 - 2\sqrt{2}$.

PROB. CLXXXV.

127. If two weights P, T keep one another in equilibrio, on the two wheels whose radii are AB, CB; the strait tooth AB of the one, acting on the crooked tooth BD of the other; to find the proportion of the weights P, T.

Draw EBF perpendicular to OD, EH perpendicular to AB, and FG perpendicular to BC. The point B of the end AB, is acted upon by three forces: 1. in direction AB; 2. in direction BE; 3. in direction EH by the weight P; and these forces are as BH, BE, EH.

Again, the point B of the tooth BD is acted on by these three forces: 1. in direction BC; 2. in direction FB; 3. in direction FG by the weight T, and these forces are as BG, BF and FG. But the action and reaction at the point B, being equal; we have $BE = BF$, and in the right-angled triangle BHE, $\text{rad. } (1) : EB :: S.ABE : HE = EB \times S.ABE$. And in the triangle BGF, $\text{rad. } (1) : BF \text{ or } EB :: S.FBG : GF = EB \times S.FBG$. Whence $P : T :: HE : GF :: EB \times S.ABE : EB \times S.FBG$; that is, $P : T :: \text{cof. ABD} : \text{cof. CBD}$, when the weights are in equilibrio.

Whence if ABC is a right line, $P = T$; and if $\angle CBD = 0$, then $P : T :: \text{cof. ABO} : \text{radius}$.

PROB. CLXXXVI.

To find proper numbers for the wheels and pinions of a clock, to go eight days; and to shew hours by the great wheel, minutes by the second wheel, seconds by the ballance wheel, and to beat seconds.

For the moving part.

Suppose four wheels in the moving part A, B, 128. C, D, and let the numbers for the wheels and pinions be denoted as in the figure, and let $f=12$, b =height the weight descends, t =time of going down in hours.

It is plain $\frac{A}{p}$ = number of revolutions B has for one of A, and $\frac{B}{q}$ = number of revolutions C has for one of B; whence $\frac{AB}{pq}$ = number of revolutions C has for one of A. And likewise $\frac{ABC}{pqr}$ = number of revolutions D has for one of A.

Since the arbor of D carries an index to shew seconds, therefore $D=30$, because for every tooth there are two beats, and $2D=60$.

Because the arbor of B carries an index to shew minutes, and of A to shew hours; consequently A goes about in 12 hours, and B in 1, whence $\frac{A}{p} = 12$.

And because D goes 60 times round for B's once, therefore $\frac{BC}{qr} = 60$.

Therefore the two equations $\frac{A}{p} = 12$, and $\frac{BC}{qr} = 60$, will resolve the question; which being

Fig. unlimited, many of them may be taken at pleasure, provided they be all whole numbers.

Suppose $r=6$, $p=8$, $q=8$; then $A=96$, and $BC=6 \times 8 \times 60$, and if $B=60$, then $C=48$; or $B=72$, and $C=40$. It will be better if B and q , C and r be prime to one another.

To find the diameter of the wheel for the rope, it will be $t : b :: f : \frac{fb}{t} = \text{circumference}$, and

$$\frac{fb}{3.1416t} = \text{diameter.}$$

For the striking part.

Let L be the fly, K the warning-wheel, I the detent wheel, H the pin-wheel, G the great wheel, F the count-wheel, their teeth and pinions as in the figure; n = number of striking pins, and there are 78 strokes in 12 hours: F goes round in 12 hours, I goes round for every stroke of the clock.

Now $\frac{78}{n} = \text{number of revolutions of } H \text{ in } 12$ hours, and $\frac{FG}{ab} = \text{number of revolutions of } H$ to one of F , that is, in 12 hours; therefore $\frac{FG}{ab} = \frac{78}{n}$.

Again, I goes round n times for H 's once, and therefore $\frac{H}{c} = n$. Therefore from these two equations $\frac{FG}{ab} = \frac{78}{n}$, and $H = cn$, all the requisites may be found; but being unlimited most of the numbers may be taken at pleasure, so as they be all convenient whole numbers.

Because the pin in the warning-wheel must always come to the same place when the clock has struck

struck out, therefore $\frac{I}{d} =$ a whole number. L Fig. 128.
 and e may be any numbers, because there is no phenomenon to be shewn by them.

The train, or beats in an hour is $= \frac{ABC}{pqr} \times \frac{60}{12}$
 $= \frac{5ABC}{pqr}$. Suppose $n=12$, $a=6$, $b=8$, $c=6$;

then $H=72$, and $FG = \frac{78 \times 48}{12} = 4 \times 78$, there-

fore F may be $=13$ and $G=24$. But note $\frac{FG}{ab}$
 may be put into one wheel or more as one pleases.

If the string go about the axis of F , its diameter is found as in the other. But if it go round the axis of G , it must be made less in proportion as a to F . If one weight carry both parts, their diameters must be but half the former quantities.

P R O B. CLXXXVII.

Supposing with BORELLI (part. I. prop. 22. de motu animalium), that a strong man can but bear 26 lb. at arm's end, and that the weight of his whole arm is equivalent to 4 lb. at arm's end; from the length of his arm given; to find the dimensions of that man's arm, that can bear no more than its own weight.

Suppose 4 lb. at arm's end equivalent to 8, the weight of the arm. And suppose the two arms, similar solids, and the arm $=$ half the length of the body. Put $a =$ length of a common man's arm, $b=4$ lb. $w=26$ lb, $x =$ length of the great man's arm.

The weight of like bodies are as the cubes of the sides, $a^3 : x^3 :: 2b : \frac{2bx^3}{a^3} =$ weight of the

H h 3

great

Fig.
128.

great man's arm, and $\frac{bx^3}{a^3}$ = the weight at arm's end, producing the same stress.

And the stress being as the length and weight, we have $\overline{w+b} \times a$ = stress of the common man's arm; and $\frac{bx^3}{a^3} x$ = stress of the great man's arm.

But (by mechanics) the stress in this case, is as the strength, that is, as the cubes of the like sides.

Therefore $a^3 : x^3 :: \overline{w+b} \times a : \frac{bx^4}{a^3}$, whence bx^4

$$= \overline{w+b} \times ax^3, \text{ or } x = \frac{w+b}{b} a = \frac{30a}{4} = 7\frac{1}{2}a.$$

Now if $a=1$ yard; then if there be a man whose height is above 15 yards; he will not be able to stretch out his arm.

P R O B. CLXXXVIII.

129. Given the length and position of the beam AD, leaning against the wall DE; to find the position of the plane BE, on which it may stand without moving.

Let G be the center of gravity of the beam together with any weight it carries. Through G, draw the horizontal line BH. And suppose DA put into the position da , infinitely near the former. Now since the beam is to have no inclination of moving from the position DA, or da ; the center of gravity G, g must be in the horizontal line BH, by the principles of mechanics. Draw Gn , dm , Ar perpendicular to ad or AD. And let $DG=b$, $AG=c$, $b=S.DHG$, p, q =sine and cof. ADH; s, f =sine and cosine DGH, x =tang. DAE, $v=DF$.

Since $DG=dg=mn$, and $AG=ag=rn$, therefore $Dm=ng=ar$. In the triangle Ddm, $S.mdD$ (q)

(q) : S.mDd (p) :: mD : md = $\frac{p}{q} \times mD$ or $\frac{p}{q}$ Fig. 129.

$\times gn$. And in the triangle Ggn, S.gGn (f) :

S.Ggn (s) :: gn : Gn = $\frac{s}{f} \times gn$. By the similar triangles Fdm, FGn, Fd (v) : FG

(b-v) :: md $\left(\frac{p}{q} \times gn \right)$: n G

$\left(\frac{s}{f} \times gn \right)$, whence $\frac{sv}{f} = \frac{pb-pv}{q}$, and $qsv = pbf - pfv$, and $v = \frac{pbf}{pf+qs}$. But (Trigon. I. 5.)

$pf+qs=b$. Therefore $v = \frac{pbf}{b}$. In the triangle

Aar, 1 : ar or ng :: x : rA = x \times ng. And in the similar triangles FDM, FAR, Fd (v) : md

$\left(\frac{p}{q} \times gn \right)$:: FA (b+c-v) : rA (x \times ng) ;

therefore $vx = \frac{p}{q} \times \overline{b+c-v}$, and $vqx = pb + pc - pv$, and $vqx + pv = pb + pc$, and substituting the

value of v, $\frac{pbf}{qx+p} \times \frac{pbf}{b} = pb + pc$, and $bfqx +$

$bfp = bb + cb$, whence $x = \frac{bb+cb-bfp}{bfq} = \frac{b+c}{bfq} b$

$-\frac{p}{q}$. Whence

1. If DH be perpendicular to the horizon, $b=1$,

$$s=q, f=p, \text{ and } x = \frac{b+c}{bpq} - \frac{p}{q}.$$

2. If DH nearly coincides with DA, $b=s, p=0$,

$$q=1, \text{ then } x = \frac{b+c}{b} \times \frac{s}{f}, \text{ or } \frac{b+c}{b} \times \text{tang.}$$

DGH.

PROB. CLXXXIX.

Having given, the specific gravity of two things, and likewise the specific gravity of a mixture of them; to find the proportion of the things mixed.

Let A, B be the two things, and M the mixture, a, b, c the specific gravity of A, B, M; A, B, M their magnitudes. Then since the absolute weight is as the magnitude and specific gravity; therefore aA, bB, mM will be the weight of A, B, M. And $aA + bB = mM = m \times \frac{A+B}{m}$, and transposing $aA - mA = mB - bB$. Whence $\frac{a-m}{a-m} : A : B$.

PROB. CXC.

Having given the weights and velocities of two spherical bodies perfectly elastic, meeting one another in a right line; to determine their velocities, after reflexion.

Let A, B, be the weights of the bodies, a, b , their velocities towards different parts, x and y their velocities the contrary way, after reflexion. Then Aa, Bb are the quantities of motion in their respective directions, before reflexion; and Ax, By after. As the bodies are elastic, they will recede from one another, with the same relative velocity they met, whence $a + b = y + x$. And (by mechanics) the difference of the motions, moving the same way, will remain the same after as before the stroke, therefore $Aa - Bb = B_1 - Ax$, but $y = a + b - x$, therefore $Aa - Bb = Ba + Bb - Bx - Ax$; and transposing, $Ax + Bx = aB + bB - Aa + bB$, and $x = \frac{B - A \times a + 2Bb}{A + B}$, and $y = \frac{A - B \times b + 2Aa}{A + B}$.

PROB.

P R O B. CXCI.

ACDB is a thread fixed at A and B, at the points C, D of this thread are fixed the two threads CE, DF, with the weight EF; having given the weight F, and the position of the points C, D; to find the weight E.

Let the weight $F = w$, weight $E = x$, $S. \angle CDB = 130^\circ$, $= s$, $S. FDB = t$, $S. DCA = p$, $S. ECA = q$.

The point D is kept in equilibrio by 3 forces in directions DB, DC, DF, which are to one another, as the sines of the angles they pass through (Mechan. 8. cor. 2.) : therefore $S. CDB (s) : \text{force at F } (w) :: S. FDB (t) : \text{force in DC} = \frac{tw}{s} = \text{force in CD}$, because action and reaction are equal and contrary.

Again, the point C is drawn by three forces, in directions CD, CA, CE; therefore, $S. ECA (q) : \text{force CD } \left(\frac{tw}{s}\right) :: S. ACD (p) : \text{force at E } (x)$,

therefore $qx = \frac{ptw}{s}$, and $x = \frac{pt}{qs} w$.

P R O B. CXCII.

Three points of the ceiling, A, B, C are given, to which are fixed the threads AF, BF, CF whose lengths are given; to these is fixed the thread FD, with the given weight D; to find the tension of all the threads.

Because the triangle ABC is given, and the lengths of the threads; the point O will be given, where DF produced cuts the ceiling. Produce AO to E, and draw EF, which will be $= \sqrt{FO^2 + OE^2}$.

All

Fig. 131. All the sides of the triangles CFE, EFB, are given, and consequently the angles. Now instead of the threads FC, FB, suppose the thread FE to sustain the weight. And then the whole is sustained by the two threads AF, FE acting in the perpendicular plane AOEF. Draw OL parallel to AF, in the plain AEF, and LG parallel to CF in the plane CFB.

Put $AE = a$, $AF = b$, $EO = c$, $AO = d$, $EF = f$, $OF = b$, $S.\angle CFB = p$, $S.CFE = q$, $S.EFB = s$.

Then (Mechan. 8.) the tension of the threads DF, AF, EF, will be denoted by OF, OL, LF; and taking away the thread FE, the tension of the threads CF, BF, will be LG, GF. Then to find each. By similar triangles, EA (a) : AF (b) ::

$$EO (c) : OL = \frac{bc}{a}. \text{ And } EA (a) : AO (d) ::$$

$$EF (f) : LF = \frac{fd}{a}. \text{ And in the triangle FLG,}$$

$$S.LGF (p) : LF \left(\frac{fd}{a} \right) :: S.LFG (s) : LG =$$

$$\frac{sf d}{ap}, \text{ and } :: S.FLG (q) : FG = \frac{qdf}{pa}.$$

Therefore the tensions of DF, AF, CF, BF, are respectively as b , $\frac{bc}{a}$, $\frac{sf d}{ap}$, $\frac{qfd}{ap}$.

S E C T. XI.

Philosophical or physical Problems.

P R O B. CXCIII.

Required the height of the tower, from the top of which a stone falling to the bottom, the sound will reach the ear at the top, in the time of the fall.

PUT $b = 16\frac{1}{2}$ feet, the height a body falls in a second.

$c = 1142$ feet, the space sound moves through in a second.

$a =$ time of the body's falling.

Then $1 : c :: a : ca =$ space sound moves in the time a .

And $1 : a :: b : baa =$ height the ball descends.

Therefore per qu. $baa = ca$, and $a = \frac{c}{b} = 71$ seconds.

And $baa = ca = \frac{cc}{b} = 81088$, the height.

P R O B. CXCIV.

There is a round tower, whose circumference is 100 yards, a spiral tube runs about, from bottom to top, at an elevation of $61^{\circ} : 5'$. A ball put in at the top of this tube will run down to the bottom in 8 seconds; to find the height.

Let $\angle ABD$ be $61^{\circ} : 5'$, AC perpendicular to BD , and BC perpendicular to AB . Then whilst a body falls through AC , another would descend through

Fig. through AB in the same time (Mechan. 34. cor. 1.)
 132. Put $b=16.1$ feet, $d=8''$, $s=S.ABD$. Then by
 the laws of falling bodies, $1 : b :: dd : bdd =$
 height fallen in $8'' = AC$. And rad. (1) : AC
 (bdd) :: S.C (s) : $bdds = AB$. And rad. (1) :
 AB ($bdds$) :: S.ABD (s) : $bddss = AD$, the height
 required $= 789$.

P R O B. CXC.V.

*Given the distance of the earth and the moon, and
 their quantities of matter; to find the place where
 a body will be attracted to neither of them.*

Let d = distance of their centers, l = matter in
 the moon, t = matter in the earth, x = distance
 from the earth where the body is, then $d-x$ = its
 distance from the moon.

Then since the force of attraction is as the mat-
 ter directly, and the square of the distance inver-
 sely; therefore we have $\frac{t}{xx} =$ earth's attraction, and

$\frac{l}{d-x^2} =$ moon's attraction; but *per quest.* these

are equal, therefore $\frac{t}{xx} = \frac{l}{dd-2dx+xx}$; which
 reduced is $t-l.xx-2dtx+ddt=0$.

P R O B. CXC.VI.

*A clock that keeps true time on the surface of the
 earth; being carried to the top of a certain moun-
 tain, lost 2 minutes in a day. What was the
 mountain's height?*

Let r = earth's radius = 6982000 yards, $b=1440$
 minutes, $c=2$ minutes, a = height of the mountain.
 But

But (Mechan. 40. cor. 6.) the length of a pendulum is as the force of gravity, and the square of the time of vibration; and the length being given, the force of gravity is reciprocally as the square of the time of vibration. Fig. 132.

But the force of gravity is also as the square of the distance from the earth's center; therefore the time of vibration of the same pendulum, is as the distance from the earth's center: and the number of vibrations in a given time, reciprocally as that

distance. Therefore $r : \frac{1}{b} :: r+a : \frac{1}{b-c}$, and

$$\frac{r+a}{b} = \frac{r}{b-c}, \text{ and } r+a = \frac{br}{b-c}. \text{ Whence}$$

$$a = \frac{br}{b-c} - r = \frac{cr}{b-c} = 9697.$$

P R O B. CXC VII.

A ball projected from the top of a tower, at an elevation of 31 deg. above the horizon, did in 9½ seconds fall 2000 feet from its base; to find the height. 133.

Let $X=VB$ the tower's height, $BA=d$ the distance, $b=\text{tang. DVC}=31^\circ$, $t=9\frac{1}{2}$ the time, $f=16.1$; then

In the time t , the ball without gravity would arrive at D , and in the same time it would descend through DA . Whence $1 : f :: tt : ttf = DA$ by the laws of falling bodies.

And in the triangle DVC , $1 : b :: d : db = DC$, and $DC + CA = DA$, or $db + x = ttf$, and $x = ttf - db = 149$.

Fig.

PROB. CXCVIII.

If a ball be dropped from the top of a tower a mile high, on the side facing the east, in latitude $51\frac{1}{2}$; where will it fall?

134. Let the body fall at D, whilst the tower by the rotation of the earth is carried to IC. Now by the laws of centripetal force, the area AIE, which the body, moving in the circle AIF describes; is equal to the area AGDE, which the body moving in the curve AGD describes in the same time, that is, in the time of falling through AB. Hence the area $AGI = \text{area } EGD$; and $AGDF = EIF$. But by reason of the small distance BD, the curve AGD (which otherwise would be an ellipsis) is nearly a parabola; and the area of $AFD = \frac{1}{2} AF \times AB = \frac{1}{2} FI \times AE$, the area of the sector EIF. First, let A be a place in the equinoctial.

Put $BE = r = 21000000$ feet, $AB = m = 5280$, $f = 16.1$, $t = 24$ hours $= 86400''$, $c = 3.1416$, $a = DC$, $p = \text{cof. } 51\frac{1}{2}$. Then by the laws of falling

bodies, $\sqrt{f} : 1 :: \sqrt{m} : \sqrt{\frac{m}{f}} = \text{time of falling}$

through AB. And $t : 2rc :: \sqrt{\frac{m}{f}} : BC =$

$\frac{2rc}{t} \sqrt{\frac{m}{f}} = d$, and $BD = a + d$; also by similar

sectors, $r : r + m :: a + d : \frac{m+r}{r} \times \overline{a+d} = AF$.

And $r : a :: r + m : \frac{r+m}{r} a = FI$. Therefore

$\frac{r+m}{3r} \times \overline{a+d} \times m = \frac{r+m}{2r} a \times \overline{r+m}$; therefore

$\overline{a+d} \times \frac{m}{3} = \frac{r+m}{2} a$, and $2am + 2dm = 3ra +$

$3ma$, and reduced $a = \frac{2dm}{3r+m} = 4.64$; and $pa =$

2.88 for the lat. $51\frac{1}{2}$.

PROB.

PROB. CXCIX.

There are two islands A, C; at C is a castle. A ship from A to C keeps pace with the waves of the sea, 100 in number, from A to B. At B she fires a gun, which echoes back from the castle to B, in 3 seconds; and the time of sailing from B to C was 3 minutes; to find the distance AC.

Let $b=100$, $c=3''$, $d=3'$, $f=39.2$ In. the length of a second pendulum, $a=1142$ feet, the velocity of sound in a second, $AC=y$, and x =breadth of a wave.

Then by the motion of pendulums, $\sqrt{f}:1::\sqrt{x}:\sqrt{\frac{x}{f}}$ =time of vibration of the pendulum x .

And $\sqrt{\frac{x}{f}}:x::1'':\sqrt{fx}$ =a space. But (by the principles of philosophy) while the pendulum x vibrates once, the ship or a wave runs through the breadth x ; or in 1 second runs through the space \sqrt{fx} .

And $1'':\sqrt{fx}::d':d\sqrt{fx}$ =CB.

Also by the motion of sound $1'':a::\frac{1}{2}c:\frac{ca}{2}$ =CB, for the echo returns with the same velocity the sound went. Therefore $d\sqrt{fx}=\frac{ca}{2}$, and

$ddf x = \frac{ccaa}{4}$, and $x = \frac{ccaa}{4ddf}$; therefore $AB =$

$bx = \frac{bccaa}{4ddf}$, and $y = \frac{ca}{2} + \frac{bccaa}{4ddf}$.

PROB. CC.

136. *Supposing a planet and its satellite to move in circular orbits, it is required to find, whether the path of a satellite is concave or convex to the sun, when it is in a line between the sun and its primary.*

At the time of the conjunction, if the planet and satellite, both describe very small arches in the same time, whose versed sines are equal, the satellite will then move in a right line. Let ABC be the orbit of the planet, EF that of the satellite; whilst the planet moves through AB, the orbit of the satellite EF is moved into the position ef, and the satellite has moved from e to o. Draw BD, on perpendicular to AE, B ϵ ; and CG perpendicular to AG. Now put AD = ne, then since the center (of the orbit EF,) A is advanced to B, nearer to the line CG, by the distance AD; and the point o is receded from the same line CG, by the distance en equal to AD; it is plain, E and o are equidistant from GC, and Eo is a right line, or the satellite E, o, at that time moves in a right line.

Let $a = eo$, $b = AB$, r , $s =$ the radii of eo , AB .
Hence $AD = \frac{bb}{2s}$, and $en = \frac{aa}{2r}$, and $\frac{bb}{2s} = \frac{aa}{2r}$.
Put $p =$ periodic time of the satellite, $q =$ that of the primary; $c = 3.1416 \times 2$. Then $cr : p :: a :$
 $\frac{pa}{cr} =$ time of describing a ; and $\frac{bq}{cs} =$ time of describing b , then $\frac{pa}{r} = \frac{qb}{s}$, and $\frac{ppaa}{2rr} = \frac{qqbb}{2ss}$, divide this by the former equation, and $\frac{pp}{r} = \frac{qq}{s}$, or
 $pp = \frac{rqq}{s}$. Therefore as pp is greater, equal, or lesser

less than $\frac{rqq}{s}$, then the satellite's orbit is concave, streight, or convex towards the sun, in its conjunction. Fig. 136.

P R O B. CCI.

To find the divisions of a monochord, to sound all the half notes, according to equal intervals of sound; and also to find the variations between these and the strict harmonic divisions.

It is well known an octave is divided into 6 ¹³⁷ whole tones, or 12 semitones. Let BA be the monochord or vibrating string, C the middle point; then BC will be an octave above BA. Let Bd, Be, Bf, Bg, &c. be the several lengths of the strings sounding the half notes, gradually ascending, above AB, by equal degrees of sound. Then will Ad, de, ef, &c. be all unequal in length; and whatever part Bd is of BA, the same part will Be be of Bd, and Bf of Be, and Bg of Bf, &c. to make the several sounds ascend equally. Therefore BA, Bd, Be, Bf, &c. are a set of geometrical proportionals decreasing, continued to 13 terms, the last of which is BC. Also Ad, de, ef, &c. are a set of geometrical proportionals in the same ratio. Also Ad, Ae, Af, Ag, &c. are also a set of geometrical proportionals increasing.

Put BA = 1, BC = $\frac{1}{2}$, Bd = x. Then BA (1) : Bd (x) :: Bd (x) : Be = xx; likewise Bf = x³, Bg = x⁴, &c. and BC = x¹² = $\frac{1}{2}$. And $x = \sqrt[12]{\frac{1}{2}} = .9439$.

Or, put X = log : x. Then $X = \frac{\log : \frac{1}{2}}{12} = -1.9749142$, consequently 2X, 3X, 4X, &c. = logarithms of x², x³, x⁴, &c. Therefore x = $\frac{1}{11} .9439$.

Fig. 9439, $x^1 = .9809$ for a mean tone, &c. and the 137. rest are as in the following table.

The harmonic divisions of the monochord, to found the pure concords will be, as follows; the lesser third $= \frac{5}{8}$, greater third $\frac{4}{3}$, fourth $\frac{3}{4}$, fifth $\frac{3}{2}$, lesser sixth $\frac{5}{4}$, greater sixth $\frac{7}{4}$, eight $\frac{1}{2}$; which see in the following table, in decimals.

Names of the chords.	Pure concords.	Equal divisions.	Errors.
whole string	1.0000	1.0000	0.
<i>b</i> second9439	...
<i>#</i> second8909	...
lesser third	.8333	.8409	<i>b</i> $\frac{1}{15}$
greater third	.8000	.7937	<i>#</i> $\frac{1}{15}$
fourth	.7500	.7492	<i>#</i> $\frac{1}{100}$
<i>#</i> fourth7071	...
fifth	.6666	.6674	<i>b</i> $\frac{1}{100}$
lesser sixth	.6250	.6300	<i>b</i> $\frac{1}{15}$
greater sixth	.6000	.5946	<i>#</i> $\frac{1}{15}$
<i>b</i> seventh5612	...
<i>#</i> seventh5297	...
Eight	.5000	.5000	0.

Then to find the errors or variation of the correspondent cords. Let Bt = cord by column 2d, Br = cord by column 3d, rp = a whole tone, n = number of mean proportionals between Br and Bp , then $\frac{1}{n}$ will be the error, for it shews what part rt

is of the whole note rp . Here then $\frac{Bt^n}{Br^{n-1}} = Bp$
 $= Br \times .8909$. For .8909 being a whole note for the

the string 1, $Br \times .8909$ will be a note for the string Br . Therefore $\left(\frac{Br}{Br}\right)^n = .8909$. And $n \times \log:$

$$\frac{Br}{Br} = \log : .8909 = -1.94983; \text{ and } n =$$

$$\frac{-1.94983}{\log:Br - \log:Br} = \frac{.05017}{\log:Br - \log:Br}.$$

$$\log:Br - \log:Br = 000500, \text{ and } n = \frac{.05017}{.000500} =$$

$$100; \text{ whence the error} = \frac{1}{100}.$$

But as this variation bears but a small proportion to the length of the string, there will be no need to make use of logarithms. For since $1 - .8909 = .1091$ is the length of a note when the string is 1; therefore $.1091 \times Br =$ a note for the

string Br . Whence $\frac{rt}{rp}$ or $\frac{rt}{tp} =$ the error, or

$$\text{which is the same thing } \frac{Br - Bt}{.1091 Br} = \text{the error.}$$

$$\text{As in the fifth, } Br - Bt = .6674 - .6666\bar{6} = .0007,$$

$$\text{and } .1091 \times Br = .0727, \text{ and } \frac{.0727}{.0007} = 100, \text{ nearly,}$$

$$\text{or } \frac{7}{727} = \frac{1}{100} = \text{the error.}$$

Or shorter thus. Since $Br - Bt =$ twice the difference of two adjoining numbers in col. 3. or = difference of two numbers 2 degrees distant, taking one greater and the other less than the proper

$$\text{note; therefore } \frac{Br - Bt}{Br - Bp} = \text{the error.}$$

$$\text{As in the fifth, } \frac{6674 - 6666}{7071 - 6300} = \frac{8}{771} = \frac{1}{97}$$

$$\text{the error. And in a greater third, } \frac{.8000 - .7937}{.8409 - .7492}$$

$$= \frac{.0063}{.0917} = \frac{1}{14\frac{1}{2}} \text{ the error.}$$

Fig. 138. The errors for each concord being thus computed, are set down in the fourth column, which shews the error of the third column, as it differs from the second; those below denoted by (*b*), these above, by (*#*).

In tuning a harpsichord, since the fifth must be 12 times repeated to make 7 octaves, therefore the variation, by tuning by true fifths, will be $\frac{12}{100}$ or about $\frac{1}{8}$ of a note, which is an error that a good ear can discover; and being too sharp, the fifths therefore ought to be tuned as flat as the ear will bear.

Hence the equal division of the notes in an octave is the best system, for the greatest error is in the lesser third and greater sixth, which only amounts to $\frac{1}{13}$ of a note.


P R O B. CCII.


To find the number of beats made in any imperfect concord, in music.

I call that an *imperfect concord* that varies a little from the perfect one, which is made by a harmonical division of the monochord. Thus when the lengths of the strings are 4 and 5, you have the perfect cord (a greater third), but vary one length as 4, making it 3.99, and you will have an imperfect cord attended with beats.

A *beat* is a jarring sound made by the irregular vibrations of two strings, sounding together, when the due period, or coincidence of their vibrations is interrupted. Its noise is such as this *waw, aw, aw, aw*, or *yá, yá, yá, yá, yá*. Our business is to find in how many vibrations this perturbation happens, or how many *yaws* in a second of time.

Let





Let the line AZ represent one second of time; Fig. 139. and suppose it divided on the under side, into the number of vibrations of the lower note or base, at A, B, C, D, &c. and the upper side into the number of vibrations of the upper note, at *a*, *b*, *c*, *d*, &c. Now if any number of divisions on the under side coincide with any number of divisions on the upper, constantly and regularly, as at C and *d*, E and *g*, &c. then the concord is pure, and there is no beat. But when the points *b*, *c*, *d*, &c. are any of them dislocated, and gets to the other side of its corresponding one; then the succession of the short harmonic periods of coincidence is disturbed; and this causes the noise called a beat, such as happens at X and Y. For *c*, *f*, *i*, &c. are continually approaching to B, D, F, &c. till they fall in at X, Y, and change sides: where Bc or B*b*, is supposed the least distance, in the first harmonical period AC, supposing *ad* was to coincide with AC. Therefore at the points X, Y, the succession of the harmonical periods are confused, (and that periodically,) which spoils the harmony.

Now to find the length of this period. Let AC be one harmonical period, that is, when *d* coincides with C, as in the pure concord. In this false cord we must find the time *dC*, which is gained or lost in the time AC. And from thence compute in what time, Bc (the nearest distance), would be gained or lost; in and that will be the time required.

Let *n* = number of parts AB, BC, &c. or its number of vibrations.

t = number of vibrations of the upper string in the perfect cord.

c = length of its string on the monochord.

r = number of parts *ab*, *bc*, *cd*, &c. or its number of vibrations.

I i 3 b =

Fig.

139.

 b = length of its string on the monochord. $\frac{p}{q}$ = number expressing the concord, = $\frac{3}{4}$ for the fourth, or = $\frac{2}{3}$ for the fifth, &cThen $AB = \frac{1}{n}$, and $AC = \frac{p}{n}$; also $ab = \frac{1}{r}$,and $ad = \frac{q}{r}$. Then $AC - ad = \frac{p}{n} - \frac{q}{r} = dC$.Then if $dC \left(\frac{p}{n} - \frac{q}{r} \right)$ be lost or gained in the time $AC \left(\frac{p}{n} \right) :: AB$ will be lost in thetime $\frac{\frac{p}{n}}{\frac{p}{n} - \frac{q}{r}} \times AB = \frac{pr}{pr - qn} \times AB = \frac{pr}{pr - pt}$ $\times AB = \frac{r}{r-t} \times AB$, and Bc will be lost sooner, in proportion of AB to Bc , that is, in the time $\frac{r}{r-t} \times Bc$, which is the time of the period. Butby the laws of vibration, $r : t :: \frac{1}{b} : \frac{1}{c} :: c : b$,and $r-t : r :: c-b : c$; whence $\frac{r}{r-t} \times Bc =$ $\frac{c}{c-b} \times Bc =$ the periodic time of the beats. Andif AZ be divided by the periodic time, you willhave $\frac{c-b}{c} \times \frac{AZ}{Bc} =$ number of beats in a second.But $Bc = \frac{AB}{q}$, $AZ = n \times AB$, therefore $\frac{AZ}{Bc} =$ $\frac{nq \times AB}{AB} = nq$. Whence $\frac{c-b}{c} \times nq = \frac{c-b}{c} \times pt =$

number of beats in a second.

Hence,

Hence, from the length of the string or division Fig. of the monochord, as given in the table of the last 139. problem, and having also the number of vibrations; the beats will be found, as in this table. Where the ground or lowest note is F the cliff-note of the base.

Cords.	Vibrations.	<i>b</i>	<i>c</i>	<i>p.</i> <i>q.</i>	Beats.
Eight	600	5000	5000	1 . 2	0
g. sixth	500	5946	6000	3 . 5	13 ♯
f. sixth	480	6300	6250	5 . 8	18 <i>b</i>
fifth	450	6674	6666	2 . 3	1 <i>b</i>
fourth	400	7492	7500	3 . 4	1½ ♯
g. third	375	7937	8000	4 . 5	21 ♯
f. third	360	8409	8333	5 . 6	15 <i>b</i>
Base F	300	10000	10000	1 . 1	0

This table shews the beats for all the concords, reckoning upwards from F; when the instrument is tuned according to an equal ascent of notes; where the flats and sharps (*b*, ♯) shew whether the upper note is lower or higher, than the true concord in the last column. In the octave above, the beats will be twice as many; and in the octave below, but half as many; being always proportional to the number of vibrations of the base note. The fifth is most serviceable in tuning, and the number of beats in one second, for the

fifth is $\frac{n}{300}$.

If it be supposed that the beat is not made at the points X, Y, but at some intermediate place, where they fall thicker and more confused; and that at the points X, Y, there is the least imperfection. Yet the periodic time will still be the same, whatever part of the cycle XY it falls in. When the

Fig. cycle XY is very short, the single beats are imperceptible, and we hear nothing but a disagreeable noise. All the concords beat, but being exceeding quick, they are not perceived singly; and being regular throughout, they exhibit an agreeable harmony.

When the pitch of the two notes are not altered, the beats succeed one another in equal times, but altering either of them nearer to a perfect harmony, the beats succeed in longer times, and the nearer the longer, till at last they vanish, when the concord is perfect. All the beats are heard in organs; but only half of them are heard in stringed instruments.



S E C T. XII.

Problems relating to Series.

P R O B. CCIII.

Given the diameter of a circle; to find the side of any regular poligon, inscribed in it.

LET d = diameter, n = number of sides, 140.
 x = side of the figure, EB. By Trigonometry, $\frac{3 \cdot 14159d}{2n} = \text{arch DE} = a$, by substitution.

And (Trig. I. 12.) half the side, or EA = $a - \frac{4aa}{6dd}$ A — $\frac{4aa}{20dd}$ B — $\frac{4aa}{42dd}$ C, &c.

And 2EA or EB or $x = 2a - \frac{4aa}{6dd}$ A — $\frac{4aa}{20dd}$ B — $\frac{4aa}{42dd}$ C —

$\frac{4aa}{72dd}$ D, &c.

Or thus,

By a table of natural sines, find the sine of $\frac{180}{n} = s$; then $x = ds$.

P R O B. CCIV.

Suppose $x^3 - cx^{\frac{1}{2}} + x^{\frac{2}{3}} + bx^{\frac{3}{4}} = dccx$; to find x .

Divide by the least power of x , that is, by $x^{\frac{1}{2}}$, and $x^{\frac{5}{2}} - c + x^{\frac{1}{6}} + bx^{\frac{5}{4}} = dccx^{\frac{1}{2}}$. Take r the nearest root, and put $r + e = x$.

Then

Fig. Then

$$\left. \begin{aligned} x^5 &= r^5 + \frac{5}{2} r^3 e + \frac{15}{8} r^1 e^3 \\ -c &= -c \\ + x^{\frac{5}{6}} &= r^{\frac{5}{6}} + \frac{1e}{6r^{\frac{5}{6}}} - \frac{5ee}{72r^{\frac{1}{6}}} \\ + bx^{\frac{5}{4}} &= br^{\frac{5}{4}} + \frac{be}{4r^{\frac{3}{4}}} - \frac{3bee}{32r^{\frac{1}{4}}} \\ - dcx^{\frac{5}{2}} &= -dcr^{\frac{5}{2}} - \frac{dcce}{2r^{\frac{1}{2}}} + \frac{dccee}{8r^{\frac{3}{2}}} \end{aligned} \right\} = 0.$$

That is, $p + qe + see = 0$, by substitution.

Whence

$$e = \frac{-\frac{p}{s}}{\frac{q}{s} + e}, \text{ which may be repeated for}$$

more exactness.

Or thus,

Seek the least common dividend of the denominators of the indices of x , and reduce the equation, which will become $x^{\frac{10}{12}} - c + x^{\frac{2}{12}} + bx^{\frac{3}{12}} - dccc^{\frac{6}{12}} = 0$. Put $x = e^{12}$; then the equation becomes $e^{10} - c + e^2 + be^3 - dccc^6 = 0$, or $e^{10} - dccc^6 + be^3 + e^2 = c$, and the root extracted gives e , and consequently x is had.

PROB. CCV.

141. Given the sides AC, CB, of the triangle ACB; and the ratio of AB to the arch CE is given; to find AB.

Let AC = $r = 14$, CB = $s = 22$, AB = x , and AB : CE :: 10 : 4, whence arch CE = $\frac{1}{10} x$. Let $y = \text{cof. CAB}$. Then

Then (Trig. case 5.) $rr + xx - 2rsy = ss$, whence Fig.

$$y = \frac{rr + xx - ss}{2rs} = \text{cof. A to the radius } 1, \text{ and } ry = \text{141.}$$

$$\frac{rr - ss + xx}{2s} = \text{cof. A to the radius } r. \text{ But (Trig.}$$

$$\text{I. 12. cor. 1.) } \text{cof. A} = r - \frac{aa}{2r} + \frac{a^4}{24r^3}, \text{ \&c.} = r$$

$$- \frac{\frac{-2}{4x}}{200r} + \frac{\frac{-4}{4x^3}}{24 \cdot 10000r^3} - \frac{\frac{4^6}{4x^5}}{720 \cdot 1000000r^5} \text{ \&c.}$$

$$= \frac{rr - ss}{2s} + \frac{xx}{2s}, \text{ and transposing,}$$

$$\frac{\frac{1}{2s} + \frac{4^2}{200r} \times x^2 - \frac{4^4}{240000r^3} \times x^4 + \frac{4^6}{720000000r^5} \times x^6$$

$$\text{\&c.} = r - \frac{rr - ss}{2s}, \text{ or } Ax^2 + Bx^4 + Cx^6 \text{ \&c.} = b,$$

by substitution, then (Prob. lxii. I.) $xx =$
 $\frac{b}{A} - \frac{B}{A^3} b^2 + \frac{2BB - AC}{A^5} b^3 \text{ \&c.} = 836.95$, and
 $x = 28.93$.

P R O B. CCVI.

Given the arch of the circle BHE, and the sine BD; 142.
to find the radius BC.

Let BHE = $d = 8$, BD = $s = 3$. Take an angle p nearly equal to ACB, $a =$ sine, $b =$ its cosine, rad. = $1.n = .0174533$, $c = 3.1415926$, then $np =$ arch belonging to the angle p . Let $p + x =$ true angle ACB; then $np + nx$ or $np + z =$ correspondent arch, (putting $z = nx$): And (Trig. I. 13.), the sine of $np + z = a + bz - \frac{az^2}{2} - \frac{bz^3}{6} + \frac{az^4}{24} +$
 $\text{\&c.} = \frac{s}{d} c - \frac{s}{d} np - \frac{s}{d} z$ per quest. that is,

Fig. $\frac{s}{d} + b \cdot z - \frac{a}{2}zx - \frac{b}{6}z^2 + \frac{a}{24}z^4 \&c. = \frac{s}{d}c -$
 $\frac{s}{d}np - a$, or $Az + Bzz + Cz^3 + Dz^4, \&c. = R$.

Affume $p = 55^\circ$, then a and b will be known, and $R = .0010292$, and (Prob. lxii. I.) x or

$$ux = \frac{R}{A} - \frac{B}{A^3}R^2 + \frac{2BB-AC}{A^5}R^3 \&c. = -$$

$.001084$, and $x = -.0621$ degrees, which is, $3' 43''\frac{1}{2}$; therefore $p+x = 54^\circ 56' 16''\frac{1}{2} = \angle ACB$. Hence $BC = 3.66513$.

P R O B. CCVII.

143. The ordinate AC, and curve BC are given; and the equation of the curve is $\frac{y}{a} = \text{hyp. log.} \frac{z + \sqrt{aa + zz}}{a}$, and $a+x = \sqrt{aa + zz}$; where $AC = y$, $BC = z$; to find AB.

Let $y = 6 = b$, $z = 9 = g$, $a = r + e$, taking r the nearest value of a . Then h. log :

$$\frac{g + \sqrt{rr + 2re + ee + gg} \&c.}{r + e} = \frac{b}{r + e} = \frac{b}{r} - \frac{be}{rr} +$$

$$\frac{bee}{r^3} \&c. \text{ And (putting } ff = rr + gg), \text{ by evolu-}$$

$$\text{tion, log: } \frac{g + f + \frac{re}{f} + \frac{gg}{f^3}ee}{r + e} = \frac{b}{r} - \frac{be}{rr} + \frac{bee}{r^3}, \text{ and}$$

$$\frac{f + g + \frac{re}{f} + \frac{gg}{f^3}ee}{r + e} = \text{number of the hyp. log :}$$

$$\frac{b}{r} - \frac{be}{rr} + \frac{bee}{r^3} = n - \frac{nbe}{rr} + \frac{nbee}{r^3}$$

$$+ \frac{nbh}{2r^4} ee \text{ (putting } n =$$

h. log: $\frac{b}{r}$) (Prob. lxxxv. I.) And

And multiplying by $r+e$,

Fig.
143.

$$g+f+\frac{re}{f} + \frac{gg}{2f^3} ee = rn - \frac{nbe}{r} + \frac{nb^2 ee}{2r^3} \\ + ne + \frac{nbce}{rr} - \frac{nbce}{rr}$$

And reduced

$$\frac{r}{f} + \frac{nb}{r} - n \times e + \frac{gg}{2f^3} - \frac{nbb}{2r^3} \times ee = rn - f - g.$$

Assume $r=3.7$; then $e=-.00112$, and $a=3.69885$, and $x=\sqrt{aa+zz}-a=6.0316$; substitute this value of a for r in the last equation, and the operation repeated, gives a still more exact.

P R O B. CCVIII.

Given the length of a pendulum, and the arch it describes; to find the time lost by describing a greater arch.

Let r =length of the pendulum, c =cord of half the arch it describes, C =any other cord, t =time of falling through $2r$. $P = \frac{3.1416}{2} t$. Then by mechanics it is found that the time of 1 vibration,

is $= P \times \left(1 + \frac{cc'}{16rr} + \frac{9c^4}{4^5 r^4} \right) \&c.$ for the cord c .

And $= P \times \left(1 + \frac{CC}{16rr} + \frac{9C^4}{4^5 r^4} \right) \&c.$ for the cord C .

and $P \times \left(\frac{CC-cc}{16rr} + \frac{9}{4^5 r^4} \times \overline{C^4-c^4} \right) \&c. =$
time lost in one vibration for the cord c . But when

Fig. when c is 0, P is the time of one vibration, which does not sensibly differ from a vibration for the cord c . Therefore since $86400 =$ the number of seconds in 24 hours, therefore $\frac{86400}{P} =$ number of vibrations for the cord c , in 24 hours. Therefore $\frac{86400}{P} \times P \times \frac{CC-cc}{16rr}$ &c. or $86400 \times \frac{CC-cc}{16rr} + \frac{9}{4^5 r^4} \times \overline{C^4 - c^4}$ &c. $=$ seconds lost in 24 hours; that is, $\frac{5400}{rr} \times :$

$\frac{CC-cc}{16rr} + \frac{9}{64rr} \times \overline{C^4 - c^4}$ &c. $=$ seconds lost in 24 hours; and $\frac{5400}{rr} \times \overline{CC-cc}$, is nearly $=$ the seconds lost in 24 hours.

If r swings seconds, then $r = 39.2$, and the time lost in 24 hours is nearly $= 3.52 \times \overline{CC-cc}$.

Cor. If $c = 0$, and $C =$ cord of 90° . A pendulum vibrating in the double arch of 90° , will lose 4 h. 20 min. in 24 hours time.

And if $c = 0$, then to find the length of a pendulum vibrating in the arch of C in the same time. Let $r =$ pendulum vibrating in the very small arch, $x =$ pendulum vibrating in the arch of C . Then the lengths being as the squares of the times of vibration, we shall have in the first case \sqrt{r} for t , and in the second \sqrt{x} for t ; whence in the first case $P = \frac{3.1416}{2} \sqrt{r}$, in the second $P = \frac{3.1416}{2} \sqrt{x}$. And the times being equal we shall have $P \times 1$ or $\frac{3.1416}{2} \sqrt{r} = P \times$

Fig. $1 + \frac{CC}{16rr}$ or $\frac{3.1416}{2} \sqrt{x} \times 1 + \frac{CC}{16rr}$, or \sqrt{r}
 $= \sqrt{x} \times 1 + \frac{CC}{16xx}$, and $r = x + \frac{CC}{8x}$, which re-
 duced is $rx - xx = \frac{1}{2}CC$; whence x will be found.
 And on the contrary x being given, r will be
 found.

P R O B. CCIX.

Given the latitude sailed from, the departure, and
 difference of longitude; to find the difference of
 latitude.

Let d = departure, l = diff. longitude, x = arch of
 latitude come to, z = its mer. parts, a = the given
 lat. m = its mer. parts.

Then by *Mercator's Sailing*; as diff. lat. ($a - x$)
 : mer. diff. latitude ($m - z$): : d : l , whence $al - lx =$
 $dm - dz$, and $dz - lx = dm - al$. But Dr. *Halley's*
 Series for the meridional parts of x , is

$x + \frac{1}{6}x^3 + \frac{1}{24}x^5 + \frac{61}{5040}x^7$ &c. Therefore

$dx - lx + \frac{d}{6}x^3 + \frac{d}{24}x^5 + \frac{61d}{5040}x^7$ &c. =

$dm - al$. And by reversion of series (Prob. lxii.)
 x will be found; then $3438x$ = latitude in mi-
 nutes.

Or thus,

Seek another latitude, by the table of meridio-
 nal parts, such, that the proper difference of la-
 titude divided by the mer. diff. latitude, will be
 equal to the quotient $\frac{d}{l}$, which is easily done by
 a few trials; and that is the other latitude.

P R O B.

PROB. CCX.

144. The curve BMD is described with a pair of compasses upon the surface of a cylinder, which is afterwards stretched into a plane; to find the ordinate PM.

Let d = diameter of the cylinder, $a = AB$ the extent of the compasses, $AP = x$, $PM = y$, $v =$ chord, whose arch is y . Then (Geom. II. 21.) $aa - xx = vv$. But (Trig. I. 12. cor. 2.) $v = y -$

$$\frac{yy}{2.3dd} A - \frac{y^2}{4.5dd} B - \text{\&c.} \quad \text{Whence } \sqrt{aa - xx}$$

$$= y - \frac{yy}{2.3dd} A - \frac{yy}{4.5dd} B - \frac{yy}{6.7dd} \text{\&c.} \quad \text{and}$$

by reversion of series y is had.

If the arch was in a given ratio to the chord, the figure would be an ellipsis; but as this is not so, the curve will be a mechanical one.



S E C T. XIII.

Problems concerning exponential quantities.

P R O B. CCXI.

Some maids driving a flock of sheep, were asked, how many they had? To which they answered, that if the flock was equally divided among them, the share of each would be twice as many as there were maids. And if the terms of this double progression 1, 2, 4, 8, &c. be counted, as often as there are maids; the last term will be the number of sheep.

LET a = sheep, e = maids. Then $\frac{a}{e} = 2e$, and the e^{th} term of the progression 1, 2, 4, 8, &c. = 2^{e-1} (Propor. 25.), therefore $2^{e-1} = a$, per quest. Whence $a = 2ee$, and $2^e = 2a$, and expunging a , $2^e = 4ee$, or $2^{e-2} = ee$. Therefore $e^{-2} \times \log: 2 = 2 \log: e$, or $.30103e - .60206 = 2 \log: e$, and $.150515e - \log: e = .30103$. Then to find e (by Rule 5. Prob. xcii.), assume $e = 6$, then $.1505e - \log: e = .125$, which should be $.301$, and the error is $-.176$.

Again, assume $e = 7$, then $.1505e - \log: e = .208$, and the error is $-.092$. Then $\frac{1 \times .092}{.176 - .092} = 1.1$; therefore $e = 8.1$ nearly.

Suppose $e = 8.1$; then $.1505e - \log: e = .3106$, and the error = $+.0096$. And the correction = 1.04 , and $e = 8.1 - 1.04 = 7.996$; and $e = 8$ exact, and $a = 128$.

PROB. CCXII.

Two travellers at 150 miles distance set out to meet one another. In the several days, A goes at this rate, 5, 10, 20, 40, &c. B goes 6, 10, 14, 18, &c. miles; to find in what time they will meet.

Let x = days, (by Geom. Prog.) A's last day, will be $5 \times 2^{x-1}$; and his journey $5 \times 2^x - 5$. And (Arith. Prog.) B's last day is $6 + 4x - 4$ or $4x + 2$, and $\frac{12 + 4x - 4}{2} \times x$, or $2x^2 + 4x =$ B's journey. Whence $2^x \times 5 - 5 + 4x + 2x^2 = 150$, and $2^x + \frac{4}{10}x^2 + \frac{8}{10}x = 31$. And $2^x = 31 - \frac{8}{10}x - \frac{4}{10}xx$. And $\log: 2^x$ or $x \times \log: 2 = \log: 31 - \frac{8}{10}x - \frac{4}{10}xx$. By trials x will be found greater than 4; let $n=4$, and $n+v=x$, $b=4$, $c=\log: 2$. Then $cn + cv = \log: 31 - \frac{8}{10}n - \frac{8}{10}v - \frac{4}{10}nn - \frac{8}{10}nv - \frac{4}{10}vv$. But the number belonging to $cn + cv = cn \times : 1 + mcv + \frac{mcv^2}{2}$ &c.

(Prob. lxxxv.) whence $cn + ccmmv + \frac{c^2nmv^2}{2}$ &c.

$$= 31 - \frac{8}{10}n - \frac{4}{10}nn - \frac{8}{10} + \frac{8}{10}n \times v - \frac{4}{10}vv.$$

And

And reduced, $ccmnv + \frac{ccmn}{2} vv = 31$

$$\begin{array}{r}
 + \frac{8}{10} \quad + \frac{4}{10} \quad - \frac{8}{10} n \\
 + \frac{8}{10} n \quad - \frac{4}{10} nn \\
 - cn
 \end{array}$$

Which put into numbers, and reverting the series, (Prob. lxii.), v is had $=.32$; then put new n for $x+v$ or 4.32 , and repeat the operation; and at last $v=.3256$, and $x=4.3256$.

PROB. CCXIII.

To find x in this equation, $x^x = 123456789$.

Here x will be found between 8 and 9. Put $n=8$, $n+v=x$, $b=\log:n$, $c=\log:123456789$; then $x \log:x = \log:123456789 = c$, or $n+v \times \log:n+v = c$.

But (Prob. lxxxiv.) $\log:n+v = b + \frac{Mv}{n} - \frac{Mv^2}{2n^2} + \frac{Mv^3}{3n^3} - \&c.$ and $\overline{n+v} \times \log:\overline{n+v} = nb + \frac{nMv}{n} - \frac{nMv^2}{2n^2} + \frac{nMv^3}{3n^3} - \&c.$

$$\left. \begin{array}{l}
 + bv + \frac{Mv^2}{n} - \frac{Mv^3}{2n^2} - \&c. \\
 \end{array} \right\} = c,$$

and transposing and reducing,

$$\frac{b}{M} + 1 \times v + \frac{v^2}{2n} - \frac{v^3}{2.3n^2} + \frac{v^4}{3.4n^3} - \frac{v^5}{4.5n^4} - \&c. = \frac{c-nb}{M}.$$

And extracting the root (Prob. xciii.)

$v=.64002$, and x or new $n=8.64002$ for another operation, which will give $x=8.6400268$.

PROB. CCXIV.

To find the value of x in the equation
 $1000-x \times \log:1000-x = x$.

Put $b=1000$, $x=a+v$, $b-a=g$, $p=\log:g$;
 then $\overline{g-v} \times \log:g-v = a+v$. And substituting
 the logarithmic series instead of $\log:g-v$ (Prob.
 lxxxiv.), $\overline{g-v} \times : p - \frac{v}{g} - \frac{v^2}{2gg} - \frac{v^3}{3g^2} - \frac{v^4}{4g^3} \&c.$
 $= a+v$, which multiplied and reduced is, $gp-a$
 $- \overline{p+2} \cdot v + \frac{vv}{2g} + \frac{v^3}{6g^2} + \frac{v^4}{12g^3} + \frac{v^5}{4.5g^4} + \frac{v^6}{5.6g^5}$
 $\&c. = 0$. Assume $a=836$, and extracting the root
 (Prob. xciii.) $v=.05315$; and $x=836,05315$.

PROB. CCXV.

To find x in the equation

$$\log:1000-x = \frac{x}{1000-x} - \frac{1000}{x}$$

Put $n=1000$, $x=a+v$, $g=n-a$, $p=\log:g$.
 Since x is nearly $=860$, assume $a=860$; then
 the equation is $\log:g-v = \frac{a+v}{g-v} - \frac{n}{a+v}$. But

$$\log:g-v = p - \frac{v}{g} - \frac{vv}{2g^2} \&c. \text{ Whence}$$

$$\frac{a+v-ng+nv}{a+v \times g-v} = p - \frac{v}{g} - \frac{vv}{2gg} - \frac{v^3}{3g^2} \&c.$$

Which equation reduced gives

$$agp - 3av + \frac{a}{2g}v^2 + \frac{a}{6gg}v^3 \&c. = 0.$$

$$\begin{array}{r} -aa - n \\ +ng + gp \\ -ap \end{array} \quad \begin{array}{r} -p \\ -2 \end{array} \quad + \frac{1}{2g}$$

In numbers,

Fig.

$4626.3 + 7138v + 3.8702v^2 - .01088v^3 = 0$;
 or $1 + 1.543v + .000836v^2 - .0000023v^3 = 0$;
 whence, by extracting the root, $v = .64822$, and
 $x = 859,35178$.

PROB. CCXVI.

Having given the equations $x^{x+y} = y^n$, and
 $y^{x+y} = x^m$; to find x and y .

From the first equation $y = x^{\frac{x+y}{n}}$, and from the
 second, $y = x^{\frac{m}{x+y}}$; therefore $x^{\frac{x+y}{n}} = x^{\frac{m}{x+y}}$. And
 equating the indices $\frac{x+y}{n} = \frac{m}{x+y}$, and $x+y =$
 \sqrt{mn} . Whence $y = x^{\frac{\sqrt{mn}}{n}} = x^{\frac{\sqrt{m}}{n}}$. Therefore
 by the first equation, $x^{x+x^{\frac{\sqrt{m}}{n}}} = y^n = x^{\sqrt{mn}}$,
 and again equating their indices, $x + x^{\frac{\sqrt{m}}{n}} = \sqrt{mn}$.
 Then x being had y is known from the equa-
 tion $y = x^{\frac{\sqrt{m}}{n}}$.

To find x put $x = v^n$, then $x^{\frac{\sqrt{m}}{n}}$ or $x^{\frac{\sqrt{mn}}{n}}$
 $= v^{\sqrt{mn}}$, and $v + v^{\sqrt{mn}} = \sqrt{mn}$. And the root
 may be extracted by logarithms.

P R O B. CCXVII.

To find the value of x in this equation, $X^2 + X = \frac{1}{x}$
 X being the hyperbolic log: of x .

Here x is between 1 and 2, therefore put $x = 1 + v$, then (Prob. lxxxiv. cor. 1.) $X = v - \frac{v^2}{2} + \frac{v^3}{3} - \frac{v^4}{4} \&c.$ Whence $v - \frac{v^2}{2} + \frac{v^3}{3} \&c. + v - \frac{v^2}{2} + \frac{v^3}{3} \&c. = \frac{1}{1+v}$, and multiplying and reducing $v + \frac{3}{2}v^2 - \frac{1}{6}v^3 \&c. = 1$, and by reversion (Prob. lxii.) $v = .56$, and $x = 1.56$.

But because this does not converge fast enough; put $n = 1.56$, and $n + v = x$, $l = .4446858 = \text{hyp. log: } n = m \times \text{log: } n$; then (Prob. lxxxiv. cor. 2.)

$X = l + \frac{v}{n} - \frac{v^2}{2nn} + \frac{v^3}{3n^2}$, whence we shall have

$$l + \frac{v}{n} - \frac{vv}{2nn} \&c. + l + \frac{v}{n} - \frac{vv}{2nn} : \times n + v = 1.$$

And when multiplied and reduced,

$$l + 1 \times nl + l + 1 + l \times v + \frac{l + \frac{1}{2}}{n} vv \&c. = 1.$$

In numbers,

$1.0021921 + 2.5318022v + 1.246593v^2 = 1$; or
 $2.031v + vv = -.0017586$. Whence (Prob. 88.)
 $v = -.0008661$, and $n + v$ or $x = 1.5591339$.

Otherwise thus,

Let $l = .4446858$ the h. log: 1.56, or n , as before, $l + s = X$; then the number (x) belonging to

to $l+s$ or $X=n \times : 1+s + \frac{1}{2}ss + \frac{1}{6}s^3 \text{ \&c.}$

(Prob. lxxxv.); whence $\overline{l+s} + l+s : \times n+ns + \frac{1}{2}nss \text{ \&c.} = 1$; and by reduction,

$$\overline{l+1} \times \overline{ln} + \overline{l+1} + l \times ns + \frac{1}{2}l + 2\frac{1}{2}l + 2 \times ns^2 \text{ \&c.} = 1.$$

In numbers,

$$1.0021921 + 3.949611s + 5.008515s^2 \text{ \&c.} = 1.$$

$$\text{or } .78858s + ss = .00043768;$$

And extracting the root (Prob. lxxxviii.) $s = .0005549$, and $l+s$ or $X = .4441309$, and $\frac{X}{m} =$

$.1928836$ the com. log: x ; or else $\frac{s}{m} =$

$.0002410$, and since com. log: $1.56 = .1931246$; therefore $.1931246 - .0002410 = .1928836$ the common log: x . Whence $x = 1.559134$.

P R O B. CCXVIII.

To find x in the equation $x^{xx} = 123456789 = b$.

Put $b = 123456789$, and by a few trials you will find x near 2.8 , put $n = 2.8$, $n+v = x$, $l = \log: n$, $ml = \text{hyp. log: } n$.

Then (Prob. lxxxv. cor. 6.) $x^x = \frac{n^n \times}{1+mlv+v}$. Put $r = n^n$, $e = \frac{n^n \times}{mlv+v}$, then $x^x = r+e$; let this be an index, then $x^{x^x} = \frac{x^{r+e}}{n+v} = \frac{r^{r+e}}{n+v} =$ (by the same cor.) $n^r \times \frac{1+mlv+v}{n} = b$ per quest. Then

restoring the values of r and e , $n^n \times : 1 + mln^n$

Fig.

$$\times \overline{mlv+v} + \frac{n^v}{n} := b. \text{ Put } g = ml^n \times \overline{ml+1},$$

then $\overline{n^n} \times 1 + gv + n^{n-1}v = b$, and by reducing,

$$v = \frac{b - n^n}{n^n \times g + n^{n-1}}; \text{ here } n^n = 97620000,$$

$ml = 1.02962$, $g = 37.3368$, $n^{n-1} = 6.3810$, therefore $v = \frac{25830000}{4266000000} = .006054$.

Or let $\frac{b}{n^n} = f$. Then $v = \frac{f-1}{g+n^{n-1}} =$

$\frac{.26466}{43.7178} = .006054$; then $n+v$ or $x = 2.806054$ nearly; or put $n = 2.806054$ for another operation.

This problem is easily resolved by rule 5, problem xcii. by making several suppositions for the value of x , and finding the correction every time; and so you will continually approximate to the true value.

P R O B. CCXIX.

If X be the log: x , it is required to find x , in the equation $x^X + X^x = 100$.

Let $n+v=x$, $l=\log:n$, $l+s=\log:n+v$, $L=\log:l$. Then (Prob. lxxxv.) $n+v=n+nms$ &c. whence $\frac{n+nms}{n+nms}^{l+s} + \frac{l+s}{l+s}^{n+nms} = 100$. But (Prob. lxxxv. cor. 6.) $\frac{n+nms}{n+nms}^{l+s} = n^l \times \frac{1+m's+lms}{1+m's+lms}$.

$$\text{And } \frac{l+s}{l+s}^{n+nms} = l^n \times \frac{1+m'Lns + \frac{ns}{l}}{1+m'Lns + \frac{ns}{l}}.$$

Therefore

Therefore $n^l \times \frac{1+2mls}{l} + l^n \times \frac{1+m^2Lns}{l} + \frac{ns}{l} = 100.$

Or $n^l \times 2mls + l^n \times m^2Lns + \frac{ns}{l} = 100 - n^l - l^n = d.$

$$\text{And } s = \frac{d}{2mln^l + nLl^n m^2 + nl^{n-1}}.$$

To approach nearly to the value of X, we shall have $X \log: x$ or $XX = \log. x^X$, and $x \log: X = \log: X^x$. Therefore num. of $XX +$ num. of $x \log: X = 100$. By a few trials X is found between 1.25 and 1.26, but nearer 1.26; therefore suppose $l = 1.257$, then $n = 18.072$, $L = .09933$, $n^l = 38.02$, $l^n = 62.41$, $d = -.43$, $2mln^l = 220.1$, $nLl^n m^2 = 594.0$, $nl^{n-1} = 897.4$. Whence $s = \frac{-.43}{1711.5} = -.000251$, and $X = 1.256749$; and $x = 18.0613$.

Here we have sought the logarithm X, for variety; but the number x might have been found, after the manner of the last problem.

PROB. CCXX.

Given $n^{\bar{x}^x} + x^x - x^{-x} + x^{\frac{1}{x}} = 200$; to find x .

Take n very near the root, to be found by frequent trials, and put $n+v=x$, $l = \log:n$, $r = n^n$, $f = ml + 1$, $p = n^n$, $q = mlrf + \frac{r}{n}$. $t = n^{\frac{1}{n}}$, $a = \frac{m^l - 1}{nn}$.

Then

Fig. Then $x^n = n+v^{n+v} = n^n \times \overline{1+mv+v}$ (Prob. lxxxv. cor. 2.) $= r+frv$.

$$\text{And } x^{x^n} = x^{r+frv} = \overline{n+v}^{r+frv} = n^r \times \overline{1+mlrfv+\frac{rv}{n}} = p+pqv, \text{ (ib. cor. 6.)}$$

$$\text{Also } x^{-x} = \frac{1}{x^x} = \frac{1}{r+frv} = \frac{1-fv}{r}$$

$$\text{And } \frac{1}{x} = \frac{1}{n+v} = \frac{n-v}{nn}$$

$$\text{Also } x^{\frac{1}{x}} = \overline{n+v}^{\frac{n-v}{nn}} = n^{\frac{1}{n}} \times \overline{1-\frac{mv}{nn}+\frac{v}{nn}}$$

(ib. cor. 6.) $= t-atv$.

Therefore writing for the several powers of x , their respective values, we have

$$p+pqv+r+frv + \frac{fv-1}{r} + t-atv = 200.$$

$$\text{reduced } v = \frac{200-p-r-t+\frac{1}{r}}{pq+rf+\frac{f}{r}-ta}$$

It easily appears that x is greater than 2, and trying $2\frac{1}{2}$, it will be found a little too small; therefore assume $n=2.27$, whence there will come out $v=-.0009463$, and therefore $x=2.2690537$, which may be put for n , for another operation.

S E C T. XIV.

Problems of Maxima and Minima.

P R O B. CCXXI.

The line AE, and the two points B, C, being given ¹⁴⁵ in position; to find the point P, so that BP + PC may be the least possible.

TAKE the point p extremely near P , and draw Bp , Cp , and also $pD \perp$ to BP , and $pO \perp$ to CP . Then pD is the increment of BP ; and PO the decrement of CP , therefore $DP = OP$, by the nature of the question. And since the hypotenuse Pp is common, $pD = pO$. And $\angle pPD = pPO$, that is $BPA = CPE$; whence the triangles BAP , CEP , are similar. Put $AE = b$, $AB = p$, $CE = q$, $AP = x$, $EP = b - x$; then $AB (p) : AP (x) :: CE (q) : EP (b - x)$ therefore $qx = pb - px$, and $px + qx = pb$, and $x = \frac{pb}{p+q}$, and $b - x = \frac{bq}{p+q}$.

P R O B. CCXXII.

The lines ABC, and CE, being given in position, and ¹⁴⁶ the points A, B, being given; to find the point D in the line CE, where the angle ADB is the greatest possible.

About AB describe a circle to touch the line CE ; then the point of contact D is the point required.

For

Fig. For to any other point E, in the line CE, draw
146. AE, BE, and draw BF. Then the angle AEB is
less than AFB, or its equal ADB (Geom. IV. 13.)

Let $BC=b$, $AC=d$, $CD=x$. Then (Geom.
IV. 21. cor. 2.) $xx=bd$, and $x=\sqrt{bd}$.

P R O B. CCXXIII.

147. To draw the shortest line possible, through a given
point P, placed within the right angle ABC.

Let CPA be the shortest line. Draw PD pa-
rallel to AB, and PF parallel to CB, and let
 $PD=b$, $PF=c$, $CD=x$, $Cc=e$ an extremely small
quantity, $PC=z$.

By the similar triangles CDP, PFA, $x : z ::$

$c : \frac{cz}{x} = AP$, and by the similar triangles CDP,

cHC , $z : x :: e : \frac{xe}{z} = Hc$. Also $z : b :: e : \frac{be}{z}$
 $= HC$. And by the similar triangles PCH, PGA,

$x : \frac{be}{z} :: \frac{cz}{x} : \frac{bce}{zx} = aG$. And by the similar

triangles CDP, aGA, $x : b :: \frac{bce}{zx} : \frac{bbce}{zxx} = AG$.

But $Hc=AG$, that is $\frac{xe}{z} = \frac{bbce}{zxx}$, or $x = \frac{bbc}{xx}$, and

$x^3=bbc$, whence $x=\sqrt[3]{bbc}$.

P R O B. CCXXIV.

148. Given the line EF, and two points A, B; to find
the point D, so that $a \times AD + b \times BD$, may be the
least possible; a, b being given numbers.

Take d infinitely near D, and draw Ad , Bd ;
on which let fall the perpendiculars Dr , Df . Then
will

will $a \times AD + b \times BD = a \times Ad + b \times Bd$; and by subtraction $a \times AD - Ad = b \times Bd - BD$, or $a \times dr = b \times df$. But in the triangles Ddr , Ddf , the hypotenuse Dd is common; therefore $dr : df :: S.dDr : S.dDf :: \text{cof. ADF} : \text{cof. BDE}$. Whence $a : b :: \text{cof. BDE} : \text{cof. ADF}$.

Let AF , BE be perpendicular to EF ; and put $AF = c$, $BE = d$, $EF = n$, $DF = x$, $DE = v$. Then

$$DA (\sqrt{cc+xx}) : 1 : DF (x) : \frac{x}{\sqrt{cc+xx}} = S.DAF$$

$$= \text{cof. ADF}; \text{ then } b : a :: \frac{x}{\sqrt{cc+xx}} :$$

$$\frac{ax}{b\sqrt{cc+xx}} = \text{cof. BDE} = S.DBE. \text{ And } \frac{ax}{b\sqrt{cc+xx}}$$

$$: v :: 1 : (BD) \sqrt{aa+vv}; \text{ therefore } v = \frac{ax \sqrt{aa+vv}}{b\sqrt{cc+xx}}, \text{ and } bv\sqrt{cc+xx} = ax\sqrt{aa+vv}.$$

And squaring, $bbccvv + bbxxvv = aaddxx + aaxxvv$,

but $v = n - x$, put $p = bb - aa$, then $vcc + pxxvv$

$= aaddxx$, or $bbcc + pxx \times \frac{nn - 2nx + xx}{n} = aaddxx$;

reduced, $px^4 - 2pnx^3 + pnxx - 2nbccx + b^2c^2n^2 = 0$.

$$\begin{aligned} &+bbcc \\ &-aadd \end{aligned}$$

P R O B. CCXXV.

Three points A , B , C being given; to find a fourth point D , so that $a \times AD + b \times BD + c \times CD$, may be the least possible; where a , b , c , are given numbers. 149.

Let D be the point sought; with radius CD , describe the circle GDH . Take the point d infinitely near D , and draw Ad , Bd ; on AD , BD , let fall the perpendiculars dr , df . Then supposing

Fig. *ling* CD to be given, $a \times AD + b \times BD$ will be a
 149. minimum. But $a \times Dr$ is the increment of $a \times AD$,
 and $b \times Df$ is the decrement of $b \times BD$, therefore
 $a \times Dr = b \times Df$. But in the right-angled triangles Ddr ,
 Ddf , $Dr : Df :: S.Ddr : S.Ddf ; : S.rDC$, or ADC
 $: S.mDf$ or BDC . Therefore $b : a :: S.ADC :$
 $S.BDC$.

After the same manner, supposing BD given,
 we shall have $c : a :: S.ADB : S.BDC$. There-
 fore when $a \times AD + b \times BD + c \times CD = \text{minimum}$;
 a, b, c , are respectively as the sines of $BDC, ADC,$
 ADB ; or of BDm, ADm, BDr , which makes 180° .
 Therefore if a triangle be made of the 3 lines a, b, c ;
 the angles of this triangle will be equal to the an-
 gles at D , *viz.* that opposite to $a = mDB$, to
 $b = mDA$, to $c = BDr$. Therefore all the angles
 about the point D being given ; the distances $AD,$
 BD, CD will be found by Prob. cxxxii.

P R O B. CCXXVI.

150. *Given the triangle ABD, and the circle CFK whose
 center is A ; to find the point F in the circumfe-
 rence CFK, that the angle BFD may be the great-
 est possible.*

Through the points B, D , describe the circle
 BFD to touch the circle CFK in F , the point
 required. For to any other point C , in the cir-
 cle CK , draw DC cutting BFD in S , and draw
 BS, BC . Then the $\angle BSD$ or $BFD = \angle BCD +$
 CBS ; therefore BFD is greater than BCD .

On BD let fall the perpendiculars, $AH, FI,$
 GE ; G being the center of the circle BFD . Then
 to find its radius GF ; let $BE = ED = b, HE = c,$
 $AH = p, AF = r, GE = x$. Then $AG = \sqrt{p^2 + x^2} + c$

=

$= \sqrt{pp+cc+2px+xx}$, and $GF = \sqrt{pp+cc+2px+xx}-r$, and $BG = \sqrt{bb+xx}$,
 whence $\sqrt{pp+cc+2px+xx}-r = \sqrt{bb+xx}$, and
 $\sqrt{pp+cc+2px+xx} = r + \sqrt{bb+xx}$, which squared
 is $cc+pp+2px+xx = rr+bb+xx+2r \times \sqrt{bb+xx}$
 Put $s = cc+pp-rr-bb$; then $s+2px = 2r \times$
 $\sqrt{bb+xx}$, and $ss+4spx+4ppxx = 4rrbb+4rrxx$;
 reduced, $4ppxx + 4spx = 4rrbb$.
 $\quad -4rr \qquad \qquad \qquad -ss$

Then x being found, it will be $EG(x) : \text{rad.}$
 $(1) :: BE(b) : \text{tang. BGE, or its supplement}$
 $BFD.$

P R O B. CCXXVII.

To find the greatest area contained under any number of
 right lines given, and another line unknown. 151²

Let ABCDE be the figure; then since ABE +
 BCDE is a maximum; it is evident, whatever the
 figure BCDE is, ABE must be a right-angled tri-
 angle, right-angled at B.

Again, since ABC + CDE + ACE is a maxi-
 mum; it is evident whatever ABC and CDE are,
 ACE must be a right-angled triangle, right-
 angled at C.

Also since ABCD + ADE is a maximum; it is
 plain, whatever the figure ABCD is, ADE must
 be a right-angled triangle, right-angled D. And
 so on if there were never so many lines. And
 therefore all the angles ABE, ACE, ADE, sub-
 tended by AE, must be right angles; and conse-
 quently the whole figure is inscribed in a semi-
 circle, whose diameter is AE, so that the whole
 may be a maximum.

Therefore if it be required to find the area,
 we must find the diameter AE, and then find
 the

Fig. the area of the polygon ABCDE inscribed in a femicircle.

P R O B. CCXXVIII.

152. To find a line, which with three given lines, will contain the greatest area possible.

It is plain the line sought is the diameter of the femicircle in which the three given lines are inscribed.

Let ABCD, be the quadrangle, draw the diagonals AC, BD, on which let fall the perpendiculars CP, BF.

Let $AB=b$, $BC=c$, $CD=d$, diameter $AD=y$. Then $BD=\sqrt{yy-bb}$, and $AC=\sqrt{yy-dd}$. But (Geom. IV. 28.) $CP=\frac{cd}{y}$, and $BF=\frac{bc}{y}$. There-

fore $b + \frac{cd}{y} \sqrt{yy-bb} = 2 \text{ area } ABCD$, and

$d + \frac{bc}{y} \sqrt{yy-dd} = 2 \text{ area} = b + \frac{cd}{y} \sqrt{yy-bb}$,

and $by+cd \times \sqrt{yy-bb} = dy+bc \sqrt{yy-dd}$, and squaring and multiplying, $bb^2y^4 + 2bcdy^3 + ccddy^2 - b^2yy - 2b^2cdy - bbccdd = ddy^4 + 2bcdy^3 + bbccyy - d^2yy - 2bcd^2y - bbccdd$. And reducing

$$\begin{array}{r} bb^2y^3 - b^2y + 2bcd^2 = 0. \\ -dd \quad - bbcc \quad - 2b^2cd \\ \quad \quad + ccdd \\ \quad \quad + d^2 \end{array}$$

And dividing by $bb-dd$

$$\begin{array}{r} y^3 - bb y = 2bcd, \text{ and } y \text{ being known, the} \\ -cc \\ -dd \end{array}$$

area is known from the foregoing steps.

P R O B.

a b c d e f g h i
 A B C D E F X Y Z
 Fig. 139




Fig. 17



P R O B. CCXXIX.

SP is perpendicular to PM, and there is given SP, ^{153.}
SN; and drawing NL, so that the angle LDM
may be equal to SCP; to find CD, a maximum.

Draw NA perpendicular to CD, then CA=AD,
and CA is a maximum. Put SN=b, SP=d,
SC=y, then CN=b-y, CP= $\sqrt{yy-dd}$. Then
by similar triangles, $y : \sqrt{yy-dd} : b-y : CA =$
 $\frac{b-y}{y} \sqrt{yy-dd} = \text{max.}$ and $\overline{b-y} \times \frac{yy-dd}{yy} =$
 $\text{max.} = \overline{b-y}^2 - \frac{dd}{yy} \times \overline{b-y}^2$. Increase y by a
very small quantity e, then $\overline{b-y-e} = \overline{b-y} - 2e \times$
 $\overline{b-y}$. Also $y+e = yy+2ye$, and by division
 $\frac{dd}{yy+2ye} = \frac{dd}{yy} - \frac{2dde}{y^3}$. Whence $\overline{b-y}^2 - \frac{dd}{yy} \times$
 $\overline{b-y}^2 = \overline{b-y}^2 - 2e \times \overline{b-y} - \frac{dd}{yy} + \frac{2dde}{y^3} \times$
 $\overline{b-y}^2 - 2e \times \overline{b-y}$, and transposing, $2e \times \overline{b-y}$
 $= \frac{2dde}{y^3} \times \overline{b-y} + \frac{2dde}{yy} \times \overline{b-y}$, and dividing
by $2e \times \overline{b-y}$, $1 = \frac{dd}{y^3} \times \overline{b-y} + \frac{dd}{yy}$, and mul-
tiplying by y^3 , we have $y^3 = ddb - ddy + ddy$, or
 $y^3 = bdd$, and $y = \sqrt[3]{bdd}$.

PROB. CCXXX.

154. Given the situation of the two places A, E, and the river BD; and suppose a traveller going from A to C, can travel 6 miles an hour on this side the river from A to C; and 9 miles an hour on the other side from C to E; it is required to know where he must cross the river BD, so that he may go from A to E in the least time possible.

Let AB, ED be perpendicular to BD; let $AB=a, DE=b, BD=d, m=6, n=9, BC=x$. Then

$$CD=d-x, AC=\sqrt{aa+xx}, CE=\sqrt{bb+d-x}^2$$

And per quest. $m : 1 :: \sqrt{aa+xx} : \sqrt{aa+xx}$

$$= \text{time in AC, and } n : 1 :: \sqrt{bb+d-x}^2 : \sqrt{bb+d-x}^2 = \text{time in CE. Therefore}$$

$$\frac{\sqrt{aa+xx}}{m} + \frac{\sqrt{bb+d-x}^2}{n} = \text{minimum. Or}$$

$$n\sqrt{aa+xx} + m\sqrt{bb+d-x}^2 = \text{min. Write } x+e$$

for x ; then $xx=xx+2xe$, and $\frac{d-x-e}{d-x} =$

$\frac{d-x}{d-x} - 2e \times \frac{1}{d-x}$. Therefore we have

$$n\sqrt{aa+xx+2xe} + m\sqrt{bb+d-x}^2 - 2e \times \frac{d-x}{d-x}$$

$$= n\sqrt{aa+xx} + m\sqrt{bb+d-x}^2. \text{ But}$$

$$\sqrt{aa+xx+2xe} = \sqrt{aa+xx} + \frac{xe}{\sqrt{aa+xx}}, \text{ and}$$

$$\sqrt{bb+d-x}^2 - 2e \times \frac{d-x}{d-x} = \sqrt{bb+d-x}^2 -$$

$$\frac{exd-x}{\sqrt{bb+d-x^2}}. \text{ Therefore } n\sqrt{aa+xx} + \frac{nx}{\sqrt{aa+xx}} \quad 154.$$

$$+ m\sqrt{bb+d-x^2} - \frac{mexd-x}{\sqrt{bb+d-x^2}} = n\sqrt{aa+xx}$$

$$+ m\sqrt{bb+d-x^2}. \text{ Therefore } \frac{nx}{\sqrt{aa+xx}} =$$

$$\frac{mexd-x}{\sqrt{bb+d-x^2}} = 0. \text{ And multiplying,}$$

$nx\sqrt{bb+dd-2dx+xx} = md-mx\sqrt{aa+xx}$; and squaring $nbbxx+nnddxx-2nndxx^2+nnx^4=mmddaax-2mmdaax+mmaaxx+mmddxx-2mmdxx^2+mmx^4$. And being reduced is,

$$\begin{aligned} nnx^4 - 2nndxx^2 + nnddxx + 2mmdaax - m^2d^2a^2 &= 0. \\ -mm + 2mmd + nbb & \\ -mdd & \\ -mmaa & \end{aligned}$$

P R O B. CCXXXI.

Within the given angle ACB, to cut off a given area, 155. with the shortest line AB.

Let the area = b ; s , c = sine and cof. C; $CA = x$, $CB = y$, then per quest. $sxy = b$, and by Trigonometry $AB = \sqrt{xx+yy-2cxy} = \text{min.}$; therefore $xx+yy-2cxy = \text{min.}$ but $xy = \frac{b}{s}$, and $y = \frac{b}{sx}$, therefore $xx + \frac{bb}{ssxx} - \frac{2cb}{s} = \text{min.}$ or $xx + \frac{bb}{ssxx} = \text{min.}$ Put $x+e$ for x , then $x+e = x^2+2ex$, and $\frac{1}{x+e}$ or $\frac{1}{x+e} - 2 = \frac{1}{xx} - \frac{2e}{x^3}$. Whence

Fig. 155. $xx + 2ex + \frac{bb}{ss} \times \frac{1}{xx} - \frac{bb}{ss} \times \frac{2e}{x^3} = xx + \frac{bb}{ssxx}$, and

$$2ex - \frac{2bbe}{ssx^3} = 0, \text{ and } x = \frac{bb}{ssx^3}, \text{ whence } x^4 = \frac{bb}{ss}.$$

But $y^4 = \frac{b^4}{s^4x^4} = \frac{b^4}{s^4} \times \frac{ss}{bb} = \frac{bb}{ss}$; therefore

$$y^4 = x^4, \text{ and } y = x = \sqrt{\frac{b}{s}}.$$

P R O B. CCXXXII.

156. To find the greatest parallelogram inscribed in a triangle.

Let the parallelogram BDEF be inscribed in the triangle ABC. Put $AB = a$, $BC = b$, $DB = x$, $DE = y$. Then by the similar triangles ABC, ADE, $a - x : y :: a : b$, and $ay = ba - bx$, and $y = \frac{ba - bx}{a} = b - \frac{bx}{a}$. But $xy = \max$, or $bx - \frac{bxx}{a} = \max$.

Put e for the small increment of x , then the increment of bx is be , and the decrement of xx is $x + e^2 - xx = 2xe$, and the decrement of $\frac{bxx}{a} = \frac{2bxe}{a}$, whence $be = \frac{2bxe}{a}$, and $1 = \frac{2x}{a}$, and $x = \frac{1}{2}a$. Therefore $y = \frac{1}{2}b$.

P R O B. CCXXXIII.

157. Given the point P within the right angle ACB; to draw the line APB, so that $AP \times PB$ may be a minimum.

Draw DP, PF parallel to CB, CA; and put $CF = b$, $CD = c$, $AD = x$. Then by similar triangles

angles $x : b :: c : y$, and $xy = bc$, and $y = \frac{bc}{x}$. Fig. 157.

Then $AP = \sqrt{bb + xx}$, and $PB = \sqrt{cc + yy}$; and

$AP \times PB = \sqrt{bb + xx} \times \sqrt{cc + \frac{bbcc}{xx}} = \text{min.}$ and

squaring, $bbcc + \frac{b^4cc}{xx} + ccxx + bbcc = \text{min.}$ and

$ccxx + \frac{b^4cc}{xx} = \text{min.}$ Whence $cc \times \overline{xx + e^2} +$

$\frac{b^4cc}{x+e} = ccxx + \frac{b^4cc}{xx}$, or $cc \times \overline{xx + 2xe} + b^4cc \times$

$\frac{1}{xx} - \frac{2e}{x^3} = ccxx + \frac{b^4cc}{xx}$. Whence $2ccxe - \frac{2b^4cce}{x^3}$

$= 0$, $x^3 = \frac{b^4}{cc}$, or $x^4 = b^4$, and $x = b$, whence $y = c$.

And $AC = x + c = b + c$, and $CB = b + y = b + c$.
Therefore $AC = CB$.

And if it be required to have $AP + PB$, a minimum, we shall have $\sqrt{bb + xx} + \sqrt{cc + yy} =$

min. or $\sqrt{bb + xx} + \sqrt{cc + \frac{bbcc}{xx}} = \text{min.}$ But

$\sqrt{bb + x + e} - \sqrt{bb + xx} = \frac{xe}{\sqrt{bb + xx}}$ = the

increment of $\sqrt{bb + xx}$. And in like manner

$\frac{-bbce}{xx \sqrt{cc + \frac{bbcc}{xx}}}$ is the increment of $\sqrt{cc + \frac{bbcc}{xx}}$, or

$\frac{bbce}{xx \sqrt{xx + bb}}$ its decrement. Therefore

$\frac{xe}{\sqrt{bb + xx}} = \frac{bbce}{xx \sqrt{bb + xx}}$, and $x^3 = bbc$, or

$x = \sqrt[3]{bbc}$, as in Prob. ccxiii. by another method.

P R O B. CCXXXIV.

Given the sum of the legs of a right-angled triangle; to find the legs, so as to contain the greatest area possible.

Let $a =$ sum of the legs, $x =$ one of them; then $x(a-x) = 2 \text{ area} = \text{max.}$ therefore $x + e \times a - x - e = x \times a - x$, that is, $ax - xx - xe + ae - xe = ax - xx$, and $ae - 2xe = 0$, or $2x = a$, whence $x = \frac{1}{2}a$, and $a - x = \frac{1}{2}a$. Therefore the legs are equal. And therefore when the area is given; the sum of the legs will be the least, when they are equal.

P R O B. CCXXXV.

Given the area of a right-angled triangle; to find the sides, when the perimeter is the least possible.

Let $a =$ area, $x =$ sum of the legs, $v, y =$ the two legs; then $vv + yy + 2vy = xx$, but $vy = 2a$, and $vv + yy = xx - 2vy = xx - 4a$, and the hypotenuse $= \sqrt{vv + yy} = \sqrt{xx - 4a}$; therefore $x + \sqrt{xx - 4a} =$ perimeter $=$ min. write $x + e$ for x ; then $\sqrt{x+e}^2 - 4a = \sqrt{xx + 2xe - 4a} = \sqrt{xx - 4a} + \frac{xe}{\sqrt{xx - 4a}}$; whence $x + e + \sqrt{xx - 4a} + \frac{xe}{\sqrt{xx - 4a}} = x + \sqrt{xx - 4a}$, and $e + \frac{xe}{\sqrt{xx - 4a}} = 0$. And $e\sqrt{xx - 4a} = -xe$, and $eex - 4aee = -xxe$, and $4aee = 0$, and $ee = \frac{0}{4a}$, or $e = 0$. And therefore since the increment of x is nothing; therefore x is a minimum, and when x or the sum

sum is a minimum; then the legs are equal, by Fig. the last problem; therefore $v=y=\frac{1}{2}x$, and $\frac{1}{2}xx=2a$, or $x=\sqrt{8a}$.

P R O B. CCXXXVI.

Given the solidity of a square pyramid DF; to find 158. the slant side AB the least possible.

Let b =solidity, $x=CB$ the height, $y=2AC$ the breadth, then $AB=\sqrt{xx+\frac{1}{4}yy}$. But $\frac{1}{3}xyy=b$,

and $yy=\frac{3b}{x}$; therefore $AB=\sqrt{xx+\frac{3b}{4x}}=$

minimum, and $xx+\frac{3b}{4x}=\text{min.}$ Put $x+e$ for

x , then $xx=xx+2xe$, and $\frac{3b}{4x}=\frac{3b}{4x+4e}=\frac{3b}{4x}$

$-\frac{3be}{4xx}$; whence $xx+2xe+\frac{3b}{4x}\rightarrow\frac{3be}{4xx}=\frac{3b}{4x}$,

and $2xe-\frac{3be}{4xx}=0$, and $8x^3=3b$, whence

$$x=\sqrt[3]{\frac{3b}{8}}=\frac{1}{2}\sqrt[3]{3b}.$$

P R O B. CCXXXVII.

Given the solidity of the square pyramid DF, to find 158. that which has the least surface, excluding the base.

Let b =solidity, $x=CB$ the height, $y=2AC$, or $2AD$ the breadth: Then $AB=\sqrt{xx+\frac{1}{4}yy}$,

and $\frac{1}{3}y \times \sqrt{xx+\frac{1}{4}yy}=DBL$, and $2y\sqrt{xx+\frac{1}{4}yy}=\text{surface}$.

But $\frac{1}{3}xyy=b$, and $yy=\frac{3b}{x}$. Whence the

$$\text{surface} = 2\sqrt{\frac{3b}{x}} \times \sqrt{xx+\frac{3b}{4x}} = 2\sqrt{3bx+\frac{9bb}{4xx}}$$

Fig. 158. = maximum. And $3bx + \frac{9bb}{4xx} = \text{maximum}$, or

$$12bx + \frac{9bb}{xx} = \text{max. write } x+e \text{ for } x, \text{ then } \frac{9bb}{xx}$$

$$= \frac{9bb}{xx+2xe} = \frac{9bb}{xx} - \frac{18bbe}{x^3}. \text{ Therefore } 12bx +$$

$$12be + \frac{9bb}{xx} - \frac{18bbe}{x^3} = 12bx + \frac{9bb}{xx}; \text{ and } 12be -$$

$$\frac{18bbe}{x^3} = 0, \text{ or } 12b = \frac{18bb}{x^3}, \text{ and } 12x^3 = 18b, \text{ and}$$

$$x^3 = \frac{3b}{2}, \text{ whence } x = \sqrt[3]{\frac{3}{2}b}.$$

P R O B. CCXXXVIII.

159. To find the greatest cylinder, inscribed in a given cone.

Let axis $AB=a$, BC or $BF=b$, $c=3.1416$, $DB=x$, DE or $DG=y$. Then by the similar triangles ABC , ADE , $a-x : y :: a : b$, and $ay = ba - bx$, or $bx = ab - ay$,

and $x = \frac{ab-ay}{b}$. But $cyy = \text{maximum}$, or

$$cyy \times \frac{ab-ay}{b} = \frac{abcyy - acy^2}{b} = \text{max. that is } \frac{ac}{b} \times$$

$b\overline{yy} - y^2 = \text{max. and } b\overline{yy} - y^2 = \text{max. put } y+e \text{ for } y, \text{ then } b\overline{yy} \text{ becomes } b\overline{yy} + 2b\overline{ye}, \text{ and } y^2 \text{ becomes } y^2 + 3y^2e. \text{ Whence } b\overline{yy} + 2b\overline{ye} - y^2 - 3y^2e = b\overline{yy} - y^2. \text{ And } 2b\overline{ye} - 3y^2e = 0, \text{ or } 2by = 3yy, \text{ and } y = \frac{2}{3}b.$

$$\text{Whence } x = \frac{ab - \frac{2}{3}ab}{b} = \frac{1}{3}a.$$

P R O B. CCXXXIX.

Given the weights of two elastic bodies A, C; to find the weight of the intermediate body B; so that A striking B at rest, and B with the motion acquired, striking C at rest, may make C's motion the greatest possible.

Let x = weight of B, a = velocity of A.
 y = velocity of B, acquired by the stroke.
 v = velocity of C, by the stroke.

Then Aa is the motion of both A and B, after the stroke, as well as before; and a is the difference of their velocities; therefore $y - a$ is the velocity of A after the stroke. And since the sum of their motions remains the same (Mechan. 10.), therefore $xy + y - a \times A = Aa$, or $xy - Aa + Ay = Aa$, that is, $xy + Ay = 2aA$, and $y = \frac{2aA}{A+x}$

Again, xy is the motion of both B and C, and y the difference of their velocities, as well after as before the stroke of B. Therefore $v - y$ is the velocity of B after its striking C. Whence $Cv + v - y \times x = xy$, or $Cv + xv = 2yx$. Whence $v = \frac{2yx}{C+x} = \frac{4aAx}{A+x \times C+x} =$ maximum per question; or $\frac{x}{A+x \times C+x} =$ maximum; or $\frac{A+x \times C+x}{x} =$ minimum, that is, $\frac{AC + A + C.x + xx}{x} =$ minimum, or $\frac{AC}{x} + A + C$

Fig.

$A+C+x$ = minimum; therefore $\frac{AC}{x} + x$ = minimum; put $x+e$ for x ; then $\frac{AC}{x+e} = \frac{AC}{x} - \frac{ACe}{xx}$, and therefore $\frac{AC}{x} - \frac{ACe}{xx} + x + e = \frac{AC}{x} + x$, and throwing out the superfluous quantities, $e - \frac{ACe}{xx} = 0$, and $xx = AC$, whence $x = \sqrt{AC}$.

P R O B. CCXL:

To find $x^m - x^n$ the greatest possible, supposing n greater than m .

Write $x+e$ for x , then $\overline{x+e}^m = x^m + m^{m-1}e$, and $\overline{x+e}^n = x^n + nx^{n-1}e$; therefore $\overline{x+e}^m - \overline{x+e}^n = x^m - x^n$; that is, $x^m + mx^{m-1}e - x^n - nx^{n-1}e = x^m - x^n$. And by subtraction, $mx^{m-1}e - nx^{n-1}e = 0$, or $mx^{m-1} = nx^{n-1}$, or $mx^m = nx^n$. And n being greater than m , we have $nx^{n-m} = m$, and $x^{n-m} = \frac{m}{n}$; whence

$$x = \sqrt[n-m]{\frac{m}{n}}$$

P R O B.

P R O B. CCXLI.

To find the greatest parallelogram inscribed in the given curve AMC.

Let MPBF be the greatest parallelogram. To the point M where it touches the curve, draw the tangent TMD. Then if the subtangent PT be equal to the height of the parallelogram PB, then MPBF is the greatest parallelogram. For it is plain from Problem ccxxxii, that this parallelogram is the greatest that can be inscribed in the triangle TDB; and as this is greater than any other that can be inscribed in the triangle, so much more, is it greater than any other that can be inscribed in the curve, since the angle M which is in the curve, will in all other cases fall short of the tangent.

Therefore knowing the method of drawing a tangent to the curve; you must seek the point P, where the ordinate PM being erected, and the tangent TM drawn, TP may be equal to PB. Thus if AM be a parabola; put $AB=a$, $AP=x$, then by the nature of the curve, $AF=x$, whence $TP=2x$, $PB=a-x$, therefore $2x=a-x$, $3x=a$, or $x=\frac{1}{3}a$.

And the same will hold good, if not in all, yet in most curves which are convex to the axis. For since the parallelogram is the greatest for the triangle, it will also be greatest for the curve, since the curve at that place coincides with the tangent.

Otherwise thus,

Suppose the nature of the curve be $rx^m = y^n$, where $AP=x$, $PM=y$, also $AB=a$, $BC=b$.

Then

Fig. Then $PB = a - x$, and $\overline{a-x} \times y = \max$. But

160.

$y = r^{\frac{1}{n}} x^{\frac{m}{n}}$, therefore $\overline{a-x} \times r^{\frac{1}{n}} x^{\frac{m}{n}} = \max$. or

$a x^{\frac{m}{n}} - x^{\frac{m+n}{n}} = \max$. put $x + e = x$; then

$$a \cdot \overline{x+e}^{\frac{m}{n}} = a x^{\frac{m}{n}} + \frac{m}{n} a x^{\frac{m-n}{n}} e, \text{ and } \overline{x+e}^{\frac{m+n}{n}}$$

$$= x^{\frac{m+n}{n}} + \frac{m+n}{n} x^{\frac{m}{n}} e. \text{ Therefore,}$$

$$a x^{\frac{m}{n}} + \frac{m}{n} a x^{\frac{m-n}{n}} e - x^{\frac{m+n}{n}} - \frac{m+n}{n} x^{\frac{m}{n}} e = a x^{\frac{m}{n}} - x^{\frac{m+n}{n}};$$

$$\text{whence } \frac{m}{n} a x^{\frac{m-n}{n}} e - \frac{m+n}{n} x^{\frac{m}{n}} e = 0, \text{ or}$$

$$\overline{m a x^{\frac{m-n}{n}}} = \overline{m+n} \times \overline{x^{\frac{m}{n}}}; \text{ and dividing by}$$

$$x^{\frac{m-n}{n}}, \overline{m a} = \overline{m+n} \times x = \overline{m+n} \times x, \text{ whence}$$

$$x = \frac{m a}{m+n}. \text{ Which is general for all parabolical}$$

figures. Thus if $m=1, n=2$, as in the common parabola, then $x = \frac{1}{3} a$, and if $m=2, n=1$, then is $x = \frac{2}{3} a$, as in the same parabola, with its convexity towards the axis. If $m=1, n=1$; then $x = \frac{1}{2} a$, for the triangle, as was proved before.

P R O B. CCXLII.

161. Given the distance of the point A from the perpendicular plane BC; to find the position of the plane AC, through which a body shall descend in the shortest time possible to the plane BC.

Let AB be perpendicular to BC, AD parallel to it, and CD perpendicular to AC. Put $AB = b$, $BC = x$; then (Mechan. 34. cor. 1.) in the time a body

body descends through the inclined plane AC, and Fig-
body will fall perpendicularly through the space 161-
AD. Therefore as the time in AC must be a

minimum, the time in AD must be a minimum,
and AD itself must be a minimum. By the si-
milar triangles BAC, CAD, it is $BC (x) :$
 $CA (\sqrt{bb+xx}) :: CA (\sqrt{bb+xx}) : AD =$
 $\frac{bb+xx}{x} = \text{minimum.}$ And $\frac{bb}{x} + x = \text{min. write } x + e$

for x , then $\frac{bb}{x+e} = \frac{bb}{x} - \frac{bbe}{xx}$; therefore $\frac{bb}{x}$

$-\frac{bbe}{xx} + x + e = \frac{bb}{x} + x$, and $-\frac{bbe}{xx} + e = 0$, or

$1 = \frac{bbe}{xx}$, and $xx = bb$, or $x = b$, therefore
 $BC = BA$.

Otherwise thus,

Describe the circle AGC with the center B, and 162-
radius BA; draw AC and any other line AE,
and CGF parallel to it. Then (Mechan. 37.
cor. 1.) the times of a body's descending through
GC, AC, are equal. And the times of descend-
ing through the equal lines, of equal inclinations,
AE, FC are equal. But the time of descending
through GC is less than the time of descending
through FC. Therefore the time of descending
through AC is less than the time of descending
through any other line AE.

P R O B. CCXLIII.

AB is a horizontal line, BD an inclined plane. It is 163-
required to find the position of the plane AD, through
which a body descending from A shall arrive at the
plane BD, in the least time possible.


Suppose AD to be the plane, draw AL per-
pendicular to AB, and DH perpendicular to AL,
and


Fig. and DF perpendicular to AD. And put $b=AB$,
 163. $s, c = \text{fine and cof. B}$, $BD=x$, $AD=y$. Then
 by plain Trigonometry $\sqrt{bb+xx-2cbx}=y$; and
 $AD (y) : S.B (s) :: BD (x) : \frac{sx}{y} = S.BAD$
 or ADH. And rad. (1) : $AD (y) :: S. ADH$
 $(\frac{sx}{y}) : AH=sx$. And by similar triangles, AH
 $(sx) : AD (y) :: AD (y) : AF = \frac{yy}{sx} =$
 $\frac{bb+xx-2cbx}{sx}$. But the time of falling through
 AF is equal to the time of descending through
 AD (Mech. 34. cor. 1.). And this time is a mi-
 nimum, therefore AF is a minimum, that is,
 $\frac{bb+xx-2cbx}{sx} = \text{min.}$ and $\frac{bb+xx-2cbx}{x} = \text{min.}$
 or $\frac{bb}{x} + x - 2cb = \text{min.}$ whence $\frac{bb}{x} + x = \text{min.}$
 as in the last problem. And therefore $x=b$, or
 $AD=AB$.

Or thus,

On AF describe a semicircle ADF to touch the
 line AL in D; draw AD, which will be the line
 of shortest time. For the time of descending
 through all the cords in the semicircle will be
 equal (Mechan. 37. cor. 1.) to the time in AD.
 But the time in any chord is shorter than the time
 in the same chord when produced to the line BL,
 which lies without the circle. And therefore the
 time in AD is also shorter than in any other line
 drawn to BL.

PROB.





The first part of the book is devoted to the study of the properties of the circle and the ellipse. The author discusses the various ways in which these curves can be constructed and the relationships between their different parts. He also examines the properties of the tangents and normals to these curves, and the way in which they can be used to solve a variety of geometric problems.

The second part of the book is devoted to the study of the properties of the parabola and the hyperbola. The author discusses the various ways in which these curves can be constructed and the relationships between their different parts. He also examines the properties of the tangents and normals to these curves, and the way in which they can be used to solve a variety of geometric problems.

The third part of the book is devoted to the study of the properties of the sphere and the cylinder. The author discusses the various ways in which these solids can be constructed and the relationships between their different parts. He also examines the properties of the tangents and normals to these solids, and the way in which they can be used to solve a variety of geometric problems.

The fourth part of the book is devoted to the study of the properties of the cone and the frustum. The author discusses the various ways in which these solids can be constructed and the relationships between their different parts. He also examines the properties of the tangents and normals to these solids, and the way in which they can be used to solve a variety of geometric problems.

The fifth part of the book is devoted to the study of the properties of the sphere and the cylinder. The author discusses the various ways in which these solids can be constructed and the relationships between their different parts. He also examines the properties of the tangents and normals to these solids, and the way in which they can be used to solve a variety of geometric problems.

P R O B. CCXLIV.

To divide a given line AB, into three parts, x, y, z ; 164.
so that xyz^3 may be the greatest product possible.

First, suppose $x+y=b$ a given quantity, to find $xyy = a$ maximum. Then $x=b-y$, and $\overline{b-y}xyy$ or $\overline{byy-y^3} = \max.$ put $y+e$ for y , then $b \times \overline{y+e} - \overline{y+e} = \max.$ that is, $b \overline{yy} + 2bye - y^3 - 3yye = b \overline{yy} - y^3$, and $2bye - 3yye = 0$, and $3y = 2b$, or $y = \frac{2}{3}b$, and $x = \frac{1}{3}b$. Therefore $y = 2x$.

Again, let $x+y+z=d$, to find $xy^2z^3 = \max.$ Then by what is gone before, whatever z be, y will be $= 2x$. Whence $4x^3z^3 = \max.$ But $x+y$

$$= d - z, \text{ or } 3x = d - z, \text{ and } 4x^3z^3 = \frac{4}{3} \times \overline{d-z}^3$$

$\times z^3 = \max.$ and $\overline{d-z}^3 \times z^3 = \max.$ or $\overline{d-z} \times z = \max.$ or $\overline{dz-zz} = \max.$ put $z+e$ for z , then $\overline{dz+de-z-zz} = \overline{dz-zz}$, or $\overline{dz+de-zz-2ze} = \overline{dz-zz}$, or $\overline{de-2ze} = 0$; whence $2z = d = x+y+z$, and $z = x+y = 3x$. Therefore $x+2x+3x$ or $6x = d$, and $x = \frac{1}{6}d$, and $y = (2x) = \frac{2}{6}d$, and $z = (3x) = \frac{3}{6}d$.

F I N I S.

166644D
37
13

M I N I M A

P R O B C O M M I

E R R A T A .

Page 328. line 8. (from the bottom) read $a = \frac{3^x}{5}$.

p. 336. line 8. read $\frac{bfa - bdc}{}$

p. 430. line 12. read $\sqrt{m \times c + y} -$

p. 466. line 3. read $n \times \sqrt{2} - 1.$

M I N I M A

